

ON SEMI-INVARIANT SUBMANIFOLDS OF A NEARLY KENMOTSU MANIFOLD WITH A SEMI-SYMMETRIC NON-METRIC CONNECTION

MOBIN AHMAD* AND JAE-BOK JUN**

ABSTRACT. We define a semi-symmetric non-metric connection in a nearly Kenmotsu manifold and we study semi-invariant submanifolds of a nearly Kenmotsu manifold endowed with a semi-symmetric non-metric connection. Moreover, we discuss the integrability of distributions on semi-invariant submanifolds of a nearly Kenmotsu manifold with a semi-symmetric non-metric connection.

1. Introduction

In [4], K. Kenmotsu introduced and studied a new class of almost contact manifolds called Kenmotsu manifolds. The notion of nearly Kenmotsu manifold was introduced by A. Shukla in [8]. Semi-invariant submanifolds in Kenmotsu manifolds were studied by N. Papaghuic [6], M. Kobayashi [5] and B. B. Sinha and R. N. Yadav [9]. Semi-invariant submanifolds of a nearly Kenmotsu manifolds were studied by M. M. Tripathi and S. S. Shukla in [10]. In this paper we study semi-invariant submanifolds of a nearly Kenmotsu manifold with a semi-symmetric non-metric connection.

Let ∇ be a linear connection in an n -dimensional differentiable manifold M . The torsion tensor T and the curvature tensor R of ∇ are given respectively by

$$\begin{aligned}T(X, Y) &= \nabla_X Y - \nabla_Y X - [X, Y], \\R(X, Y)Z &= \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z.\end{aligned}$$

Received February 01, 2010; Accepted April 23, 2010.

2010 Mathematics Subject Classification: Primary 53C05, 53D12.

Key words and phrases: semi-invariant submanifolds, nearly Kenmotsu manifolds, semi-symmetric non-metric connection, Gauss and Weingarten equations, integrability conditions, distributions.

Correspondence should be addressed to Jae-Bok Jun, jbjun@kookmin.ac.kr.

The connection ∇ is *symmetric* if the torsion tensor T vanishes, otherwise it is non-symmetric. The connection ∇ is *metric* if there is a Riemannian metric g in M such that $\nabla g = 0$, otherwise it is non-metric. It is well known that a linear connection is symmetric and metric if it is the Levi-Civita connection.

In ([3], [7]) A. Friedmann and J. A. Schouten introduced the idea of a semi-symmetric linear connection. A linear connection ∇ is said to be *semi-symmetric* if its torsion tensor T is of the form

$$T(X, Y) = \eta(Y)X - \eta(X)Y,$$

where η is a 1-form.

The paper is organized as follows. In section 2, we give a brief introduction of nearly Kenmotsu manifold. In section 3, we show that the induced connection on a semi-invariant submanifolds of a nearly Kenmotsu manifold with a semi-symmetric non-metric connection is also semi-symmetric and non-metric. In section 4, we established some lemmas on semi-invariant submanifolds and in section 5, we discussed the integrability conditions of the distributions on semi-invariant submanifolds of nearly Kenmotsu manifolds with a semi-symmetric non-metric connection.

2. Preliminaries

Let \bar{M} be an $(2m + 1)$ -dimensional almost contact metric manifold [2] with a metric tensor g , a tensor field ϕ of type $(1,1)$, a vector field ξ and a 1-form η which satisfies

$$\phi^2 = -I + \eta \otimes \xi, \phi\xi = 0, \eta\phi = 0, \eta(\xi) = 1,$$

$$g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y)$$

for any vector fields X and Y on \bar{M} . If in addition to the condition for an almost contact metric structure we have $d\eta(X, Y) = g(X, \phi Y)$, then the structure is said to be a *contact metric structure*.

The almost contact metric manifold \bar{M} is called a *nearly Kenmotsu manifold* if it satisfies the condition [8]

$$(\bar{\nabla}_X \phi)Y + (\bar{\nabla}_Y \phi)X = -\eta(Y)\phi X - \eta(X)\phi Y,$$

where $\bar{\nabla}$ denotes the Riemannian connection with respect to g . If, moreover, \bar{M} satisfies

$$(2.1) \quad (\bar{\nabla}_X \phi)Y = g(\phi X, Y)\xi - \eta(Y)\phi X,$$

then it is called *Kenmotsu manifold* [3].

DEFINITION 2.1. An n -dimensional Riemannian submanifold M of a nearly Kenmotsu manifold \bar{M} is called a *semi-invariant submanifold* if ξ is tangent to M and there exists on M a pair of orthogonal distributions (D, D^\perp) such that [1]

- (i) $TM = D \oplus D^\perp \oplus \{\xi\}$,
- (ii) distribution D is invariant under ϕ , that is $\phi D_x = D_x$ for all $x \in M$,
- (iii) distribution D^\perp is anti-invariant under ϕ , that is $\phi D_x^\perp \subset T_x^\perp M$ for all $x \in M$, where $T_x M$ and $T_x^\perp M$ are respectively the tangent and normal space of M at x .

The distribution D (resp. D^\perp) is called the *horizontal* (resp. *vertical*) distribution. A semi-invariant submanifold M is said to be an *invariant* (resp. *anti-invariant*) submanifold if we have $D_x^\perp = \{0\}$ (resp. $D_x = \{0\}$) for each $x \in M$. We also call M is *proper* if neither D nor D^\perp is null. It is easy to check that each hypersurface of M which is tangent to ξ inherits a structure of the semi-invariant submanifold of \bar{M} .

Now, we define a *semi-symmetric non-metric connection* $\bar{\nabla}$ in a Kenmotsu manifold by

$$(2.2) \quad \bar{\nabla}_X Y = \bar{\bar{\nabla}}_X Y + \eta(Y)X$$

such that $(\bar{\nabla}_X g)(Y, Z) = -\eta(Y)g(X, Z) - \eta(Z)g(X, Y)$ for any X and $Y \in TM$, where $\bar{\bar{\nabla}}$ is the induced connection on M .

From (2.1) and (2.2), we have

$$(2.3) \quad (\bar{\nabla}_X \phi)Y = g(\phi X, Y)\xi - 2\eta(Y)\phi X,$$

$$(\bar{\nabla}_X \phi)Y + (\bar{\nabla}_Y \phi)X = -2\eta(X)\phi Y - 2\eta(Y)\phi X.$$

We denote by g the metric tensor of \bar{M} as well as that is induced on M . Let $\bar{\nabla}$ be the semi-symmetric non-metric connection on \bar{M} and ∇ be the induced connection on M with respect to the unit normal N .

THEOREM 2.2. *The connection induced on the semi-invariant submanifolds of a nearly Kenmotsu manifold with a semi-symmetric non-metric connection is also a semi-symmetric non-metric connection.*

Proof. Let ∇ be the induced connection with respect to the unit normal N on semi-invariant submanifolds of a nearly Kenmotsu manifold with semi-symmetric non-metric connection $\bar{\nabla}$. Then

$$\bar{\nabla}_X Y = \nabla_X Y + m(X, Y),$$

where m is a tensor field of type $(0, 2)$ on semi-invariant submanifold M . If ∇^* be the induced connection on semi-invariant submanifolds from Riemannian connection $\bar{\nabla}$, then

$$\bar{\nabla}_X Y = \nabla^*_X Y + h(X, Y),$$

where h is a second fundamental tensor. Now using (2.2), we have

$$\nabla_X Y + m(X, Y) = \nabla^*_X Y + h(X, Y) + \eta(Y)X.$$

Equating the tangential and normal components from the both sides in the above equation, we get

$$h(X, Y) = m(X, Y)$$

and

$$\nabla_X Y = \nabla^*_X Y + \eta(Y)X.$$

Thus ∇ is also a semi-symmetric non-metric connection. □

Now, the Gauss formula for a semi-invariant submanifolds of a nearly Kenmotsu manifold with a semi-symmetric non-metric connection is

$$(2.4) \quad \bar{\nabla}_X Y = \nabla_X Y + h(X, Y)$$

and the Weingarten formula for M is given by

$$(2.5) \quad \bar{\nabla}_X N = -(A_N + a)X + \nabla^\perp_X N$$

for $X, Y \in TM, N \in T^\perp M$, where $a = \eta(N)$ is a function on M , h (resp. A_N) is the second fundamental tensor (resp. form) of M in \bar{M} and ∇^\perp denotes the operator of the normal connection. Moreover, we have

$$(2.6) \quad g(h(X, Y), N) = g(A_N X, Y).$$

Any vector X tangent to M is given as

$$(2.7) \quad X = PX + QX + \eta(X)\xi,$$

where PX and QX belong to the distribution D and D^\perp respectively.

For any vector field N normal to M , we put

$$(2.8) \quad \phi N = BN + CN,$$

where BN (resp. CN) denotes the tangential (resp. normal) component of ϕN .

The Nijenhuis tensor $N(X, Y)$ for a semi-symmetric non-metric connection is defined as

$$(2.9) \quad N(X, Y) = (\bar{\nabla}_{\phi X} \phi)Y - (\bar{\nabla}_{\phi Y} \phi)X - \phi(\bar{\nabla}_X \phi)Y + \phi(\bar{\nabla}_Y \phi)X$$

for any X and $Y \in T\bar{M}$.

From (2.3), we have

$$(2.10) \quad (\bar{\nabla}_{\phi X}\phi)Y = 2\eta(Y)X - 2\eta(X)\eta(Y)\xi - (\bar{\nabla}_Y\phi)\phi X.$$

Also, we have

$$(2.11) \quad (\bar{\nabla}_Y\phi)(\phi X) = ((\bar{\nabla}_Y\eta)X)\xi + \eta(X)\bar{\nabla}_Y\xi - \phi(\bar{\nabla}_Y\phi)X.$$

Now, using (2.11) in (2.10), we have

$$(2.12) \quad (\bar{\nabla}_{\phi X}\phi)Y = 2\eta(Y)X - 2\eta(X)\eta(Y)\xi - ((\bar{\nabla}_Y\eta)X)\xi \\ - \eta(X)\bar{\nabla}_Y\xi + \phi(\bar{\nabla}_Y\phi)X.$$

By virtue of (2.12) and (2.9), we get

$$(2.13) \quad N(X, Y) = -2\eta(Y)X - 2\eta(X)Y + 8\eta(X)\eta(Y)\xi + \eta(Y)\bar{\nabla}_X\xi \\ - \eta(X)\bar{\nabla}_Y\xi + 2g(\phi X, Y)\xi + 4\phi(\bar{\nabla}_Y\phi X)$$

for any X and $Y \in T\bar{M}$.

3. Basic Lemmas

LEMMA 3.1. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric non-metric connection. Then we have*

$$2(\bar{\nabla}_X\phi)Y = \nabla_X\phi Y - \nabla_Y\phi X + h(X, \phi Y) - h(Y, \phi X) - \phi[X, Y].$$

Proof. By the Gauss formula we have

$$(3.1) \quad \bar{\nabla}_X\phi Y - \bar{\nabla}_Y\phi X = \nabla_X\phi Y - \nabla_Y\phi X + h(X, \phi Y) - h(Y, \phi X).$$

Also by use of (2.4), the covariant differentiation yields

$$(3.2) \quad \bar{\nabla}_X\phi Y - \bar{\nabla}_Y\phi X = (\bar{\nabla}_X\phi)Y - (\bar{\nabla}_Y\phi)X + \phi[X, Y].$$

From (3.1) and (3.2) we get

$$(3.3) \quad (\bar{\nabla}_X\phi)Y - (\bar{\nabla}_Y\phi)X = \nabla_X\phi Y - \nabla_Y\phi X + h(X, \phi Y) - h(Y, \phi X) - \phi[X, Y].$$

Using $\eta(X) = 0$ for each $X \in D$ in (2.3), we get

$$(3.4) \quad (\bar{\nabla}_X\phi)Y + (\bar{\nabla}_Y\phi)X = 0.$$

Adding (3.3) and (3.4) we get the result. □

Similar computations also yields the following.

LEMMA 3.2. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric non-metric connection. Then we have*

$$2(\bar{\nabla}_X\phi)Y = -A_{\phi Y}X + \nabla_X^\perp\phi Y - \nabla_Y\phi X - h(Y, \phi X) - \phi[X, Y]$$

for any $X \in D$ and $Y \in D^\perp$.

LEMMA 3.3. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric non-metric connection. Then we have*

$$(3.5) \quad Q\nabla_X(\phi PY) + Q\nabla_Y(\phi PX) - QA_{\phi QY}X - QA_{\phi QX}Y = -2\eta(Y)\phi QX \\ -2\eta(X)\phi QY + 2Bh(X, Y),$$

$$(3.6) \quad h(X, \phi PY) + h(Y, \phi PX) + \nabla_X^\perp\phi QY + \nabla_Y^\perp\phi QX = 2Ch(X, Y) \\ + \phi Q\nabla_XY + \phi Q\nabla_YX,$$

$$\eta(\nabla_X\phi PY + \nabla_Y\phi PX - A_{\phi QY}X - A_{\phi QX}Y) = 0$$

for all X and $Y \in TM$.

Proof. Differentiating (2.7) covariantly and using (2.4) and (2.5), we have

$$(3.7) \quad (\bar{\nabla}_X\phi)Y + \phi(\nabla_XY) + \phi h(X, Y) = P\nabla_X(\phi PY) + Q\nabla_X(\phi PY) \\ + \eta(\nabla_X\phi PY)\xi - PA_{\phi QY}X - QA_{\phi QY}X \\ - \eta(A_{\phi QY}X)\xi + \nabla_X^\perp\phi QY + h(X, \phi PY).$$

Similarly, we have

$$(3.8) \quad (\bar{\nabla}_Y\phi)X + \phi(\nabla_YX) + \phi h(Y, X) = P\nabla_Y(\phi PX) + Q\nabla_Y(\phi PX) \\ + \eta(\nabla_Y\phi PX)\xi - PA_{\phi QX}Y - QA_{\phi QX}Y \\ - \eta(A_{\phi QX}Y)\xi + \nabla_Y^\perp\phi QX + h(Y, \phi PX).$$

Adding (3.7) and (3.8) and using (2.3) and (2.8) we have

$$(3.9) \quad -2\eta(Y)\phi PX - 2\eta(Y)\phi QX - 2\eta(X)\phi PY - 2\eta(X)\phi QY + \phi P\nabla_XY \\ + \phi Q\nabla_XY + \phi P\nabla_YX + \phi Q\nabla_YX + 2Bh(Y, X) + 2Ch(Y, X) \\ = P\nabla_X(\phi PY) + P\nabla_Y(\phi PX) + Q\nabla_Y(\phi PX) - PA_{\phi QY}X \\ + Q\nabla_X(\phi PY) + \nabla_X^\perp\phi QY - PA_{\phi QX}Y - QA_{\phi QY}X \\ - QA_{\phi QX}Y + \nabla_Y^\perp\phi QX + h(Y, \phi PX) + h(X, \phi PY) \\ + \eta(\nabla_X\phi PY)\xi + \eta(\nabla_Y\phi PX)\xi - \eta(A_{\phi QX}Y)\xi - \eta(A_{\phi QY}X)\xi.$$

Equations from (3.1) to (3.4) follows the results by the comparison of the tangential, normal and vertical components of (3.9). \square

DEFINITION 3.4. *The horizontal distribution D is said to be parallel with respect to the connection ∇ on M if $\nabla_X Y \in D$ for all vector fields X and $Y \in D$.*

PROPOSITION 3.5. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric and non-metric connection. If the horizontal distribution D is parallel, then $h(X, \phi Y) = h(Y, \phi X)$ for all X and $Y \in D$.*

Proof. Since D is parallel, therefore, $\nabla_X \phi Y \in D$ and $\nabla_Y \phi X \in D$ for each X and $Y \in D$. Now from (3.5) and (3.6), we get

$$(3.10) \quad h(X, \phi Y) + h(Y, \phi X) = 2\phi h(X, Y).$$

Replacing X by ϕX in the above equation, we have

$$(3.11) \quad h(\phi X, \phi Y) - h(Y, X) = 2\phi h(\phi X, Y).$$

Replacing Y by ϕY in (3.10), we have

$$(3.12) \quad -h(X, Y) + h(\phi X, \phi Y) = 2\phi h(X, \phi Y).$$

Comparing (3.11) and (3.12), we have

$$h(X, \phi Y) = h(\phi X, Y)$$

for all X and $Y \in D$. □

DEFINITION 3.6. *A semi-invariant submanifold is said to be mixed totally geodesic if $h(X, Z) = 0$ for all $X \in D$ and $Z \in D^\perp$.*

LEMMA 3.7. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric non-metric connection. Then M is mixed totally geodesic if and only if $A_N X \in D$ for all $X \in D$.*

Proof. If $A_N X \in D$, then $g(h(X, Y), N) = g(A_N X, Y) = 0$, which gives $h(X, Y) = 0$ for $Y \in D^\perp$. Hence M is mixed totally geodesic. □

4. Integrability conditions for distributions

THEOREM 4.1. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric non-metric connection. Then the distribution $D \oplus \{\xi\}$ is integrable if the following conditions are satisfied*

$$S(X, Y) \in D \oplus \{\xi\},$$

and

$$h(X, \phi Y) = h(\phi X, Y)$$

for X and $Y \in D \oplus \{\xi\}$.

Proof. The torsion tensor $S(X, Y)$ of almost contact structure is given by

$$S(X, Y) = N(X, Y) + 2d\eta(X, Y)\xi,$$

where $N(X, Y)$ is the Nijenhuis tensor. Then we know

$$(4.1) \quad S(X, Y) = [\phi X, \phi Y] - \phi[\phi X, Y] - \phi[X, \phi Y] + 2d\eta(X, Y)\xi.$$

Suppose that $D \oplus \{\xi\}$ is integrable, so for X and $Y \in D \oplus \{\xi\}$, $N(X, Y) = 0$. Then $S(X, Y) = 2d\eta(X, Y)\xi \in D \oplus \{\xi\}$. Using the Gauss formula in (2.13), we get

$$(4.2) \quad N(X, Y) = 4\phi(\nabla_Y \phi X) + 4\phi h(Y, \phi X) - 4h(Y, X)$$

for all X and $Y \in D$. From (4.1) and (4.2), we get

$$\phi Q(\nabla_Y \phi X) + Ch(Y, \phi X) + h(Y, X) = 0$$

for all X and $Y \in D$. Replacing Y by ϕZ , we have

$$(4.3) \quad \phi Q(\nabla_{\phi Z} \phi X) + Ch(\phi Z, \phi X) + h(\phi Z, X) = 0,$$

where $Z \in D$. Interchanging X and Z , we have

$$(4.4) \quad \phi Q(\nabla_{\phi X} \phi Z) + Ch(\phi X, \phi Z) + h(\phi X, Z) = 0.$$

Subtracting (4.4) from (4.3), we have

$$\phi Q[\phi X, \phi Z] + h(Z, \phi X) - h(X, \phi Z) = 0,$$

from which the assertion follows. □

LEMMA 4.2. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric non-metric connection. Then it holds*

$$2\bar{\nabla}_Y \phi Z = A_{\phi Y} Z - A_{\phi Z} Y + \nabla_Y^\perp \phi Z - \nabla_Z^\perp \phi Y - \phi[Y, Z].$$

Proof. From the Weingarten formula, we have

$$(4.5) \quad \bar{\nabla}_Y \phi Z - \bar{\nabla}_Z \phi Y = -A_{\phi Z} Y + A_{\phi Y} Z + \nabla_Y^\perp \phi Z - \nabla_Z^\perp \phi Y.$$

Also by covariant differentiation, we get

$$(4.6) \quad \bar{\nabla}_Y \phi Z - \bar{\nabla}_Z \phi Y = (\bar{\nabla}_Y \phi)Z - (\bar{\nabla}_Z \phi)Y + \phi[Y, Z].$$

From (4.5) and (4.6) we have

$$(4.7) \quad (\bar{\nabla}_Y \phi)Z - (\bar{\nabla}_Z \phi)Y = A_{\phi Y} Z - A_{\phi Z} Y + \nabla_Y^\perp \phi Z - \nabla_Z^\perp \phi Y - \phi[Y, Z].$$

From (2.3) we obtain

$$(4.8) \quad (\bar{\nabla}_Y \phi)Z + (\bar{\nabla}_Z \phi)Y = 0$$

for any X and $Y \in D$. Adding (4.7) and (4.8), we get the result. □

PROPOSITION 4.3. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric non-metric connection. Then it holds*

$$(4.9) \quad A_{\phi Y}Z - A_{\phi Z}Y = \frac{1}{3}\phi P[Y, Z],$$

where $[Y, Z]$ is the Lie bracket for $\bar{\nabla}$.

Proof. Let $Y, Z \in D^\perp$ and $X \in x(M)$. Then from (2.4) and (2.6), we have

$$2g(A_{\phi Z}Y, X) = -g(\bar{\nabla}_Y\phi X, Z) - g(\bar{\nabla}_X\phi Y, Z) + g((\bar{\nabla}_Y\phi)X + (\bar{\nabla}_X\phi)Y, Z).$$

By use of (2.3) and $\eta(Y) = 0$ for $Y \in D^\perp$, we have

$$2g(A_{\phi Z}Y, X) = -g(\phi\bar{\nabla}_YZ, X) + g(A_{\phi Y}Z, X).$$

Transvecting X from the both sides, we get

$$2A_{\phi Z}Y = -\phi\bar{\nabla}_YZ + A_{\phi Y}Z.$$

Interchanging Y and Z , we have

$$2A_{\phi Y}Z = -\phi\bar{\nabla}_ZY + A_{\phi Z}Y.$$

Subtracting above two equations, we get the result. □

THEOREM 4.4. *Let M be a semi-invariant submanifold of a nearly Kenmotsu manifold \bar{M} with a semi-symmetric non-metric connection. Then the distribution D^\perp is integrable if and only if*

$$A_{\phi Y}Z - A_{\phi Z}Y = 0$$

for all Y and $Z \in D^\perp$.

Proof. Suppose that the distribution D^\perp is integrable. Then $[Y, Z] \in D^\perp$ for any Y and $Z \in D^\perp$. Therefore, $P[Y, Z] = 0$ and from (4.9), we get

$$(4.10) \quad A_{\phi Y}Z - A_{\phi Z}Y = 0.$$

Conversely let (4.10) holds. Then by virtue of (4.9) we have $\phi P[Y, Z] = 0$ for all Y and $Z \in D^\perp$. Since $\text{rank } \phi = 2n$, therefore we have either $P[Y, Z] = 0$ or $P[Y, Z] = k\xi$. But $P[Y, Z] = k\xi$ is not possible as P being a projection operator on D . Hence $P[Y, Z] = 0$, which is equivalent to $[Y, Z] \in D^\perp$ for all $Z \in D^\perp$ and thus D^\perp is integrable. □

References

- [1] A. Bejancu, *Geometry of CR-submanifolds*, D. Reidel Publishing Company, Holland, 1986.
- [2] D. E. Blair, *Lecture notes in Math.* 509 (1976), Springer-verlag, Berlin.
- [3] A. Friedmann and J. A. Schouten, *Über die geometrie der halbsymmetrischen übertragung*, Math. Zeitschr. **21** (1924), 211-223.
- [4] K. Kenmotsu, *A class of almost contact Riemannian manifold*, Tohoku Math., J. **24**, (1972), 93.
- [5] M. Kobayashi, *Semi-invariant submanifolds of a certain class of almost contact manifolds*, Tensor, N. S. 43 (1986), 28-36.
- [6] N. Papaghiuc, *Semi-invariant submanifolds in a Kenmotsu manifold*, Rend. Math. **3** (1983), 607.
- [7] J. A. Schouten, *Ricci calculus*, Springer, 1954.
- [8] A. Shukla, *Nearly trans-Sasakian manifold*, Kuwait J. Sci. Eng. **23** (1996), no. 2, 139.
- [9] B. B. Sinha and R. N. Yadav, *Some results on semi-invariant submanifolds of a certain class of almost contact manifolds*, Indian J. pure appl. Math. **22** (1991), 783.
- [10] M. M. Tripathi and S.S. Shukla, *Semi-invariant submanifolds of nearly Kenmotsu manifolds*, Bull. Cal. Math. Soc. **95** (2003), no. 1, 17-30.

*

Department of Mathematics
Integral University
Kursi-Road, Lucknow-226026, India
E-mail: mobinahmad@rediffmail.com

**

Department of Mathematics
Kook-Min University
Seoul 136-702, Republic of Korea
E-mail: jbjun@kookmin.ac.kr