

ON LINEAR *BCI*-ALGEBRAS

YOUNG HEE KIM* AND KEUM SOOK SO**

ABSTRACT. In this note, we show that every linear *BCI*-algebra $(X; *, e)$, $e \in X$, has of the form $x * y = x - y + e$ where $x, y \in X$, where X is a field with $|X| \geq 3$.

1. Introduction

Y. Imai and K. Iséki introduced two classes of abstract algebras: *BCK*-algebras and *BCI*-algebras ([3, 4]). It is known that the class of *BCK*-algebras is a proper subclass of the class of *BCI*-algebras. In [1, 2] Q. P. Hu and X. Li introduced a wide class of abstract algebras: *BCH*-algebras. They have shown that the class of *BCI*-algebras is a proper subclass of the class of *BCH*-algebras. J. Neggers and H. S. Kim introduced the notion of *B*-algebras ([8, 9]), i.e., (I) $x * x = e$; (II) $x * e = x$; (III) $(x * y) * z = x * (z * (e * y))$, for any $x, y, z \in X$. C. B. Kim and H. S. Kim ([5]) introduced the notion of a *BG*-algebras which is a generalization of *B*-algebras, and constructed a *BG*-algebra from a non-empty set, which is non-group-derived. A. Walendziak ([12]) introduced a new notion, called an *BF*-algebra, i.e., (I); (II) and (IV) $e * (x * y) = y * x$ for any $x, y \in X$. In ([12]) it was shown that a *BF*-algebra is a generalizations of *B*-algebras. In this note, we show that every linear *BCI*-algebra $(X; *, e)$, $e \in X$, has of the form $x * y = x - y + e$ where $x, y \in X$, where X is a field with $|X| \geq 3$. Moreover, it is shown that every linear *BCI*-algebra is equivalent to quadratic *BF*(*B*, *BG*, *Q*)-algebras.

Received November 02, 2012; Accepted January 11, 2013.

2010 Mathematics Subject Classification: Primary 06F35.

Key words and phrases: *BCI*-algebra, *BF*-algebra, linear, quadratic.

Correspondence should be addressed to Keum Sook So, ksoo@hallym.ac.kr.

2. Preliminaries

DEFINITION 2.1. A *BCI-algebra* is an algebra $(X; *, e)$ of type $(2,0)$ satisfying the following axioms:

- (I) $x * x = e$,
- (II) $x * y = y * x = e$ implies $x = y$,
- (III) $(x * (x * y)) * y = e$,
- (IV) $((x * y) * (x * z)) * (z * y) = e$ for any $x, y, z \in X$.

A *BCK-algebra* is a *BCI-algebra* satisfying the following axiom:

- (V) $e * x = e$ for any $x \in X$.

A *BF-algebra* ([12]) is an algebra $(X; *, e)$ of type $(2,0)$ satisfying the following axioms (I), and

- (B2) $x * e = x$,
- (BF) $e * (x * y) = y * x$ for any $x, y \in X$.

Note that *BF-algebras* need not be *BCI-algebras*, since it does not satisfy the anti-symmetry condition. The following example shows a *BF-algebra* which is not a *BCI-algebra*.

EXAMPLE 2.2. ([12]) Let $(X; *, 0)$ be the algebra where X is the set of all real numbers and $*$ is defined by

$$x * y = \begin{cases} x & \text{if } y = 0 \\ y & \text{if } x = 0 \\ 0 & \text{otherwise} \end{cases}$$

Then X is a *BF-algebra*, but not a *BCI-algebra*, since $3 * 4 = 0 = 4 * 3$, but $3 \neq 4$.

3. Constructions of linear BCI-algebras

Let X be a field with $|X| \geq 3$. An algebra $(X; *)$ is said to be *linear* if $x * y$ is defined by $x * y = Ax + By + C$ where $A, B, C \in X$, for any $x, y \in X$. A linear algebra $(X; *)$ is said to be a *linear BCI-algebra* if it satisfies the conditions (I)~(IV).

THEOREM 3.1. Let X be a field with $|X| \geq 3$. Then every linear *BCI-algebra* $(X; *, e)$, $e \in X$, has of the form $x * y = x - y + e$ where $x, y \in X$.

Proof. Define $x * y := Ax + By + C$ where $A, B, C \in X$ for any $x, y \in X$. If we consider (I), then $e = x * x = (A + B)x + C$ for any $x \in X$. Assume $A + B \neq 0$. Then $x = \frac{e-C}{A+B}$ for any $x \in X$, a contradiction. Hence $A + B = 0$ and $C = e$. Hence

$$(3.1) \quad x * y = A(x - y) + e$$

Consider (II). Suppose $x * y = 0 = y * x$. Then by applying (1), we obtain

$$A(x - y) + e = e = A(y - x) + e$$

Hence $A(x - y) = 0 = A(y - x)$. If we assume that $A \neq 0$, then $x = y$. This means that the necessary condition to hold (III) is that $A \neq 0$.

Consider (IV). Given $x, y \in X$,

$$\begin{aligned} e &= (x * (x * y)) * y \\ &= A(x * (x * y)) + By + C \\ &= A^2(1 + B)x + B(AB + 1)y + C(AB + A + 1) \end{aligned}$$

It follows that $A^2(1 + B) = 0, B(AB + 1) = 0, C(AB + A + 1) = e$.

Case 1: $B = 0$. Since $A^2(1 + B) = 0, A^2 = 0$ and hence $A = 0$, a contradiction.

Case 2: $B \neq 0$. Since $B(AB + 1) = 0, C(AB + A + 1) = e$, we obtain $AB + 1 = 0, CA = e$, i.e., $A = -\frac{1}{B}$ and $C = -Be$. Hence

$$(3.2) \quad x * y = -\frac{1}{B}x + By - Be$$

Subcase (2.1): $C \neq 0$, i.e., $e \neq 0$. For any $x \in X$,

$$\begin{aligned} e &= x * x \\ &= -\frac{1}{B}x + Bx - Be \\ &= \left(-\frac{1}{B} + B\right)x - Be \end{aligned}$$

It follows that $B = -1, A = 1$ and $C = e$. Hence $x * y = x - y + e$. Subcase (2.2): $C = 0$, i.e., $e = 0$. By (2), we have $x * y = -\frac{1}{B}x + By$. If we let $y := x$, then $0 = e = x * x = (B - \frac{1}{B})x$ for any $x \in X$. It follows that $B = \pm 1$. Since $A^2(B + 1) = 0$, we have $B = -1$. In fact, if $B = 1$, then $0 = A^2(1 + B) = 2A^2$, a contradiction, since $A \neq 0$. Thus $x * y = x - y = x - y + 0$, proving the theorem. \square

EXAMPLE 3.2. Let \mathbf{R} be the set of all real numbers. Then $(\mathbf{R}, *, 0)$ is a linear BCI-algebra where $x * y := x - y$, for any $x, y \in \mathbf{R}$.

PROPOSITION 3.3. Let X be a field with $|X| \geq 3$. If X is a linear BCI-algebra, then $(x * z) * (y * z) = x * y$ for any $x, y, z \in X$.

Proof. Straightforward. \square

Let X be a field with $|X| \geq 3$. An algebra $(X; *)$ is said to be quadratic if $x * y$ is defined by $x * y = a_1x^2 + a_2xy + a_3y^2 + a_4x + a_5y + a_6$, where $a_1, a_2, a_3, a_4, a_5, a_6 \in X$, for any $x, y \in X$. A quadratic algebra $(X; *)$ is said to be a quadratic BF-algebra if it satisfies the condition (I), (B2) and (BF).

THEOREM 3.4. ([7]) Let X be a field with $|X| \geq 3$. Then every quadratic BF-algebra $(X; *, e)$, $e \in X$, has of the form $x * y = x - y + e$ where $x, y \in X$.

H. K. Park and H. S. Kim ([11]) proved that every quadratic B-algebra $(X; *, e)$, $e \in X$, has of the form $x * y = x - y + e$, where X is a field with $|X| \geq 3$. J. Neggers, S. S. Ahn and H. S. Kim ([10]) introduced the notions of Q -algebra, and obtained that every quadratic Q -algebra $(X; *, e)$, $e \in X$, has of the form $x * y = x - y + e$, $x, y \in X$, where X is a field with $|X| \geq 3$. Also, H. S. Kim and H. D. Lee ([6]) showed that every quadratic BG-algebra $(X; *, e)$, $e \in X$, has of the form $x * y = x - y + e$, $x, y \in X$, where X is a field with $|X| \geq 3$. We summarize:

THEOREM 3.5. Let X be a field with $|X| \geq 3$. Then the followings are equivalent :

- (1) $(X; *, e)$ is a quadratic BF-algebra,
- (2) $(X; *, e)$ is a quadratic BG-algebra,
- (3) $(X; *, e)$ is a quadratic Q -algebra,
- (4) $(X; *, e)$ is a quadratic B-algebra.

THEOREM 3.6. ([11]) Let X be a field with $|X| \geq 3$. Then every quadratic B-algebra on X is a BCI-algebra.

From Theorem 3.4 and Theorem 3.6, we obtain the following.

COROLLARY 3.7. Let X be a field with $|X| \geq 3$. Then every quadratic BF(BG, Q , B)-algebra on X is a linear BCI-algebra and vice versa.

Acknowledgements

The authors are grateful for the referee's valuable suggestions and help.

References

- [1] Q. P. Hu and X. Li, *On BCH-algebras*, Math. Seminar Notes **11** (1983), 313-320.
- [2] Q. P. Hu and X. Li, *On proper BCH-algebras*, Math. Japonica **30** (1985), 659-661.
- [3] K. Iséki and S. Tanaka, *An introduction to theory of BCK-algebras*, Math. Japonica **23** (1978), 1-26.
- [4] K. Iséki, *On BCI-algebras*, Math. Sem. Notes **8** (1980), 125-130.
- [5] C. B. Kim and H. S. Kim, *On BG-algebras*, Demonstratio Math. **41** (2008), 497-505.
- [6] H. S. Kim and H. D. Lee, *On quadratic BG-algebras*, Int. Math. J. **5** (2004), 529-535.
- [7] H. S. Kim and N. R. Kye, *On quadratic BF-algebras*, Sci. Math. Japo. **65** (2007), 287-290.
- [8] J. Neggers and H. S. Kim, *On B-algebras*, Math. Vesnik **54** (2002), 21-29.
- [9] J. Neggers and H. S. Kim, *A fundamental theorem of B-homomorphism for B-algebras*, Int. Math. J. **2** (2002), 207-214.
- [10] J. Neggers, S. S. Ahn and H. S. Kim, *On Q-algebras*, Int. J. Math Math. Sci. **27** (2001), 749-757.
- [11] H. K. Park and H. S. Kim, *On quadratic B-algebras*, Quasigroups and Related Systems **8** (2001), 67-72.
- [12] A. Walendziak, *On BF-algebras*, Math. Slovaca **57** (2007), 119-128.

*

Department of Mathematics
Chungbuk National University
Chongju 361-791, Republic of Korea
E-mail: yhkim@chungbuk.ac.kr

**

Department of Mathematics
Hallym University
Chuncheon 200-702, Republic of Korea
E-mail: kssso@hallym.ac.kr