

HILBERT 3-CLASS FIELD TOWERS OF REAL CUBIC FUNCTION FIELDS

HWANYUP JUNG*

ABSTRACT. In this paper we study the infiniteness of Hilbert 3-class field tower of real cubic function fields over $\mathbb{F}_q(T)$, where $q \equiv 1 \pmod{3}$. We also give various examples of real cubic function fields whose Hilbert 3-class field tower is infinite.

1. Introduction

Let $k = \mathbb{F}_q(T)$ be a rational function field over the finite field \mathbb{F}_q of q and $\mathbb{A} = \mathbb{F}_q[T]$. For a finite separable extension F of k , write \mathcal{O}_F for the integral closure of \mathbb{A} in F . Let ∞ be the infinite place of k and $S_\infty(F)$ be the set of places of F lying over ∞ . Let ℓ be a prime number. The *Hilbert ℓ -class field of F* is the maximal unramified abelian ℓ -extension of F in which every places $\nu_\infty \in S_\infty(F)$ splits completely. Let $F_0^{(\ell)} = F$ and inductively, $F_{n+1}^{(\ell)}$ be the Hilbert ℓ -class field of $F_n^{(\ell)}$ for $n \geq 0$ (cf. [4]). Then the sequence of fields

$$F_0^{(\ell)} = F \subset F_1^{(\ell)} \subset \dots \subset F_n^{(\ell)} \subset \dots$$

is called the *Hilbert ℓ -class field tower of F* and we say that the Hilbert ℓ -class field tower of F is infinite if $F_n^{(\ell)} \neq F_{n+1}^{(\ell)}$ for each $n \geq 0$. For a multiplicative abelian group A , let $r_\ell(A) = \dim_{\mathbb{F}_\ell}(A/A^\ell)$ be the ℓ -rank of A . Let Cl_F and \mathcal{O}_F^* be the ideal class group and the group of units of \mathcal{O}_F , respectively. Schoof [6] proved that the Hilbert ℓ -class field tower of F is infinite if

$$(1.1) \quad r_\ell(Cl_F) \geq 2 + 2\sqrt{r_\ell(\mathcal{O}_F^*) + 1}.$$

Received March 18, 2013; Accepted July 08, 2013.

2010 Mathematics Subject Classification: Primary 11R58, 11R60, 11R18.

Key words and phrases: Hilbert 3-class field tower, real cubic function field.

This work was conducted during the research year of the Chungbuk National University in 2012 .

In [1], Ahn and Jung studied the infiniteness of the Hilbert 2-class field tower of imaginary quadratic function fields. Assume that q is odd with $q \equiv 1 \pmod 3$. In [3], we studied the infiniteness of the Hilbert 3-class field tower of imaginary cubic function fields. By a *real cubic function field*, we always mean a finite (geometric) cyclic extension F over k of degree 3 in which ∞ splits completely. The aim of this paper is to study the infiniteness of the Hilbert 3-class field tower of real cubic function fields and give a various examples of real cubic function fields which have infinite Hilbert 3-class field tower.

2. Preliminaries

2.1. Rédei matrix and the invariant λ_2

Assume that q is odd with $q \equiv 1 \pmod 3$. Fix a generator γ of \mathbb{F}_q^* . Let $\mathcal{P}(\mathbb{A})$ be the set of all monic irreducible polynomials in \mathbb{A} . Assume that we have given a total ordering “ $<$ ” on the set $\mathcal{P}(\mathbb{A})$ such that $P < Q$ for any $P, Q \in \mathcal{P}(\mathbb{A})$ with $\deg P < \deg Q$. Then any real cubic function field F can be written as $F = k(\sqrt[3]{D})$, where $D = P_1^{r_1} \cdots P_t^{r_t}$ with $P_i \in \mathcal{P}(\mathbb{A})$, $r_i \in \{1, 2\}$ for $2 \leq i \leq t$ and $3 \mid \deg D$. If we assume that $P_1 < \cdots < P_t$ and $r_1 = 1$, then D is uniquely determined by F , and we denote $D_F = D$. We say that D_F is *special* if $3 \mid \deg P_i$ for all $1 \leq i \leq t$. Let σ be a generator of $G = \text{Gal}(F/k)$. We have

$$(2.1)_3(Cl_F) = \dim_{\mathbb{F}_3} (Cl_F/Cl_F^{(1-\sigma)}) + \dim_{\mathbb{F}_3} (Cl_F^{(1-\sigma)}/Cl_F^{(1-\sigma)^2}).$$

Put

$$\lambda_i(F) = \dim_{\mathbb{F}_3} (Cl_F^{(1-\sigma)^{i-1}}/Cl_F^{(1-\sigma)^i}) \quad (i = 1, 2).$$

For $0 \neq N \in \mathbb{A}$, write $\omega(N)$ for the number of distinct monic irreducible divisors of N . Then, by the Genus theory (see [2, Corollary 3.5]), we have

$$(2.2) \quad \lambda_1(F) = \begin{cases} \omega(D_F) - 1, & \text{if } D_F \text{ is special,} \\ \omega(D_F) - 2, & \text{otherwise.} \end{cases}$$

Let $\eta = \gamma^{\frac{q-1}{3}}$. Let M_F be the $t \times t$ matrix $(e_{ij})_{1 \leq i, j \leq t}$ over \mathbb{F}_3 , where, for $1 \leq i \neq j \leq t$, $e_{ij} \in \mathbb{F}_3$ is defined by $\eta^{e_{ij}} = (\frac{P_i}{P_j})_3$, and the diagonal entries e_{ii} are defined to satisfy the relation $\sum_{i=1}^t r_i e_{ij} = 0$. Let $d_i \in \mathbb{F}_3$ be defined by $d_i \equiv \deg P_i \pmod 3$ for $1 \leq i \leq t$. We associate a $(t+1) \times t$ matrix R_F over \mathbb{F}_3 to F as follows:

- If D_F is non-special or D_F is special with $\gamma \in N_{F/k}(\mathcal{O}_F^*)$, then R_F is the $(t+1) \times t$ matrix obtained from M_F by adjoining (d_1, \dots, d_t) in last row.
- If D_F is special with $\gamma \notin N_{F/k}(\mathcal{O}_F^*)$, then R_F is the $(t+1) \times t$ matrix obtained from M_F by adjoining (e_{B1}, \dots, e_{Bt}) in last row, where $B \in \mathbb{A}$ is a monic polynomial such that $B = \mathcal{N}(\mathfrak{B})$, $\mathfrak{B}^{\sigma-1} = x\mathcal{O}_F$, $N_{F/k}(x) \in \mathbb{F}_q^* \setminus \mathbb{F}_q^{*3}$ and $e_{Bi} \in \mathbb{F}_3$ is defined by $\eta^{e_{iB}} = (\frac{P_i}{B})$.

Then we have (see [2, Corollary 3.8])

$$(2.3) \quad \lambda_2(F) = \omega(D_F) - 1 - \text{rank } R_F.$$

Hence, by inserting (2.2) and (2.3) into (2.1), we have

$$(2.4) \quad r_3(Cl_F) = \begin{cases} 2\omega(D_F) - 2 - \text{rank } R_F, & \text{if } D_F \text{ is special,} \\ 2\omega(D_F) - 3 - \text{rank } R_F, & \text{otherwise.} \end{cases}$$

2.2. Some lemmas

Let E and K be finite (geometric) separable extensions of k such that E/K is a cyclic extension of degree ℓ , where ℓ is a prime number not dividing q . Let $\gamma_{E/K}$ be the number of prime ideals of \mathcal{O}_K that ramify in E and $\rho_{E/K}$ be the number of places $\mathfrak{p}_\infty \in S_\infty(K)$ that ramify or inert in E . If

$$(2.5) \quad \gamma_{E/K} \geq |S_\infty(K)| - \rho_{E/K} + 3 + 2\sqrt{\ell|S_\infty(K)| + (1 - \ell)\rho_{E/K} + 1},$$

then E has infinite Hilbert ℓ -class field tower (see [1, Proposition 2.1]). By using this criterion, we give some sufficient conditions for a real cubic function field to have infinite Hilbert 3-class field tower. For $D \in \mathbb{A}$, write $\pi(D)$ for the set of all monic irreducible divisors of D .

LEMMA 2.1. *Assume that q is odd with $q \equiv 1 \pmod 3$. Let F be a real cubic function field over k . If there is a nonconstant monic polynomial D' such that $3 \mid \deg D'$, $\pi(D') \subset \pi(D_F)$ and $(\frac{D'}{P_i})_3 = 1$ for $P_i \in \pi(D) \setminus \pi(D')$ ($1 \leq i \leq 5$), then F has infinite Hilbert 3-class field tower.*

Proof. Put $K = k(\sqrt[3]{D'})$ and $E = FK$. Since $\gamma_{E/K} \geq 15$, $|S_\infty(K)| = 3$ and $\rho_{E/K} = 0$, the inequality (2.5) is satisfied, so E has infinite Hilbert 3-class field tower. By hypothesis, P_1, P_2, P_3, P_4, P_5 and ∞ split completely in K , so E is contained in $F_1^{(3)}$. Hence F also has infinite Hilbert 3-class field tower. □

LEMMA 2.2. *Assume that q is odd with $q \equiv 1 \pmod 3$. Let F be a real cubic function field over k . If there are two distinct nonconstant monic polynomials D_1, D_2 such that $3 \mid \deg D_i$, $\pi(D_i) \subset \pi(D_F)$ for $i = 1, 2$ and*

$(\frac{D_1}{P_j})_3 = (\frac{D_2}{P_j})_3 = 1$ for some $P_j \in \pi(D_F) \setminus (\pi(D_1) \cup \pi(D_2))$ ($1 \leq j \leq 3$), then F has infinite Hilbert 3-class field tower.

Proof. Put $K = k(\sqrt[3]{D_1}, \sqrt[3]{D_2})$ and $E = FK$. Since $\gamma_{E/K} \geq 27$, $|S_\infty(K)| = 9$ and $\rho_{E/K} = 0$, the inequality (2.5) is satisfied, so E has infinite Hilbert 3-class field tower. By hypothesis, P_1, P_2, P_3 and ∞ split completely in K , so E is contained in $F_1^{(3)}$. Hence F also has infinite Hilbert 3-class field tower. \square

3. Hilbert 3-class field tower of real cubic function field

Assume that q is odd with $q \equiv 1 \pmod 3$. In this section we give several sufficient conditions for real cubic function fields to have infinite Hilbert 3-class field tower and examples.

Let F be a real cubic function field. Since $r_3(\mathcal{O}_F^*) = 2$, by Schoof's theorem, the Hilbert 3-class field tower of F is infinite if $r_3(\mathcal{C}l_F) \geq 6$. By (2.2), F has infinite Hilbert 3-class field tower if $\omega(D_F) \geq 7$ or $\omega(D_F) \geq 8$ according as D_F is special or not. We will consider the cases that $\omega(D_F) \leq 6$ if D_F is special and $\omega(D_F) \leq 7$ if D_F is non-special.

THEOREM 3.1. *Assume that q is odd with $q \equiv 1 \pmod 3$. Let F be a real cubic function field.*

- (i) *Assume that D_F is special. If $\omega(D_F) = 6$ with $\text{rank } R_F \leq 4$, $\omega(D_F) = 5$ with $\text{rank } R_F \leq 2$ or $\omega(D_F) = 4$ with $\text{rank } R_F = 0$, then F has infinite Hilbert 3-class field tower.*
- (ii) *Assume that D_F is non-special. If $\omega(D_F) = 7$ with $\text{rank } R_F \leq 5$, $\omega(D_F) = 6$ with $\text{rank } R_F \leq 3$ or $\omega(D_F) = 5$ with $\text{rank } R_F \leq 1$, then F has infinite Hilbert 3-class field tower.*

Proof. By using the Schoof's theorem with (2.4), we see that F has infinite Hilbert 3-class field tower if

$$\text{rank } R_F \leq \begin{cases} 2\omega(D_F) - 8, & \text{if } D_F \text{ is special,} \\ 2\omega(D_F) - 9, & \text{if } D_F \text{ is non-special.} \end{cases}$$

Hence the result follows immediately. \square

THEOREM 3.2. *Assume that q is odd with $q \equiv 1 \pmod 3$. Let F be a real cubic function field with $\omega(D_F) \geq 6$. If there exists $Q \in \pi(D_F)$ with $3 \mid \deg Q$ and $P_i \in \pi(D_F) \setminus \{Q\}$ such that $(\frac{Q}{P_i})_3 = 1$ ($1 \leq i \leq 5$), then F has infinite Hilbert 3-class field tower.*

Proof. By applying Lemma 2.1 with $D' = Q$, we see that F has infinite Hilbert 3-class field tower. \square

EXAMPLE 3.3. Let $k = \mathbb{F}_7(T)$ and $\mathbb{A} = \mathbb{F}_7[T]$. Take $Q = T^3 + 2T^2 + 1$, $P_1 = T$, $P_2 = T + 2$, $P_3 = T + 3$, $P_4 = T^2 + 2$ and $P_5 = T^3 + T + 1$. By simple computations, we see that $(\frac{Q}{P_i})_3 = 1$ for $1 \leq i \leq 5$. Then, for any $D = Q^e P_1^{e_1} P_2^{e_2} P_3^{e_3} P_4^{e_4} P_5^{e_5}$ with $e, e_i \in \{1, 2\}$ and $e_1 + e_2 + e_3 + 2e_4 \equiv 0 \pmod 3$, $k(\sqrt[3]{D})$ is a real cubic function field whose Hilbert 3-class field tower is infinite by Theorem 3.2.

Let $N(n, q)$ be the number of monic irreducible polynomials of degree n in $\mathbb{A} = \mathbb{F}_q[T]$. Then it satisfies the following one ([5, Corollary of Proposition 2.1]):

$$N(n, q) = \frac{1}{n} \sum_{d|n} \mu(d) q^{\frac{n}{d}}.$$

For $\alpha \in \mathbb{F}_q^*$, let $\mathcal{N}(n, \alpha, q)$ be the set of monic irreducible polynomials of degree n with constant term α in $\mathbb{A} = \mathbb{F}_q[T]$ and $N(n, \alpha, q) = |\mathcal{N}(n, \alpha, q)|$. Define

$$D_n = \{r : r|(q^n - 1), r \nmid (q^m - 1) \text{ for } m < n\}.$$

For each $r \in D_n$, let $r = m_r d_r$, where $d_r = \gcd(r, \frac{q^n - 1}{q - 1})$. In [7], Yucas proved that $N(n, \alpha, q)$ satisfies the following formula:

$$(3.1) \quad N(n, \alpha, q) = \frac{1}{n\phi(f)} \sum_{\substack{r \in D_n \\ m_r = f}} \phi(r),$$

where f is the order of α in \mathbb{F}_q^* .

EXAMPLE 3.4. Let $k = \mathbb{F}_7(T)$ and $\mathbb{A} = \mathbb{F}_7[T]$. We will find the number $N(3, 1, 7)$. We have

$$D_3 = \{r : r|(7^3 - 1), r \nmid (7^2 - 1), r \nmid (7 - 1)\} = \{9, 18, 19, 38, 57, 114, 171, 342\}$$

and $m_9 = 1$, $m_{18} = 6$, $m_{19} = 1$, $m_{38} = 2$, $m_{57} = 1$, $m_{114} = 2$, $m_{171} = 3$, $m_{342} = 6$. Since the order of 1 in \mathbb{F}_7^* is 1, by (3.1), we have

$$N(3, 1, 7) = \frac{1}{3\phi(1)} (\phi(9) + \phi(19) + \phi(57)) = 20.$$

Hence, $\mathcal{N}(3, 1, 7)$ consists of exactly 20 distinct monic irreducible polynomials. Take $Q = T$, $P_1 = T^2 + \alpha T + 1$ with $\alpha \in \{0, 3, 4\}$ and $P_2, P_3, P_4, P_5 \in \mathcal{N}(3, 1, 7)$. Then, for any $D = Q^e P_1^{e_1} P_2^{e_2} P_3^{e_3} P_4^{e_4} P_5^{e_5}$ with $e, e_i \in \{1, 2\}$ and $e_1 + e_2 + e_3 + 2e_4 \equiv 0 \pmod 3$, we have $(\frac{Q}{P_i})_3 = 1$

for $1 \leq i \leq 5$, so $k(\sqrt[3]{D})$ is a real cubic function field whose Hilbert 3-class field tower is infinite by Theorem 3.2.

THEOREM 3.5. *Assume that q is odd with $q \equiv 1 \pmod 3$. Let F be a real cubic function field with $\omega(D_F) \geq 5$. If there exist $Q_1, Q_2 \in \pi(D_F)$ whose degrees are divisible by 3 and $P_i \in \pi(D_F) \setminus \{Q_1, Q_2\}$ such that $(\frac{Q_1}{P_i})_3 = (\frac{Q_2}{P_i})_3 = 1$ ($1 \leq i \leq 3$), then F has infinite Hilbert 3-class field tower.*

Proof. By applying Lemma 2.2 with $D_1 = Q_1$ and $D_2 = Q_2$, we see that F has infinite Hilbert 3-class field tower. □

EXAMPLE 3.6. *Let $k = \mathbb{F}_7(T)$ and $\mathbb{A} = \mathbb{F}_7[T]$. Take $P_1 = T, P_2 = T + 1, P_3 = T^2 + 1, Q_1 = T^3 + T^2 + 1$ and $Q_2 = T^3 + T + 1$. By simple computations, we see that $(\frac{Q_1}{P_i})_3 = (\frac{Q_2}{P_i})_3 = 1$ for $1 \leq i \leq 3$. Then, for any $D = P_1^{e_1} P_2^{e_2} P_3^{e_3} Q_1^{f_1} Q_2^{f_2}$ with $e_i, f_j \in \{1, 2\}$ and $e_1 + e_2 + 2e_3 \equiv 0 \pmod 3$, $k(\sqrt[3]{D})$ is a real cubic function field whose Hilbert 3-class field tower is infinite by Theorem 3.5.*

THEOREM 3.7. *Assume that q is odd with $q \equiv 1 \pmod 3$. Let F be a real cubic function field with non-special D_F . Assume that $\omega(D_F) = 6$ or $\omega(D_F) = 7$. If there exist $Q_1, Q_2, Q_3 \in \pi(D_F)$ whose degrees are divisible by 3 and $P_i \in \pi(D_F) \setminus \{Q_1, Q_2, Q_3\}$ such that $(\frac{Q_1 Q_2}{P_i})_3 = (\frac{Q_1 Q_3}{P_i})_3 = 1$ ($1 \leq i \leq 3$), then F has infinite Hilbert 3-class field tower.*

Proof. By applying Lemma 2.2 with $D_1 = Q_1 Q_2$ and $D_2 = Q_1 Q_3$, we see that F has infinite Hilbert 3-class field tower. □

EXAMPLE 3.8. *Let $k = \mathbb{F}_7(T)$ and $\mathbb{A} = \mathbb{F}_7[T]$. Take $P_1 = T, P_2 = T + 2, P_3 = T^2 + 1, Q_1 = T^3 + T + 1, Q_2 = T^3 + T^2 + 1$ and $Q_3 = T^3 + T - 1$. By simple computations, we see that*

$$\left(\frac{Q_i}{P_1}\right)_3 = \left(\frac{Q_i}{P_3}\right)_3 = 1 \ (1 \leq i \leq 3), \quad \left(\frac{Q_1}{P_2}\right)_3 = \eta^2, \quad \left(\frac{Q_2}{P_2}\right)_3 = \left(\frac{Q_3}{P_2}\right)_3 = \eta.$$

Hence, we have

$$\left(\frac{Q_1 Q_2}{P_i}\right)_3 = \left(\frac{Q_1 Q_3}{P_i}\right)_3 = 1 \ (1 \leq i \leq 3).$$

Then, for any $D = P_1^{e_1} P_2^{e_2} P_3^{e_3} Q_1^{f_1} Q_2^{f_2}$ with $e_i, f_j \in \{1, 2\}$ and $e_1 + e_2 + 2e_3 \equiv 0 \pmod 3$, $k(\sqrt[3]{D})$ is a real cubic function field whose Hilbert 3-class field tower is infinite by Theorem 3.7.

References

- [1] J. Ahn and H. Jung, *Hilbert 2-class field towers of imaginary quadratic function fields*. J. Chungcheong Math. Soc. **23** (2010), no. 4, 699–704.
- [2] S. Bae, S. Hu, and H. Jung, *The generalized Rédei-matrix for function fields*. Finite Fields Appl. **18** (2012), no. 4, 760–780.
- [3] H. Jung, *Hilbert 3-class field towers of imaginary cubic function fields*. J. Chungcheong Math. Soc. **26** (2013), no. 1, 161–166.
- [4] M. Rosen, *The Hilbert class field in function fields*. Exposition. Math. **5** (1987), no. 4, 365–378.
- [5] ———, *Number theory in function fields*. Graduate Texts in Mathematics **210** (2002), Springer-Verlag, New York.
- [6] R. Schoof, *Algebraic curves over \mathbb{F}_2 with many rational points*. J. Number Theory **41** (1992), no. 1, 6–14.
- [7] J. L. Yucas, *Irreducible polynomials over finite fields with prescribed trace/prescribed constant term*. Finite Fields Appl. **12** (2006), no. 2, 211–221.

*

Department of Mathematics Education
Chungbuk National University
Cheongju 361-763, Republic of Korea
E-mail: hyjung@chungbuk.ac.kr