

THE RELATION BETWEEN MCSHANE INTEGRAL AND MCSHANE DELTA INTEGRAL

JAE MYUNG PARK*, DEOK HO LEE**, JU HAN YOON***, YOUNG
KUK KIM****, AND JONG TAE LIM*****

ABSTRACT. In this paper, we define an extension $f^* : [a, b] \rightarrow \mathbb{R}$ of a function $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ for a time scale \mathbb{T} and show that f is McShane delta integrable on $[a, b]_{\mathbb{T}}$ if and only if f^* is McShane integrable on $[a, b]$.

1. Introduction and preliminaries

First, we introduce some concepts related to the notion of time scales. A time scale \mathbb{T} is any closed nonempty subset of \mathbb{R} . For each $t \in \mathbb{T}$, we define the forward jump operator $\sigma(t)$ by

$$\sigma(t) = \inf\{z > t : z \in \mathbb{T}\}$$

and the backward jump operator $\rho(t)$ by

$$\rho(t) = \sup\{z < t : z \in \mathbb{T}\}$$

where $\inf \phi = \sup \mathbb{T}$ and $\sup \phi = \inf \mathbb{T}$.

If $\sigma(t) > t$, we say the t is right-scattered, while if $\rho(t) < t$, we say that t is left-scattered. If $\sigma(t) = t$, we say that t is right-dense, while if $\rho(t) = t$, we say that t is left-dense. The forward graininess function $\mu(t)$ is defined by $\mu(t) = \sigma(t) - t$, and the backward graininess function $\nu(t)$ is defined by $\nu(t) = t - \rho(t)$.

For $a, b \in \mathbb{T}$, we define the time scale interval in \mathbb{T} by

$$[a, b]_{\mathbb{T}} = \{t \in \mathbb{T} : a \leq t \leq b\}.$$

A pair $\delta = (\delta_L, \delta_R)$ of two real-valued functions on $[a, b]_{\mathbb{T}}$ is a Δ -gauge on $[a, b]_{\mathbb{T}}$ by $\delta_L(t) > 0$ on $(a, b]_{\mathbb{T}}$, $\delta_R(t) > 0$ on $[a, b)_{\mathbb{T}}$, $\delta_L(a) \geq 0$, $\delta_R(b) \geq 0$, and $\delta_R(t) \geq \mu(t)$ for each $t \in [a, b]_{\mathbb{T}}$.

Received December 24, 2013; Accepted January 16, 2014.

2010 Mathematics Subject Classification: Primary 26A39; Secondary 26E70.

Key words and phrases: time scales, McShane delta integral, Δ -gauge.

Correspondence should be addressed to Ju Han Yoon, yoonyh@cbnu.ac.kr.

A collection $\mathcal{P} = \{(\xi_i, [t_{i-1}, t_i]_{\mathbb{T}})\}_{i=1}^n$ of tagged intervals is a δ -fine McShane partition of $[a, b]_{\mathbb{T}}$ if $\bigcup_{i=1}^n [t_{i-1}, t_i]_{\mathbb{T}} = [a, b]_{\mathbb{T}}$, $[t_{i-1}, t_i]_{\mathbb{T}} \subset [\xi_i - \delta_L(\xi_i), \xi_i + \delta_R(\xi_i)]$ and $\xi_i \in [a, b]_{\mathbb{T}}$ for each $i = 1, 2, \dots, n$.

For a partition $\mathcal{P} = \{(\xi_i, [t_{i-1}, t_i]_{\mathbb{T}})\}_{i=1}^n$, we write

$$f(\mathcal{P}) = \sum_{i=1}^n f(\xi_i)(t_i - t_{i-1}),$$

whenever $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$.

2. The McShane and McShane delta integrals

DEFINITION 2.1. A function $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ is McShane delta integrable (or M_{Δ} -integrable) on $[a, b]_{\mathbb{T}}$ if there exists a number A such that for each $\epsilon > 0$ there exists a Δ -gauge δ on $[a, b]_{\mathbb{T}}$ such that

$$\left| f(\mathcal{P}) - A \right| < \epsilon$$

for every δ -fine McShane partition \mathcal{P} of $[a, b]_{\mathbb{T}}$. The number A is called the M_{Δ} -integral of f on $[a, b]_{\mathbb{T}}$, and we write $A = (M_{\Delta}) \int_a^b f$.

Recall that $f : [a, b] \rightarrow \mathbb{R}$ is McShane integrable (or M-integrable) on $[a, b]$ if there exists a number A such that for each $\epsilon > 0$ there exists a gauge $\delta : [a, b] \rightarrow \mathbb{R}^+$ on $[a, b]$ such that

$$\left| f(\mathcal{P}) - A \right| < \epsilon$$

for every δ -fine McShane partition \mathcal{P} of $[a, b]$.

Let $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ be a function on $[a, b]_{\mathbb{T}}$, and let $\{(a_k, b_k)\}_{k=1}^{\infty}$ be the sequence of intervals contiguous to $[a, b]_{\mathbb{T}}$ in $[a, b]$.

Define a function $f^* : [a, b] \rightarrow \mathbb{R}$ on $[a, b]$ by

$$f^*(t) = \begin{cases} f(a_k) & \text{if } t \in (a_k, b_k) \text{ for some } k. \\ f(t) & \text{if } t \in [a, b]_{\mathbb{T}}. \end{cases}$$

It is well-known [6] that $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ is Henstock delta(or H_{Δ})-integrable on $[a, b]_{\mathbb{T}}$ if and only if $f^* : [a, b] \rightarrow \mathbb{R}$ is Henstock(or H)-integrable, and $(H_{\Delta}) \int_a^b f = (H) \int_a^b f^*$.

THEOREM 2.2. *If $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ is M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$, then $f^* : [a, b] \rightarrow \mathbb{R}$ is M-integrable on $[a, b]$. In this case, $(M_{\Delta}) \int_a^b f = (M) \int_a^b f^*$.*

Proof. Let $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ be M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$ and let $\epsilon > 0$. Then there exists a Δ -gauge $\delta = (\delta_L, \delta_R)$ on $[a, b]_{\mathbb{T}}$ such that

- (1) $|f(\mathcal{P}) - (M_\Delta) \int_a^b f| < \epsilon$ for each δ -fine McShane partition $\mathcal{P} = \{(\xi_i, [a_i, b_i])\}$ of $[a, b]_{\mathbb{T}}$,
- (2) $\delta_R(t) = \sigma(t) - t$ if $t \in [a, b]_{\mathbb{T}}^{rs}$, where $[a, b]_{\mathbb{T}}^{rs}$ is the set of all right scattered points of $[a, b]_{\mathbb{T}}$,
- (3) $t + \delta_R(t) \in [a, b]_{\mathbb{T}}$ if $t \in [a, b]_{\mathbb{T}}^{rd}$, where $[a, b]_{\mathbb{T}}^{rd}$ is the set of all right dense points of $[a, b]_{\mathbb{T}}$,
- (4) $\delta_L(t) < \min \left\{ \frac{s_k - \rho(s_k)}{3}, \frac{\epsilon}{2^k(|f(s_k)| + |f(\rho(s_k))| + 1)} \right\}$ if $t = s_k$, where $[a, b]_{\mathbb{T}}^{ls} = \{s_1, s_2, s_k, \dots\}$, and
- (5) $t - \delta_L(t) \in [a, b]_{\mathbb{T}}$ if $t \in [a, b]_{\mathbb{T}}^{ld}$, where $[a, b]_{\mathbb{T}}^{ld}$ is the set of all left dense points of $[a, b]_{\mathbb{T}}$.

Define a gauge $\delta^1 = (\delta_L^1, \delta_R^1)$ on $[a, b]$ by

$$\delta^1(t) = \begin{cases} (\delta_L(t), \delta_R(t)) & \text{if } t \in [a, b]_{\mathbb{T}} \\ (t - \rho(s_k), s_k - t) & \text{if } t \in (\rho(s_k), s_k), \quad k = 1, 2, 3, \dots \end{cases}$$

Assume that $D = \{(\zeta_i, [u_i, v_i])\}_{i=1}^n$ is a δ^1 -fine McShane partition of $[a, b]$. Let $i \leq n$ and $k \in \mathbb{N}$ be given. Suppose That $(\rho(s_k), s_k) \not\subset (u_i, v_i)$, $(\rho(s_k), s_k) \not\supset (u_i, v_i)$ and $(\rho(s_k), s_k) \cap (u_i, v_i) \neq \emptyset$. Then we note that the substituting $(\zeta_i, [\rho(s_k), s_k] \cap [u_i, v_i])$ and $(\zeta_i, (\rho(s_k), s_k)^c \cap [u_i, v_i])$ instead of $(\zeta_i, [u_i, v_i])$ in D does not alter the value of $f^*(D)$.

Hence we may assume that

$(\rho(s_k), s_k) \subset (u_i, v_i)$ or $(\rho(s_k), s_k) \supset (u_i, v_i)$ or $(\rho(s_k), s_k) \cap (u_i, v_i) = \emptyset$ for each $i \leq n$ and $k \in \mathbb{N}$.

Let K be the set of all $k \in \mathbb{N}$ such that $(u_i, v_i) \subset (\rho(s_k), s_k)$ for some $i \leq n$. Then K is clearly a finite set. Denote $K = \{k_1, k_2, \dots, k_q\}$.

Let $A_j = \{i \leq n | (u_i, v_i) \subset (\rho(s_{k_j}), s_{k_j})\}$ for $j = 1, 2, \dots, q$,

$$A = \bigcup_{j=1}^q A_j \quad \text{and} \quad B = \{1, 2, \dots, n\} - A.$$

If $i \in B$, then $\zeta_i \in [a, b]_{\mathbb{T}}$ and $u_i, v_i \in [a, b]_{\mathbb{T}}$.

Let $A_j^1 = \{i \in A_j | \rho(s_{k_j}) \leq \zeta_i \leq s_{k_j}\}$ and $A_j^2 = A_j - A_j^1$.

Define

$$\zeta_i^1 = \begin{cases} \zeta_i & \text{if } i \in B \\ \rho(s_k) & \text{if } i \in A_j^1 \quad (j = 1, 2, \dots, q) \\ \zeta_i & \text{if } i \in A_j^2 \quad (j = 1, 2, \dots, q). \end{cases}$$

Let $D^1 = \{(\zeta_i^1, [u_i, v_i])\}_{i=1}^n$. Then clearly D^1 is a δ^1 -fine McShane partition on $[a, b]$ and $\zeta_i^1 \in [a, b]_{\mathbb{T}}$ for each $i \leq n$, and

$$\begin{aligned}
& |f^*(D^1) - f^*(D)| \\
&= \left| \sum_{i \in A} [f^*(\zeta_i^1) - f^*(\zeta_i)](v_i - u_i) \right| \\
&= \left| \sum_{j=1}^q \sum_{i \in A_j^1} [f^*(\zeta_i^1) - f^*(\zeta_i)](v_i - u_i) \right| \\
&\leq \sum_{j=1}^q \sum_{i \in A_j^1} |f^*(\zeta_i^1) - f^*(\zeta_i)|(v_i - u_i) \\
&\leq \sum_{j=1}^q \left[(|f^*(\rho(s_{k_j}))| + |f^*(s_{k_j})|) \sum_{i \in A_j^1} (v_i - u_i) \right] \\
&< \sum_{j=1}^q \frac{\epsilon (|f(\rho(s_{k_j}))| + |f(s_{k_j})|)}{2^{k_j} (|f(\rho(s_{k_j}))| + |f(s_{k_j})| + 1)} < \epsilon.
\end{aligned}$$

For each j , choose $\lambda_j, \mu_j \in A_j$ such that

$$f(\zeta_{\lambda_j}^1) = \min_{i \in A_j} f(\zeta_i^1) \quad \text{and} \quad f(\zeta_{\mu_j}^1) = \max_{i \in A_j} f(\zeta_i^1).$$

Then the partitions

$$D_1^1 = \{(\zeta_i^1, [u_i, v_i]) | i \in B\} \cup \{(\zeta_{\lambda_j}^1, [\rho(s_{k_j}), s_{k_j}]) | j = 1, 2, \dots, q\}$$

and

$$D_2^1 = \{(\zeta_i^1, [u_i, v_i]) | i \in B\} \cup \{(\zeta_{\mu_j}^1, [\rho(s_{k_j}), s_{k_j}]) | j = 1, 2, \dots, q\}$$

are both δ -fine McShane partitions on $[a, b]_{\mathbb{T}}$ and

$$\begin{aligned}
f^*(D^1) &= \sum_{i \in B} f(\zeta_i^1)(v_i - u_i) + \sum_{j=1}^q \sum_{i \in A_j} f(\zeta_i^1)(v_i - u_i) \\
&\leq \sum_{i \in B} f(\zeta_i^1)(v_i - u_i) + \sum_{j=1}^q f(\zeta_{\mu_j}^1)(s_{k_j} - \rho(s_{k_j})) \\
&= f(D_2^1) < (M_{\Delta}) \int_a^b f + \epsilon.
\end{aligned}$$

Similarly, we have

$$f^*(D^1) \geq f(D_1^1) > (M_\Delta) \int_a^b f - \epsilon,$$

so that $|f^*(D^1) - (M_\Delta) \int_a^b f| < \epsilon$.

Hence,

$$\begin{aligned} \left| f^*(D) - (M_\Delta) \int_a^b f \right| &\leq |f^*(D) - f^*(D^1)| + \left| f^*(D^1) - (M_\Delta) \int_a^b f \right| \\ &< 2\epsilon. \end{aligned}$$

Thus, f^* is McShane integrable and

$$(M) \int_a^b f^* = (M_\Delta) \int_a^b f.$$

□

THEOREM 2.3. *If $f^* : [a, b] \rightarrow \mathbb{R}$ is M-integrable on $[a, b]$, then $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ is M_Δ -integrable on $[a, b]_{\mathbb{T}}$.*

Proof. Suppose that f^* is M-integrable on $[a, b]$. Define $g : [a, b] \rightarrow \mathbb{R}$ by

$$g(t) = \begin{cases} |f(\rho(t))| + |f(t)| & \text{if } t \in [a, b]_{\mathbb{T}}^{ls} \\ 0 & \text{if } t \in [a, b] - [a, b]_{\mathbb{T}}^{ls}. \end{cases}$$

Then since $[a, b]_{\mathbb{T}}^{ls}$ is a countable set, $g = 0$ a.e. on $[a, b]$. Hence, g is M-integrable on $[a, b]$ and $(M) \int_a^b g = 0$. Let $\epsilon > 0$. Then there exists a gauge $\delta = (\delta_L, \delta_R)$ on $[a, b]$ such that

$$\left| f^*(D) - (M) \int_a^b f^* \right| < \epsilon \quad \text{and} \quad |g(D)| < \epsilon$$

for each δ -fine McShane partition D of $[a, b]$.

Define a Δ -gauge $\delta^1 = (\delta_L^1, \delta_R^1)$ on $[a, b]_{\mathbb{T}}$ by

$$\begin{aligned} \delta_L^1(t) &= \delta_L(t) & \text{if } t \in [a, b]_{\mathbb{T}} \\ \delta_R^1(t) &= \begin{cases} \delta_R(t) & \text{if } t \in [a, b]_{\mathbb{T}}^{rd} \\ \sigma(t) - t & \text{if } t \in [a, b]_{\mathbb{T}}^{rs}. \end{cases} \end{aligned}$$

Assume that $D = \{(\xi_i, [u_i, v_i])\}_{i=1}^n$ is a δ^1 -fine McShane partition of $[a, b]_{\mathbb{T}}$. Let $A = \{i \leq n \mid \xi_i \in [a, b]_{\mathbb{T}}^{rs}, [\xi_i, \sigma(\xi_i)] \subset [u_i, v_i]\}$,

$$B = \{1, 2, \dots, n\} - A.$$

For each $i \in A$, choose a δ -fine McShane partition $D_i = \{(\xi_{ij}, [u_{ij}, v_{ij}])\}_{j=1}^{p_i}$ of $[\xi_i, \sigma(\xi_i)]$.

Let $D^* = \{(\xi_i, [u_i, v_i])\}_{i \in B} \cup \{(\xi_i, [u_i, \xi_i]) \mid i \in A, u_i < \xi_i\} \cup \left[\bigcup_{i \in A} D_i \right]$. Then D^* is a δ -fine McShane partition of $[a, b]$, and

$$\begin{aligned} |f(D) - f^*(D^*)| &= \left| \sum_{i \in A} f(\xi_i)(\sigma(\xi_i) - \xi_i) - \sum_{i \in A} \sum_{j \leq p_i} f^*(\xi_{ij})(v_{ij} - u_{ij}) \right| \\ &= \left| \sum_{i \in A} \left[f(\xi_i)(\sigma(\xi_i) - \xi_i) - \sum_{\substack{j \leq p_i \\ \xi_{ij} < \sigma(\xi_i)}} f(\xi_{ij})(v_{ij} - u_{ij}) \right. \right. \\ &\quad \left. \left. - \sum_{\substack{i \leq p_i \\ \xi_{ij} = \sigma(\xi_i)}} f(\sigma(\xi_i))(v_{ij} - u_{ij}) \right] \right| \\ &= \left| \sum_{i \in A} \sum_{\substack{j \leq p_i \\ \xi_{ij} = \sigma(\xi_i)}} \left[f(\xi_i) - f(\sigma(\xi_i)) \right] (v_{ij} - u_{ij}) \right| \\ &\leq \sum_{i \in A} \sum_{\substack{j \leq p_i \\ \xi_{ij} = \sigma(\xi_i)}} \left[|f(\xi_i)| + |f(\sigma(\xi_i))| \right] (v_{ij} - u_{ij}) < \epsilon. \end{aligned}$$

Hence,

$$\begin{aligned} &\left| f(D) - (M) \int_a^b f^* \right| \\ &\leq |f(D) - f^*(D^*)| + \left| f^*(D^*) - (M) \int_a^b f^* \right| < 2\epsilon. \end{aligned}$$

Thus, f is M_Δ -integrable on $[a, b]_{\mathbb{T}}$ and

$$(M_\Delta) \int_a^b f = (M) \int_a^b f^*.$$

□

From Theorem 2.2 and 2.3, we get the following theorem.

THEOREM 2.4. *A function $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ is M_Δ -integrable $[a, b]_{\mathbb{T}}$ if and only if $f^* : [a, b] \rightarrow \mathbb{R}$ is M -integrable on $[a, b]$. In this case,*

$$(M_\Delta) \int_a^b f = (M) \int_a^b f^*.$$

We next verify the basic properties of the McShane delta integral.

THEOREM 2.5. *Let $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ and let $c \in [a, b]$.*

- (1) *If f is M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$, then f is M_{Δ} -integrable on every subinterval of $[a, b]_{\mathbb{T}}$.*
- (2) *If f is M_{Δ} -integrable on each of the intervals $[a, c]_{\mathbb{T}}$ and $[c, b]_{\mathbb{T}}$, then f is M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$ and*

$$(M_{\Delta}) \int_a^b f = (M_{\Delta}) \int_a^c f + (M_{\Delta}) \int_c^b f.$$

Proof. (1) If f is M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$, then f^* is M-integrable on $[a, b]$. Since f^* is M-integrable on every subinterval of $[a, b]$, f is M_{Δ} -integrable on every subinterval of $[a, b]_{\mathbb{T}}$ by Theorem 2.4.

(2) If f is M_{Δ} -integrable on each of $[a, c]_{\mathbb{T}}$ and $[c, b]_{\mathbb{T}}$, then f^* is M-integrable on each of $[a, c]$ and $[c, b]$. By the property of the McShane integral, f^* is M-integrable on $[a, b]$ and

$$(M) \int_a^b f^* = (M) \int_a^c f^* + (M) \int_c^b f^*.$$

By Theorem 2.4, f is M_{Δ} -integrable on $[a, c]_{\mathbb{T}}$ and $[c, b]_{\mathbb{T}}$ and

$$(M_{\Delta}) \int_a^b f = (M_{\Delta}) \int_a^c f + (M_{\Delta}) \int_c^b f.$$

□

THEOREM 2.6. *Let $f, g : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ be M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$. Then $\alpha f + \beta g$ is M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$ for every $\alpha, \beta \in \mathbb{R}$ and*

$$(M_{\Delta}) \int_a^b (\alpha f + \beta g) = \alpha (M_{\Delta}) \int_a^b f + \beta (M_{\Delta}) \int_a^b g.$$

Proof. If f and g are M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$, then f^* and g^* are M-integrable on $[a, b]$. Hence $\alpha f^* + \beta g^*$ is M-integrable on $[a, b]$, and $(M) \int_a^b (\alpha f^* + \beta g^*) = \alpha (M) \int_a^b f^* + \beta (M) \int_a^b g^*$. By Theorem 2.4, $\alpha f + \beta g$ is M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$ and

$$(M_{\Delta}) \int_a^b (\alpha f + \beta g) = \alpha (M_{\Delta}) \int_a^b f + \beta (M_{\Delta}) \int_a^b g.$$

□

THEOREM 2.7. (Monotone Convergence Theorem) *Let $\{f_n\}$ be a monotone sequence of M_{Δ} -integrable functions defined on $[a, b]_{\mathbb{T}}$ and*

suppose that $\{f_n\}$ converges pointwise to f on $[a, b]_{\mathbb{T}}$. If $\lim_{n \rightarrow \infty} (M_{\Delta}) \int_a^b f_n$ is finite, then f is M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$ and

$$(M_{\Delta}) \int_a^b f = \lim_{n \rightarrow \infty} (M_{\Delta}) \int_a^b f_n.$$

Proof. Let $\{f_n\}$ be a monotone sequence of M_{Δ} -integrable functions on $[a, b]_{\mathbb{T}}$. Then by Theorem 2.4, $\{f_n^*\}$ is a monotone sequence of M-integrable functions on $[a, b]$, and $\{f_n^*\}$ converges pointwise to f^* on $[a, b]$. Since $(M_{\Delta}) \int_a^b f_n = (M) \int_a^b f_n^*$ for each n , $\lim_{n \rightarrow \infty} (M) \int_a^b f_n^*$ is finite. By the Monotone Convergence Theorem for the M-integral, f^* is M-integrable on $[a, b]$ and

$$(M) \int_a^b f^* = \lim_{n \rightarrow \infty} (M) \int_a^b f_n^*.$$

By Theorem 2.4, f is M_{Δ} -integrable on $[a, b]_{\mathbb{T}}$ and

$$(M_{\Delta}) \int_a^b f = \lim_{n \rightarrow \infty} (M_{\Delta}) \int_a^b f_n.$$

□

References

- [1] R. Agarwal and M. Bohner, Basic calculus on time scales and some of its applications, *Results Math.* **35** (1999), 3-22
- [2] S. Avsec, B. Bannish, B. Johnson, and S. Meckler, *The Henstock-Kurzweil delta integral on unbounded time scales*, PanAmerican Math. J. **16** (2006), no. 3, 77-98.
- [3] G. Sh. Guseinov, *Intergration on time scales*, J. Math. Anal. Appl. **285** (2003), 107-127.
- [4] G. Sh. Guseinov and B. Kaymakcalan, *Basics of Riemann delta and nabla integration on time scales*, J. Difference Equations Appl. **8** (2002), 1001-1027.
- [5] J. M. Park, D. H. Lee, J. H. Yoon, and J. T. Lim, *The Henstock and Henstock delta Integrals*, J. Chungcheng Math. Soc. **26** (2013), no. 2, 291-298.
- [6] J. M. Park, D. H. Lee, J. H. Yoon, Y. K. Kim, and J. T. Lim, *The relation between Henstock integral and Henstock delta Integral on time scales*, J. Chungcheng Math. Soc. **26** (2013), no. 3, 625-630.
- [7] A. Perterson and B. Thompson, *Henstock-Kurzweil Delta and Nabla Integral*, J. Math. Anal. Appl. **323** (2006), 162-178.
- [8] C. W. Swartz and D. S. Kurtz, *Theories of Integration: The Integrals of Riemann Lebesgue, Henstock-Kurzweil, and Mcshane*, World Scientific, 2004.
- [9] B. S. Thomson, *Henstock Kurzweil integrals on time scales*, PanAmerican Math J. **18** (2008), no. 1, 1-19.

*

Department of Mathematics
Chungnam National University
Daejeon 305-764, Republic of Korea
E-mail: parkjm@cnu.ac.kr

**

Department of Mathematics Education
KongJu National University
Kongju 314-701, Republic of Korea
E-mail: dhlee@kongju.ac.kr

Department of Mathematics Education
Chungbuk National University
Chungju 360-763, Republic of Korea
E-mail: yoonjh@cbnu.ac.kr

Department of Mathematics Education
Seowon University
Chungju 361-742, Republic of Korea
E-mail: ykkim@dragon.seowon.ac.kr

Department of Mathematics
Chungnam National University
Daejeon 305-764, Republic of Korea
E-mail: ssongtae@edurang.dje.go.kr