

## LORENTZIAN ALMOST $r$ -PARA-CONTACT STRUCTURE IN TANGENT BUNDLE

MOHAMMAD NAZRUL ISLAM KHAN\* AND JAE-BOK JUN\*\*

ABSTRACT. Almost contact and almost complex structures in the tangent bundle have been studied by K. Yano and S. Ishihara[5] and others. In the present paper, we have studied Lorentzian almost  $r$ -para-contact structure in the tangent bundle. Some results related to Lie-derivative have been studied.

### 1. Introduction

Let  $M$  be a  $n$ -dimensional differentiable manifold of  $C^\infty$  class and  $T_p(M)$  the tangent space of  $M$  at a point  $p$  of  $M$ . Then the set  $T(M) = \bigcup_{p \in M} T_p(M)$  is called the *tangent bundle* over the manifold  $M$ . For any point  $\tilde{p}$  of  $T(M)$ , the correspondence  $\tilde{p} \rightarrow p$  determines the bundle projection  $\pi : T(M) \rightarrow M$ . Thus  $\pi(\tilde{p}) = p$ , where  $\pi : T(M) \rightarrow M$  defines the bundle projection of  $T(M)$  over  $M$ . The set  $\pi^{-1}(p)$  is called the *fibre* over  $p \in M$  and  $M$  the *base space*.

#### Vertical lifts:

If  $f$  is a function in  $M$ , then we write  $f^V$  for the function in  $T(M)$  obtained by forming the composition of  $\pi : T(M) \rightarrow M$  and  $f : M \rightarrow R$  so that  $f^V = f \circ \pi$ . Thus, if a point  $\tilde{p} \in \pi^{-1}(U)$  has induced coordinates  $(x^h, y^h)$ , then

$$f^V(\tilde{p}) = f^V(x, y) = f \circ \pi(\tilde{p}) = f(p) = f(x).$$

Thus the value of  $f^V(\tilde{p})$  is constant along each fibre  $T_p(M)$  and equal to the value  $f(p)$ . We call  $f^V$  the *vertical lift* of the function  $f$ .

---

Received August 21, 2013; Revised September 29, 2013; Accepted January 16, 2014.

2010 Mathematics Subject Classification: Primary 53D15, 53B35.

Key words and phrases: tangent bundle, vertical lifts, complete lifts, Lie derivative.

Correspondence should be addressed to Jae-Bok Jun, [jbjun@kookmin.ac.kr](mailto:jbjun@kookmin.ac.kr).

Vertical lifts to a unique algebraic isomorphism of the tensor algebra  $\mathfrak{S}(M)$  into the tensor algebra  $\mathfrak{S}(T(M))$  with respect to constant coefficients by the conditions

$$(P \otimes Q)^V = P^V \otimes Q^V, \quad (P + R)^V = P^V + R^V,$$

where  $P, Q$  and  $R$  are arbitrary elements of  $\mathfrak{S}(M)$ .

### Complete lifts:

If  $f$  is a function in  $M$ , then we write  $f^C$  for the function in  $T(M)$  defined by  $f^C = i(df)[4]$  and call  $f^C$  the *complete lift* of the function  $f$ . The complete lift  $f^C$  of a function  $f$  has the local expression  $f^C = y^i \partial_i f = \partial f$  with respect to the induced coordinates in  $T(M)$ , where  $\partial f$  denotes  $y^i \partial_i f$ .

Suppose that  $X \in \mathfrak{S}_0^1(M)$ . We define a vector field  $X^C$  in  $T(M)$  by  $X^C f^C = (Xf)^C$ , where  $f$  is an arbitrary function in  $M$  and call  $X^C$  the *complete lift* of  $X$  in  $T(M)$ . The complete lift  $X^C$  of  $X$  with components  $x^h$  in  $M$  has components

$$X^C = \begin{pmatrix} x^h \\ \partial x^h \end{pmatrix}$$

with respect to the induced coordinates in  $T(M)$ .

Suppose that  $\omega \in \mathfrak{S}_1^0(M)$ . Then a 1-form  $\omega^C$  in  $T(M)$  is defined by  $\omega^C(X^C) = (\omega(X))^C$ , where  $X$  is an arbitrary vector field in  $M$ . We call  $\omega^C$  the *complete lift* of  $\omega$ .

The complete lifts to a unique algebra isomorphism of the tensor algebra  $\mathfrak{S}(M)$  into the tensor algebra  $\mathfrak{S}(T(M))$  with respect to constant coefficients is given by the conditions

$$(P \otimes Q)^C = P^C \otimes Q^V + P^V \otimes Q^C, \quad (P + R)^C = P^C + R^C,$$

where  $P, Q$  and  $R$  are arbitrary elements of  $\mathfrak{S}(M)$ .

### Horizontal lifts:

The *horizontal lift*  $f^H$  of  $f \in \mathfrak{S}_0^0(M)$  to the tangent bundle  $T(M)$  is defined by  $f^H = f^C - \nabla_\gamma f$ , where  $\nabla_\gamma f = \gamma(\nabla f)$ .

Let  $X \in \mathfrak{S}_0^1(M)$ . Then the horizontal lift  $X^H$  of  $X$  is defined in  $T(M)$  by

$$X^H = X^C - \nabla_\gamma X,$$

where  $\nabla_\gamma X = \gamma(\nabla X)$ . The horizontal lift  $X^H$  of  $X$  has the components

$$X^H = \begin{pmatrix} x^h \\ -\Gamma_i^h x^i \end{pmatrix}$$

with respect to the induced coordinates in  $T(M)$ , where  $\Gamma_i^h = y^j \Gamma_{ji}^h$ .

The horizontal lift  $S^H$  of a tensor field  $S$  of arbitrary type in  $M$  to  $T(M)$  is defined by

$$S^H = S^C - \nabla_\gamma S.$$

For any  $P, Q \in \mathfrak{S}(M)$ , we have

$$\nabla_\gamma(P \otimes Q) = (\nabla_\gamma P) \otimes Q^V + P^V \otimes (\nabla_\gamma Q)$$

or

$$(P \otimes Q)^H = P^H \otimes Q^V + P^V \otimes Q^H.$$

## 2. Almost product structure

Let  $M$  be a  $n$ -dimensional differentiable manifold of  $C^\infty$  class. If there exists a tensor field  $F$  of type (1,1) and of  $C^\infty$  class on  $M$  such that

$$F^2 = I,$$

where  $I$  denotes the unit tensor field. Then we say that  $F$  gives to  $M$  an *almost product structure*.

## 3. Complete lifts of almost product structure and Lorentzian almost $r$ -para-contact structure in the tangent bundle

Let  $\bar{M}$  be a  $(2n + r)$ -dimensional differentiable manifold of  $C^\infty$  class and  $T(\bar{M})$  denotes the tangent bundle of  $\bar{M}$ . Suppose that there are given a tensor field  $F$  of type (1,1), a vector field  $U_\alpha$  and a 1-form  $\omega^\alpha$  on  $T(\bar{M})$  satisfying

$$(3.1) \quad F^2 = I + \sum_{\alpha=1}^r U_\alpha \otimes \omega^\alpha,$$

where

$$(3.2) \quad \begin{aligned} FU_\alpha &= 0, \\ \omega^\alpha \circ F &= 0, \\ \omega^\alpha(U_\beta) &= \delta_\beta^\alpha. \end{aligned}$$

Thus the manifold  $\bar{M}$  satisfying conditions (3.1) and (3.2) will be said to possess *Lorentzian almost  $r$ -para-contact structure* ([1], [3]).

**THEOREM 3.1.** *Let  $\bar{M}$  be a differentiable manifold endowed with Lorentzian almost  $r$ -para-contact structure  $(F, U_\alpha, \omega^\alpha)$ . Then*

$$\tilde{J} = F^C + (U_\alpha^V \otimes \omega^{\alpha V} - U_\alpha^C \otimes \omega^{\alpha C})$$

*is an almost product structure on  $T(\bar{M})$ .*

*Proof.* From (3.1) and (3.2), we have [2]

$$(3.3) \quad (F^C)^2 = I + (U_\alpha^V \otimes \omega^{\alpha C} - U_\alpha^C \otimes \omega^{\alpha V})$$

and

$$(3.4) \quad \begin{aligned} F^C U_\alpha^V &= 0, \quad F^C U_\alpha^C = 0, \\ \omega^{\alpha V} \circ F^C &= 0, \quad \omega^{\alpha C} \circ F^V = 0, \quad \omega^{\alpha C} \circ F^C = 0, \\ \omega^{\alpha V}(U_\alpha^V) &= 0, \quad \omega^{\alpha C}(U_\alpha^C) = 1, \quad \omega^{\alpha C}(U_\alpha^V) = 1, \quad \omega^{\alpha C}(U_\alpha^C) = 0. \end{aligned}$$

Let us define an element  $\tilde{J}$  of  $J(T(\bar{M}))$  by

$$(3.5) \quad \tilde{J} = F^C + (U_\alpha^V \otimes \omega^{\alpha V} - U_\alpha^C \otimes \omega^{\alpha C}).$$

Then we find by (3.3), (3.4) and (3.5),

$$\tilde{J}^2 = I.$$

Thus  $\tilde{J}$  is an almost product structure in  $T(\bar{M})$ . □

In view of equation (3.5), we have

$$(3.6) \quad \begin{aligned} \tilde{J}X^V &= -(FX)^V + (\omega^\alpha(X))^V U_\alpha^C, \\ \tilde{J}X^C &= -(FX)^C - (\omega^\alpha(X))^V U_\alpha^C - (\omega^\alpha(X))^C U_\alpha^C. \end{aligned}$$

In particular, we have

$$(3.7) \quad \tilde{J}X^V = -(FX)^V, \quad \tilde{J}X^C = -(FX)^C,$$

$$(3.8) \quad \tilde{J}U_\alpha^V = U_\alpha^C, \quad \tilde{J}U_\alpha^C = U_\alpha^C,$$

where  $X$  is an arbitrary vector field in  $M$  such that  $\omega^\alpha(X) = 0$ .

**THEOREM 3.2.** *Let the tangent bundle  $T(M)$  of the manifold  $M$  admits  $\tilde{J}$  defined in (3.5). Then for any vector fields  $X, Y$  such that  $\omega^\alpha(Y) = 0$ , we have*

- (i)  $(L_X V \tilde{J})Y^V = 0$ ,
- (ii)  $(L_X V \tilde{J})Y^C = -((L_X F)Y)^V + ((L_X \omega^\alpha)Y)^V U_\alpha^C$ ,
- (iii)  $(L_X V \tilde{J})U_\alpha^V = (L_X U_\alpha)^V$ ,
- (iv)  $(L_X V \tilde{J})U_\alpha^C = -((L_X F)U_\alpha)^V + (L_X \omega^\alpha(U_\alpha))^V U_\alpha^C$

and

$$\begin{aligned}
\text{(i)'} \quad & (L_X C \tilde{J}) Y^V = -((L_X F) Y)^V + (L_X \omega^\alpha(Y))^V U_\alpha^C, \\
\text{(ii)'} \quad & (L_X C \tilde{J}) Y^C = -((L_X F) Y)^C - ((L_X \omega^\alpha) Y) U_\alpha + ((L_X \omega^\alpha) Y)^C U_\alpha^C, \\
\text{(iii)'} \quad & (L_X C \tilde{J}) U_\alpha^V = ((L_X F) U_\alpha)^C + [X, U_\alpha]^C + ((L_X \omega^\alpha) U_\alpha)^V U_\alpha^C, \\
\text{(iv)'} \quad & (L_X C \tilde{J}) U_\alpha^C = ((L_X F) U_\alpha)^C - [X, U_\alpha]^V - (L_X \omega^\alpha(U_\alpha))^V U_\alpha \\
& \quad \quad \quad + ((L_X \omega^\alpha) U_\alpha)^C U_\alpha^C,
\end{aligned}$$

where  $L$  is the Lie-derivative and  $[ , ]$  is the Lie-Bracket.

*Proof.* The proof follows easily from (3.4), (3.6), (3.8) and [5].  $\square$

#### 4. Horizontal lifts of Lorentzian almost $r$ -para-contact structure

Let  $(F, U_\alpha, \omega^\alpha)$  be Lorentzian almost  $r$ -para-contact structure in  $\bar{M}$  with an affine connection. Then we have from (3.1) and (3.2) and [5],

$$\begin{aligned}
& (F^H)^2 = (I + U_\alpha \otimes \omega^\alpha)^H, \\
(4.1) \quad & (F^H)^2 = I + (U_\alpha \otimes \omega^\alpha)^H, \\
& (F^H)^2 = I + U_\alpha^H \otimes \omega^{\alpha V} + U_\alpha^V \otimes \omega^{\alpha H}.
\end{aligned}$$

Also,

$$\begin{aligned}
& F^H U_\alpha^H = 0, \quad F^H U_\alpha^V = 0, \\
(4.2) \quad & \omega^{\alpha H}(U_\alpha^H) = 0, \quad \omega^{\alpha H}(U_\alpha^V) = 1, \quad \omega^{\alpha V}(U_\alpha^H) = 1, \\
& \omega^{\alpha H} \circ f^H = 0, \quad \omega^{\alpha V} \circ f^H = 0.
\end{aligned}$$

Let us define a tensor field  $\tilde{J}^*$  of type (1,1) in  $T(\bar{M})$  by

$$\tilde{J}^* = F^H + U_\alpha^V \otimes \omega^{\alpha V} - U_\alpha^H \otimes \omega^{\alpha H}.$$

Then it is easy to show from (4.1) and (4.2) that

$$\tilde{J}^{*2} = I,$$

which means that  $\tilde{J}^*$  is an almost product structure in  $T(\bar{M})$ . Thus we have

**THEOREM 4.1.** *Let  $(F, U_\alpha, \omega^\alpha)$  be Lorentzian almost  $r$ -para-contact structure in  $\bar{M}$  with an affine connection  $\nabla$ . Then  $\tilde{J}^*$  is a Lorentzian almost product structure in  $T(\bar{M})$ .*

### References

- [1] T. Adati, *An almost  $r$ -para contact Riemannian manifold and an almost product Riemannian Manifold*, Tensor, N. S. **43** (1986), 239-247.
- [2] Lovejoy S. Das and Mohd. Nazrul Islam Khan, *Almost  $r$ -contact structures on the tangent bundle*, Differential Geometry-Dynamical Systems **7** (2005), 34-41.
- [3] K. Matsumoto, *On Lorentzian paracontact manifolds*, Bull. of Yamagata Univ. Nat. Sci. **12** (1989), no. 2, 151-156.
- [4] K. Yano and S. Ishihara, *Almost complex structures induced in tangent bundles*, Kodai Math. Sem. Rep. **19** (1967), 1-27.
- [5] K. Yano and S. Ishihara, *Tangent and cotangent Bundles*, Marcel Dekker, Inc. New York, 1973.

\*

Department of Computer Science  
Qassim University Kingdom of Saudi Arabia  
*E-mail*: mnazrul@rediffmail.com

\*\*

Department of Mathematics  
Kookmin University  
Seoul 136-702, Republic of Korea  
*E-mail*: jbjun@kookmin.ac.kr