

DOUBLE PAIRWISE $(r, s)(u, v)$ -SEMICONTINUOUS MAPPINGS

EUN PYO LEE* AND SEUNG ON LEE**

ABSTRACT. We introduce the concepts of $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double (r, s) (u, v) -semiclosures and $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiinteriors. Using the notions, we investigate some of characteristic properties of double pairwise $(r, s)(u, v)$ -semicontinuous, double pairwise $(r, s)(u, v)$ -semiopen and double pairwise $(r, s)(u, v)$ -semiclosed mappings.

1. Introduction

Chang [2] defined fuzzy topological spaces. These spaces and its generalizations are later studied by several authors, one of which, developed by Šostak [13], used the idea of degree of openness. This type of generalization of fuzzy topological spaces was later rephrased by Chat-topadhyay, Hazra, and Samanta [3], and by Ramadan [12].

As a generalization of fuzzy sets, the concept of intuitionistic fuzzy sets was introduced by Atanassov [1]. Çoker and his colleagues [4, 6, 7] introduced intuitionistic fuzzy topological spaces using intuitionistic fuzzy sets. Using the idea of degree of openness and degree of nonopenness, Çoker and M. Demirci [5] defined intuitionistic fuzzy topological spaces in Šostak's sense as a generalization of smooth fuzzy topological spaces and intuitionistic fuzzy topological spaces.

Kandil [8] introduced and studied the notion of fuzzy bitopological spaces as a natural generalization of fuzzy topological spaces.

In this paper, we introduce the concepts of $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosures and $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiinteriors. Using the

Received August 09, 2014; Accepted October 10, 2014.

2010 Mathematics Subject Classification: Primary 54A40, 03E72.

Key words and phrases: double $(r, s)(u, v)$ -semiclosures, double $(r, s)(u, v)$ -semiinteriors, double pairwise $(r, s)(u, v)$ -semicontinuous.

Correspondence should be addressed to Seung On Lee, solee@chungbuk.ac.kr.

This work was supported by the research grant of Chungbuk National University in 2013.

notions, we investigate some of characteristic properties of double pairwise $(r, s)(u, v)$ -semicontinuous, double pairwise $(r, s)(u, v)$ -semiopen and double pairwise $(r, s)(u, v)$ -semiclosed mappings.

2. Preliminaries

Let I be the unit interval $[0, 1]$ of the real line. A member μ of I^X is called a fuzzy set of X . For any $\mu \in I^X$, μ^c denotes the complement $1 - \mu$. By $\tilde{0}$ and $\tilde{1}$ we denote constant maps on X with value 0 and 1, respectively. All other notations are standard notations of fuzzy set theory.

Let X be a nonempty set. An *intuitionistic fuzzy set* A is an ordered pair

$$A = (\mu_A, \gamma_A)$$

where the functions $\mu_A : X \rightarrow I$ and $\gamma_A : X \rightarrow I$ denote the degree of membership and the degree of nonmembership, respectively, and $\mu_A + \gamma_A \leq \tilde{1}$.

Obviously every fuzzy set μ on X is an intuitionistic fuzzy set of the form $(\mu, \tilde{1} - \mu)$.

DEFINITION 2.1. [1] Let $A = (\mu_A, \gamma_A)$ and $B = (\mu_B, \gamma_B)$ be intuitionistic fuzzy sets on X . Then

- (1) $A \subseteq B$ iff $\mu_A \leq \mu_B$ and $\gamma_A \geq \gamma_B$.
- (2) $A = B$ iff $A \subseteq B$ and $B \subseteq A$.
- (3) $A^c = (\gamma_A, \mu_A)$.
- (4) $A \cap B = (\mu_A \wedge \mu_B, \gamma_A \vee \gamma_B)$.
- (5) $A \cup B = (\mu_A \vee \mu_B, \gamma_A \wedge \gamma_B)$.
- (6) $0_\sim = (\tilde{0}, \tilde{1})$ and $1_\sim = (\tilde{1}, \tilde{0})$.

Let f be a mapping from a set X to a set Y . Let $A = (\mu_A, \gamma_A)$ be an intuitionistic fuzzy set of X and $B = (\mu_B, \gamma_B)$ an intuitionistic fuzzy set of Y . Then:

- (1) The image of A under f , denoted by $f(A)$, is an intuitionistic fuzzy set in Y defined by

$$f(A) = (f(\mu_A), \tilde{1} - f(\tilde{1} - \gamma_A)).$$

- (2) The inverse image of B under f , denoted by $f^{-1}(B)$, is an intuitionistic fuzzy set in X defined by

$$f^{-1}(B) = (f^{-1}(\mu_B), f^{-1}(\gamma_B)).$$

An *intuitionistic fuzzy topology* on X is a family T of intuitionistic fuzzy sets in X which satisfies the following properties:

- (1) $0_{\sim}, 1_{\sim} \in T$.
- (2) If $A_1, A_2 \in T$, then $A_1 \cap A_2 \in T$.
- (3) If $A_i \in T$ for all i , then $\bigcup A_i \in T$.

The pair (X, T) is called an *intuitionistic fuzzy topological space*.

Let $I(X)$ be a family of all intuitionistic fuzzy sets of X and let $I \otimes I$ be the set of the pair (r, s) such that $r, s \in I$ and $r + s \leq 1$.

DEFINITION 2.2. [13] Let X be a nonempty set. An *intuitionistic fuzzy topology in Šostak's sense* $\mathcal{T}^{\mu\gamma} = (\mathcal{T}^{\mu}, \mathcal{T}^{\gamma})$ on X is a mapping $\mathcal{T}^{\mu\gamma} : I(X) \rightarrow I \otimes I$ ($\mathcal{T}^{\mu}, \mathcal{T}^{\gamma} : I(X) \rightarrow I$) which satisfies the following properties:

- (1) $\mathcal{T}^{\mu}(0_{\sim}) = \mathcal{T}^{\mu}(1_{\sim}) = 1$ and $\mathcal{T}^{\gamma}(0_{\sim}) = \mathcal{T}^{\gamma}(1_{\sim}) = 0$.
- (2) $\mathcal{T}^{\mu}(A \cap B) \geq \mathcal{T}^{\mu}(A) \wedge \mathcal{T}^{\mu}(B)$ and $\mathcal{T}^{\gamma}(A \cap B) \leq \mathcal{T}^{\gamma}(A) \vee \mathcal{T}^{\gamma}(B)$.
- (3) $\mathcal{T}^{\mu}(\bigcup A_i) \geq \bigwedge \mathcal{T}^{\mu}(A_i)$ and $\mathcal{T}^{\gamma}(\bigcup A_i) \leq \bigvee \mathcal{T}^{\gamma}(A_i)$.

The $(X, \mathcal{T}^{\mu\gamma}) = (X, \mathcal{T}^{\mu}, \mathcal{T}^{\gamma})$ is said to be an *intuitionistic fuzzy topological space in Šostak's sense*. Also, we call $\mathcal{T}^{\mu}(A)$ a *gradation of openness* of A and $\mathcal{T}^{\gamma}(A)$ a *gradation of nonopenness* of A .

DEFINITION 2.3. [11] Let A be an intuitionistic fuzzy set in an intuitionistic fuzzy topological space in Šostak's sense $(X, \mathcal{T}^{\mu}, \mathcal{T}^{\gamma})$ and $(r, s) \in I \otimes I$. Then A is said to be

- (1) a $\mathcal{T}^{\mu\gamma}$ -fuzzy (r, s) -open set if $\mathcal{T}^{\mu}(A) \geq r$ and $\mathcal{T}^{\gamma}(A) \leq s$,
- (2) a $\mathcal{T}^{\mu\gamma}$ -fuzzy (r, s) -closed set if $\mathcal{T}^{\mu}(A^c) \geq r$ and $\mathcal{T}^{\gamma}(A^c) \leq s$.

Let $(X, \mathcal{T}^{\mu}, \mathcal{T}^{\gamma})$ be an intuitionistic fuzzy topological space in Šostak's sense. For each $(r, s) \in I \otimes I$ and for each $A \in I(X)$, the $\mathcal{T}^{\mu\gamma}$ -fuzzy (r, s) -closure is defined by

$$\begin{aligned} \mathcal{T}^{\mu\gamma}\text{-cl}(A, (r, s)) &= \bigcap \{B \in I(X) \mid A \subseteq B, B \text{ is } \mathcal{T}^{\mu\gamma}\text{-fuzzy } (r, s)\text{-closed}\} \end{aligned}$$

and the $\mathcal{T}^{\mu\gamma}$ -fuzzy (r, s) -interior is defined by

$$\begin{aligned} \mathcal{T}^{\mu\gamma}\text{-int}(A, (r, s)) &= \bigcup \{B \in I(X) \mid A \supseteq B, B \text{ is } \mathcal{T}^{\mu\gamma}\text{-fuzzy } (r, s)\text{-open}\}. \end{aligned}$$

A system $(X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ consisting of a set X with two intuitionistic fuzzy topologies in Šostak's sense $\mathcal{T}^{\mu\gamma}$ and $\mathcal{U}^{\mu\gamma}$ on X is called a *double bitopological space*.

DEFINITION 2.4. [10] Let A be an intuitionistic fuzzy set of a double bitopological space $(X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ and $(r, s), (u, v) \in I \otimes I$. Then A is said to be

- (1) a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiopen set if there is a $\mathcal{T}^{\mu\gamma}$ -fuzzy (r, s) -open set B in X such that $B \subseteq A \subseteq \mathcal{U}^{\mu\gamma}\text{-cl}(B, (u, v))$,
- (2) a $(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiopen set if there is an $\mathcal{U}^{\mu\gamma}$ -fuzzy (u, v) -open set B in X such that $B \subseteq A \subseteq \mathcal{T}^{\mu\gamma}\text{-cl}(B, (r, s))$,
- (3) a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosed set if there is a $\mathcal{T}^{\mu\gamma}$ -fuzzy (r, s) -closed set B in X such that $\mathcal{U}^{\mu\gamma}\text{-int}(B, (u, v)) \subseteq A \subseteq B$,
- (4) a $(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiclosed set if there is an $\mathcal{U}^{\mu\gamma}$ -fuzzy (u, v) -closed set B in X such that $\mathcal{T}^{\mu\gamma}\text{-int}(B, (r, s)) \subseteq A \subseteq B$.

3. Double pairwise $(r, s)(u, v)$ -semicontinuous mappings

DEFINITION 3.1. Let $(X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ be a double bitopological space and $(r, s), (u, v) \in I \otimes I$. For each $A \in I(X)$, the $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosure is defined by

$$(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v)) = \bigcap \{B \in I(X) \mid A \subseteq B, \\ B \text{ is } (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-double } (r, s)(u, v)\text{-semiclosed}\}$$

and the $(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiclosure is defined by

$$(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(A, (u, v), (r, s)) = \bigcap \{B \in I(X) \mid A \subseteq B, \\ B \text{ is } (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-double } (u, v)(r, s)\text{-semiclosed}\}.$$

DEFINITION 3.2. Let $(X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ be a double bitopological space and $(r, s), (u, v) \in I \otimes I$. For each $A \in I(X)$, the $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiinterior is defined by

$$(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)) = \bigcup \{B \in I(X) \mid A \supseteq B, \\ B \text{ is } (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-double } (r, s)(u, v)\text{-semiopen}\}$$

and the $(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiinterior is defined by

$$(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(A, (u, v), (r, s)) = \bigcup \{B \in I(X) \mid A \supseteq B, \\ B \text{ is } (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-double } (u, v)(r, s)\text{-semiopen}\}.$$

Obviously, $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v))$ is the smallest $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosed set which contains A and $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v)) = A$ for any $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosed set A . Also $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v))$ is the greatest $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiopen set which is contained A and $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)) = A$ for any $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiopen set A . Moreover, we have

$$\begin{aligned} \mathcal{T}^{\mu\gamma}\text{-int}(A, (r, s)) &\subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)) \\ &\subseteq A \\ &\subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v)) \\ &\subseteq \mathcal{T}^{\mu\gamma}\text{-cl}(A, (r, s)). \end{aligned}$$

Also, we have the following results:

- (1) $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(0_{\sim}, (r, s), (u, v)) = 0_{\sim}$ and $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(1_{\sim}, (r, s), (u, v)) = 1_{\sim}$.
- (2) $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v)) \supseteq A$.
- (3) $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A \cup B, (r, s), (u, v)) \supseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v)) \cup (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(B, (r, s), (u, v))$.
- (4) $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v)), (r, s), (u, v)) = (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v))$.
- (5) $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(0_{\sim}, (r, s), (u, v)) = 0_{\sim}$ and $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(1_{\sim}, (r, s), (u, v)) = 1_{\sim}$.
- (6) $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)) \subseteq A$.
- (7) $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A \cap B, (r, s), (u, v)) \subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)) \cap (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(B, (r, s), (u, v))$.
- (8) $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)), (r, s), (u, v)) = (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v))$.

THEOREM 3.3. For an intuitionistic fuzzy set A of a double bitopological space $(X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ and $(r, s), (u, v) \in I \otimes I$, we have:

- (1) $((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)))^c = (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v))$.
- (2) $((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A, (r, s), (u, v)))^c = (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A^c, (r, s), (u, v))$.

Proof. (1) Since $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v))$ is a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosed set and $A^c \subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v))$, we have $((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v)))^c$ is a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiopen set of X and $((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v)))^c \subseteq A$. Thus

$$\begin{aligned} &((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v)))^c \\ &= (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v)))^c, (r, s), (u, v)) \\ &\subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)) \end{aligned}$$

and hence

$$((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)))^c \subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v)).$$

Conversely, since $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v))$ is a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiopen set and $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)) \subseteq A$, $((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)))^c$ is a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosed set of X and $A^c \subseteq ((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)))^c$. Thus

$$\begin{aligned} & (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(A^c, (r, s), (u, v)) \\ & \subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)))^c, (r, s), (u, v)) \\ & = ((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(A, (r, s), (u, v)))^c. \end{aligned}$$

(2) Similar to (1) □

Let $f : (X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma}) \rightarrow (Y, \mathcal{V}^{\mu\gamma}, \mathcal{W}^{\mu\gamma})$ be a mapping from a double bitopological space X to a double bitopological space Y and $(r, s), (u, v) \in I \otimes I$. Then f is called a *double pairwise $(r, s)(u, v)$ -continuous $((r, s)(u, v)$ -open and $(r, s)(u, v)$ -closed, respectively) mapping* if the induced mapping $f : (X, \mathcal{T}^{\mu\gamma}) \rightarrow (Y, \mathcal{V}^{\mu\gamma})$ is fuzzy (r, s) -continuous $((r, s)$ -open and (r, s) -closed, respectively) and the induced mapping $f : (X, \mathcal{U}^{\mu\gamma}) \rightarrow (Y, \mathcal{W}^{\mu\gamma})$ is fuzzy (u, v) -continuous $((u, v)$ -open and (u, v) -closed, respectively).

DEFINITION 3.4. Let $f : (X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma}) \rightarrow (Y, \mathcal{V}^{\mu\gamma}, \mathcal{W}^{\mu\gamma})$ be a mapping from a double bitopological space X to a double bitopological space Y and $(r, s), (u, v) \in I \otimes I$. Then f is called

- (1) *double pairwise $(r, s)(u, v)$ -semicontinuous* if $f^{-1}(A)$ is a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiopen set of X for each $\mathcal{V}^{\mu\gamma}$ -fuzzy (r, s) -open set A of Y and $f^{-1}(B)$ is a $(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiopen set of X for each $\mathcal{W}^{\mu\gamma}$ -fuzzy (u, v) -open set B of Y ,
- (2) *double pairwise $(r, s)(u, v)$ -semiopen* if $f(C)$ is a $(\mathcal{V}^{\mu\gamma}, \mathcal{W}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiopen set of Y for each $\mathcal{T}^{\mu\gamma}$ -fuzzy (r, s) -open set C of X and $f(D)$ is a $(\mathcal{W}^{\mu\gamma}, \mathcal{V}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiopen set of Y for each $\mathcal{U}^{\mu\gamma}$ -fuzzy (u, v) -open set D of X ,
- (3) *double pairwise $(r, s)(u, v)$ -semiclosed* if $f(C)$ is a $(\mathcal{V}^{\mu\gamma}, \mathcal{W}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosed set of Y for each $\mathcal{T}^{\mu\gamma}$ -fuzzy (r, s) -closed set C of X and $f(D)$ is a $(\mathcal{W}^{\mu\gamma}, \mathcal{V}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiclosed set of Y for each $\mathcal{U}^{\mu\gamma}$ -fuzzy (u, v) -closed set D of X .

It is obvious that every double pairwise $(r, s)(u, v)$ -continuous mapping is a double pairwise $(r, s)(u, v)$ -semicontinuous mapping but the converse need not be true which is shown by the following example.

EXAMPLE 3.5. Let $X = \{x, y\}$ and let A_1, A_2, A_3 and A_4 be intuitionistic fuzzy sets of X defined as

$$A_1(x) = (0.0, 0.7), \quad A_1(y) = (0.4, 0.3);$$

$$A_2(x) = (0.5, 0.2), \quad A_2(y) = (0.6, 0.1);$$

$$A_3(x) = (0.1, 0.4), \quad A_3(y) = (0.7, 0.1);$$

and

$$A_4(x) = (0.6, 0.1), \quad A_4(y) = (0.8, 0.0).$$

Define $\mathcal{T}^{\mu\gamma} : I(X) \rightarrow I \otimes I$ and $\mathcal{U}^{\mu\gamma} : I(X) \rightarrow I \otimes I$ by

$$\mathcal{T}^{\mu\gamma}(A) = (\mathcal{T}^\mu(A), \mathcal{T}^\gamma(A)) = \begin{cases} (1, 0) & \text{if } A = 0_\sim, 1_\sim, \\ (\frac{1}{2}, \frac{1}{5}) & \text{if } A = A_1, \\ (0, 1) & \text{otherwise;} \end{cases}$$

and

$$\mathcal{U}^{\mu\gamma}(A) = (\mathcal{U}^\mu(A), \mathcal{U}^\gamma(A)) = \begin{cases} (1, 0) & \text{if } A = 0_\sim, 1_\sim, \\ (\frac{1}{3}, \frac{1}{4}) & \text{if } A = A_2, \\ (0, 1) & \text{otherwise.} \end{cases}$$

Then clearly $(X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ is a double bitopological space on X . Define $\mathcal{V}^{\mu\gamma} : I(X) \rightarrow I \otimes I$ and $\mathcal{W}^{\mu\gamma} : I(X) \rightarrow I \otimes I$ by

$$\mathcal{V}^{\mu\gamma}(A) = (\mathcal{V}^\mu(A), \mathcal{V}^\gamma(A)) = \begin{cases} (1, 0) & \text{if } A = 0_\sim, 1_\sim, \\ (\frac{1}{2}, \frac{1}{5}) & \text{if } A = A_3, \\ (0, 1) & \text{otherwise;} \end{cases}$$

and

$$\mathcal{W}^{\mu\gamma}(A) = (\mathcal{W}^\mu(A), \mathcal{W}^\gamma(A)) = \begin{cases} (1, 0) & \text{if } A = 0_\sim, 1_\sim, \\ (\frac{1}{3}, \frac{1}{4}) & \text{if } A = A_4, \\ (0, 1) & \text{otherwise.} \end{cases}$$

Then clearly $(X, \mathcal{V}^{\mu\gamma}, \mathcal{W}^{\mu\gamma})$ is a double bitopological space on X . Consider the identity mapping $1_X : (X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma}) \rightarrow (X, \mathcal{V}^{\mu\gamma}, \mathcal{W}^{\mu\gamma})$. Then it is a double pairwise $(\frac{1}{2}, \frac{1}{5})(\frac{1}{3}, \frac{1}{4})$ -semicontinuous mapping which is not a double pairwise $(\frac{1}{2}, \frac{1}{5})(\frac{1}{3}, \frac{1}{4})$ -continuous mapping.

THEOREM 3.6. Let $f : (X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma}) \rightarrow (Y, \mathcal{V}^{\mu\gamma}, \mathcal{W}^{\mu\gamma})$ be a mapping and $(r, s), (u, v) \in I \otimes I$. Then the following statements are equivalent:

- (1) f is a double pairwise $(r, s)(u, v)$ -semicontinuous mapping.
- (2) $f^{-1}(A)$ is a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosed set of X for each $\mathcal{V}^{\mu\gamma}$ -fuzzy (r, s) -closed set A of Y and $f^{-1}(B)$ is a $(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiclosed set of X for each $\mathcal{W}^{\mu\gamma}$ -fuzzy (u, v) -closed set B of Y .

(3) For each intuitionistic fuzzy set C of X ,

$$f((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(C, (r, s), (u, v))) \subseteq \mathcal{V}^{\mu\gamma}\text{-cl}(f(C), (r, s))$$

and

$$f((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(C, (u, v), (r, s))) \subseteq \mathcal{W}^{\mu\gamma}\text{-cl}(f(C), (u, v)).$$

(4) For each intuitionistic fuzzy set A of Y ,

$$(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(f^{-1}(A), (r, s), (u, v)) \subseteq f^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(A, (r, s)))$$

and

$$(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(f^{-1}(A), (u, v), (r, s)) \subseteq f^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(A, (u, v))).$$

(5) For each intuitionistic fuzzy set A of Y ,

$$f^{-1}(\mathcal{V}^{\mu\gamma}\text{-int}(A, (r, s))) \subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (r, s), (u, v))$$

and

$$f^{-1}(\mathcal{W}^{\mu\gamma}\text{-int}(A, (u, v))) \subseteq (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (u, v), (r, s)).$$

Proof. (1) \Rightarrow (2) It is obvious.

(2) \Rightarrow (3) Let C be any intuitionistic fuzzy set of X . Then $f(C)$ is an intuitionistic fuzzy set of Y , and hence $\mathcal{V}^{\mu\gamma}\text{-cl}(f(C), (r, s))$ is $\mathcal{V}^{\mu\gamma}$ -fuzzy (r, s) -closed and $\mathcal{W}^{\mu\gamma}\text{-cl}(f(C), (u, v))$ is $\mathcal{W}^{\mu\gamma}$ -fuzzy (u, v) -closed in Y . By (2), we have $f^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(f(C), (r, s)))$ is a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiclosed set and $f^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(f(C), (u, v)))$ is a $(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiclosed set of X . Also,

$$C \subseteq f^{-1}f(C) \subseteq f^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(f(C), (r, s)))$$

and

$$C \subseteq f^{-1}f(C) \subseteq f^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(f(C), (u, v))).$$

Thus

$$\begin{aligned} & (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(C, (r, s), (u, v)) \\ & \subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(f^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(f(C), (r, s))), (r, s), (u, v)) \\ & = f^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(f(C), (r, s))) \end{aligned}$$

and

$$\begin{aligned} & (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(C, (u, v), (r, s)) \\ & \subseteq (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(f^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(f(C), (u, v))), (u, v), (r, s)) \\ & = f^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(f(C), (u, v))). \end{aligned}$$

Hence

$$\begin{aligned} f((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(C, (r, s), (u, v))) &\subseteq ff^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(f(C), (r, s))) \\ &\subseteq \mathcal{V}^{\mu\gamma}\text{-cl}(f(C), (r, s)) \end{aligned}$$

and

$$\begin{aligned} f((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(C, (u, v), (r, s))) &\subseteq ff^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(f(C), (u, v))) \\ &\subseteq \mathcal{W}^{\mu\gamma}\text{-cl}(f(C), (u, v)). \end{aligned}$$

(3) \Rightarrow (4) Let A be any intuitionistic fuzzy set of Y . Then $f^{-1}(A)$ is an intuitionistic fuzzy set of X . By (3), we have

$$\begin{aligned} f((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(f^{-1}(A), (r, s), (u, v))) &\subseteq \mathcal{V}^{\mu\gamma}\text{-cl}(ff^{-1}(A), (r, s)) \\ &\subseteq \mathcal{V}^{\mu\gamma}\text{-cl}(A, (r, s)) \end{aligned}$$

and

$$\begin{aligned} f((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(f^{-1}(A), (u, v), (r, s))) &\subseteq \mathcal{W}^{\mu\gamma}\text{-cl}(ff^{-1}(A), (u, v)) \\ &\subseteq \mathcal{W}^{\mu\gamma}\text{-cl}(A, (u, v)). \end{aligned}$$

Thus

$$\begin{aligned} &(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(f^{-1}(A), (r, s), (u, v)) \\ &\subseteq f^{-1}f((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(f^{-1}(A), (r, s), (u, v))) \\ &\subseteq f^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(A, (r, s))) \end{aligned}$$

and

$$\begin{aligned} &(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(f^{-1}(A), (u, v), (r, s)) \\ &\subseteq f^{-1}f((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(f^{-1}(A), (u, v), (r, s))) \\ &\subseteq f^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(A, (u, v))). \end{aligned}$$

(4) \Rightarrow (5) Let A be any intuitionistic fuzzy set of Y . By (4),

$$(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(f^{-1}(A)^c, (r, s), (u, v)) \subseteq f^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(A^c, (r, s)))$$

and

$$(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(f^{-1}(A)^c, (u, v), (r, s)) \subseteq f^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(A^c, (u, v))).$$

By Theorem 3.3, we have

$$\begin{aligned} f^{-1}(\mathcal{V}^{\mu\gamma}\text{-int}(A, (r, s))) &= (f^{-1}(\mathcal{V}^{\mu\gamma}\text{-cl}(A^c, (r, s))))^c \\ &\subseteq ((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dscl}(f^{-1}(A)^c, (r, s), (u, v)))^c \\ &= (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (r, s), (u, v)) \end{aligned}$$

and

$$\begin{aligned} f^{-1}(\mathcal{W}^{\mu\gamma}\text{-int}(A, (u, v))) &= (f^{-1}(\mathcal{W}^{\mu\gamma}\text{-cl}(A^c, (u, v))))^c \\ &\subseteq ((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dscl}(f^{-1}(A)^c, (u, v), (r, s)))^c \\ &= (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (u, v), (r, s)). \end{aligned}$$

(5) \Rightarrow (1) Let A be any $\mathcal{V}^{\mu\gamma}$ -fuzzy (r, s) -open set and B any $\mathcal{W}^{\mu\gamma}$ -fuzzy (u, v) -open set of Y . Then $A = \mathcal{V}^{\mu\gamma}\text{-int}(A, (r, s))$ and $B = \mathcal{W}^{\mu\gamma}\text{-int}(B, (u, v))$. By (5)

$$\begin{aligned} f^{-1}(A) &= f^{-1}(\mathcal{V}^{\mu\gamma}\text{-int}(A, (r, s))) \\ &\subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (r, s), (u, v)) \\ &\subseteq f^{-1}(A) \end{aligned}$$

and

$$\begin{aligned} f^{-1}(B) &= f^{-1}(\mathcal{W}^{\mu\gamma}\text{-int}(B, (u, v))) \\ &\subseteq (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(f^{-1}(B), (u, v), (r, s)) \\ &\subseteq f^{-1}(B). \end{aligned}$$

Thus

$$f^{-1}(A) = (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (r, s), (u, v))$$

and

$$f^{-1}(B) = (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(f^{-1}(B), (u, v), (r, s)).$$

Hence $f^{-1}(A)$ is a $(\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})$ -double $(r, s)(u, v)$ -semiopen set and $f^{-1}(B)$ is an $(\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})$ -double $(u, v)(r, s)$ -semiopen set of X . Therefore f is a double pairwise $(r, s)(u, v)$ -semicontinuous mapping. \square

THEOREM 3.7. *Let $f : (X, \mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma}) \rightarrow (Y, \mathcal{V}^{\mu\gamma}, \mathcal{W}^{\mu\gamma})$ be a bijection and $(r, s), (u, v) \in I \otimes I$. Then f is a double pairwise $(r, s)(u, v)$ -semicontinuous mapping if and only if for each intuitionistic fuzzy set C of X ,*

$$\mathcal{V}^{\mu\gamma}\text{-int}(f(C), (r, s)) \subseteq f((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(C, (r, s), (u, v)))$$

and

$$\mathcal{W}^{\mu\gamma}\text{-int}(f(C), (u, v)) \subseteq f((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(C, (u, v), (r, s))).$$

Proof. Let C be any intuitionistic fuzzy set of X . Since f is one-to-one,

$$\begin{aligned} f^{-1}(\mathcal{V}^{\mu\gamma}\text{-int}(f(C), (r, s))) &\subseteq (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}f(C), (r, s), (u, v)) \\ &= (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(C, (r, s), (u, v)) \end{aligned}$$

and

$$\begin{aligned} f^{-1}(\mathcal{W}^{\mu\gamma}\text{-int}(f(C), (u, v))) &\subseteq (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(f^{-1}f(C), (u, v), (r, s)) \\ &= (\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(C, (u, v), (r, s)). \end{aligned}$$

Since f is onto, we have

$$\begin{aligned} \mathcal{V}^{\mu\gamma}\text{-int}(f(C), (r, s)) &= ff^{-1}(\mathcal{V}^{\mu\gamma}\text{-int}(f(C), (r, s))) \\ &\subseteq f((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(C, (r, s), (u, v))) \end{aligned}$$

and

$$\begin{aligned} \mathcal{W}^{\mu\gamma}\text{-int}(f(C), (u, v)) &= ff^{-1}(\mathcal{W}^{\mu\gamma}\text{-int}(f(C), (u, v))) \\ &\subseteq f((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(C, (u, v), (r, s))). \end{aligned}$$

Conversely, let A be any intuitionistic fuzzy set of Y . Since f is onto,

$$\begin{aligned} \mathcal{V}^{\mu\gamma}\text{-int}(A, (r, s)) &= \mathcal{V}^{\mu\gamma}\text{-int}(ff^{-1}(A), (r, s)) \\ &\subseteq f((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (r, s), (u, v))) \end{aligned}$$

and

$$\begin{aligned} \mathcal{W}^{\mu\gamma}\text{-int}(A, (u, v)) &= \mathcal{W}^{\mu\gamma}\text{-int}(ff^{-1}(A), (u, v)) \\ &\subseteq f((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (u, v), (r, s))). \end{aligned}$$

since f is one-to-one, we have

$$\begin{aligned} f^{-1}(\mathcal{V}^{\mu\gamma}\text{-int}(A, (r, s))) &\subseteq f^{-1}f((\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (r, s), (u, v))) \\ &= (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (r, s), (u, v)) \end{aligned}$$

and

$$\begin{aligned} f^{-1}(\mathcal{W}^{\mu\gamma}\text{-int}(A, (u, v))) &\subseteq f^{-1}f((\mathcal{U}^{\mu\gamma}, \mathcal{T}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (u, v), (r, s))) \\ &= (\mathcal{T}^{\mu\gamma}, \mathcal{U}^{\mu\gamma})\text{-dsint}(f^{-1}(A), (u, v), (r, s)). \end{aligned}$$

Hence the theorem follows. □

References

- [1] K. T. Atanassov, *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems **20** (1986), 87-96.
- [2] C. L. Chang, *Fuzzy topological spaces*, J. Math. Anal. Appl. **24** (1968), 182-190.
- [3] K. C. Chattopadhyay, R. N. Hazra, and S. K. Samanta, *Gradation of openness : Fuzzy topology*, Fuzzy Sets and Systems **49** (1992), 237-242.
- [4] D. Çoker, *An introduction to intuitionistic fuzzy topological spaces*, Fuzzy Sets and Systems **88** (1997), 81-89.
- [5] D. Çoker and M. Demirci, *An introduction to intuitionistic fuzzy topological spaces in Šostak's sense*, BUSEFAL **67** (1996), 67-76.

- [6] D. Çoker and A. Haydar Eş, *On fuzzy compactness in intuitionistic fuzzy topological spaces*, J. Fuzzy Math. **3** (1995), 899-909.
- [7] H. Gürçay, D. Çoker, and A. Haydar Eş, *On fuzzy continuity in intuitionistic fuzzy topological spaces*, J. Fuzzy Math. **5** (1997), 365-378.
- [8] A. Kandil, *Biproximities and fuzzy bitopological spaces*, Simon Stevin **63** (1989), 45-66.
- [9] E. P. Lee and S. O. Lee, *Fuzzy intuitionistic almost (r, s) -continuous mappings*, Journal of The Chungcheong Mathematical Society **26** (2013), 125-135.
- [10] E. P. Lee and S. O. Lee, *Double semiopen sets on double bitopological spaces*, Journal of The Chungcheong Mathematical Society **26** (2013), 691-702.
- [11] S. O. Lee and E. P. Lee, *Fuzzy pairwise (r, s) -irresolute mappings*, International Journal of Fuzzy Logic and Intelligent Systems **9** (2009), 105-109.
- [12] A. A. Ramadan, *Smooth topological spaces*, Fuzzy Sets and Systems **48** (1992), 371-375.
- [13] A. P. Sostak, *On a fuzzy topological structure*, Suppl. Rend. Circ. Matem. Janos Palermo, Sr. II **11** (1985), 89-103.

*

Department of Clinical Laboratory Science
Seonam University
Namwon 590-711, Republic of Korea
E-mail: eplee@seonam.ac.kr

**

Department of Mathematics
Chungbuk National University
Cheongju 361-763, Republic of Korea
E-mail: solee@chungbuk.ac.kr