

## ON HENSTOCK INTEGRALS OF INTERVAL-VALUED FUNCTIONS ON TIME SCALES

WON TAE OH\* AND JU HAN YOON\*\*

ABSTRACT. In this paper we introduce the interval-valued Henstock integral on time scales and investigate some properties of these integrals.

### 1. Introduction and preliminaries

The Henstock integral for real functions was first defined by Henstock [2] in 1963. The Henstock integral is more powerful and simpler than the Lebesgue, Wiener and Feynman integrals. The Henstock delta integral on time scales was introduced by Allan Peterson and Bevan Thompson [5] in 2006. In 2000, Congxin Wu and Zengtai Gong introduced the concept of the Henstock integral of interval-valued functions [6].

In this paper we introduce the concept of the Henstock delta integral of interval-valued function on time scales and investigate some properties of the integral.

A time scale  $T$  is a nonempty closed subset of real number  $\mathbb{R}$  with the subspace topology inherited from the standard topology of  $\mathbb{R}$ . For  $t \in T$  we define the forward jump operator  $\sigma(t) = \inf\{s \in T : s > t\}$  where  $\inf \phi = \sup\{T\}$ , while the backward jump operator  $\rho(t) = \sup\{s \in T : s < t\}$  where  $\sup \phi = \inf\{T\}$ . If  $\sigma(t) > t$ , we say that  $t$  is right-scattered, while if  $\rho(t) < t$ , we say that  $t$  is left-scattered. If  $\sigma(t) = t$ , we say that  $t$  is right-dense, while if  $\rho(t) = t$ , we say that  $t$  is left-dense. The forward graininess function  $\mu(t)$  of  $t \in T$  is defined by  $\mu(t) = \sigma(t) - t$ , while the backward graininess function  $\nu(t)$  of  $t \in T$  is defined by  $\nu(t) = t - \rho(t)$ . For  $a, b \in T$  we denote the closed interval  $[a, b]_T = \{t \in T : a \leq t \leq b\}$ .

---

Received October 17, 2014; Accepted November 10, 2014.

2010 Mathematics Subject Classification: Primary 26A39; secondary 26E70.

Key words and phrases: Henstock delta integral, time scales.

Correspondence should be addressed to Won Tae Oh, [wntoh@chungbuk.ac.kr](mailto:wntoh@chungbuk.ac.kr).

This work was supported by the research grant of the Chungbuk National University in 2012.

$\delta = (\delta_L, \delta_R)$  is a  $\Delta$ -gauge on  $[a, b]_T$  if  $\delta_L(t) > 0$  on  $(a, b]_T$ ,  $\delta_R(t) > 0$  on  $(a, b)_T$ ,  $\delta_L(a) \geq 0$ ,  $\delta_R(b) \geq 0$  and  $\delta_R(t) \geq \mu(t)$  for each  $t \in [a, b]_T$ .

A collection  $P = \{([t_{i-1}, t_i], \xi_i) : 1 \leq i \leq n\}$  of tagged intervals is  $\delta$ -fine Henstock partition of  $[a, b]_T$  if  $U_{i=1}^n [t_{i-1}, t_i] = [a, b]_T$ ,  $[t_{i-1}, t_i]_T \subset [\xi_i - \delta_L(\xi_i), \xi_i + \delta_R(\xi_i)]$  and  $\xi_i \in [t_{i-1}, t_i]_T$  for each  $i = 1, 2, \dots, n$ .

DEFINITION 1.1 ([5]). A function  $f : [a, b] \rightarrow \mathbb{R}$  is Henstock delta integrable (or  $H_\Delta$ -integrable) on  $[a, b]$  if there exists a number  $A$  such that for each  $\epsilon > 0$  there exists a  $\Delta$ -gauge  $\delta$  on  $[a, b]$  such that

$$\left| \sum_{i=1}^n f(\xi_i)(t_i - t_{i-1}) - A \right| < \epsilon$$

for every  $\delta$ -fine Henstock partition  $P = \{([t_{i-1}, t_i], \xi_i) : 1 \leq i \leq n\}$  of  $[a, b]$ . The number  $A$  is called the Henstock delta integral of  $f$  on  $[a, b]$  and we write  $A = (H_\Delta) \int_a^b f$ .

DEFINITION 1.2. Let  $I_\mathbb{R} = \{I = [I^-, I^+]$  is the closed bounded interval on the real  $\mathbb{R}\}$ , where  $I^- = \min\{x : x \in I\}$ ,  $I^+ = \max\{x : x \in I\}$ . For  $A, B, C \in I_\mathbb{R}$ , we define  $A \leq B$  iff  $A^- \leq B^-$  and  $A^+ \leq B^+$ ,  $A + B = C$  iff  $A^- + B^- = C^-$  and  $A^+ + B^+ = C^+$ , and  $AB = \{ab : a \in A, b \in B\}$ , where  $(AB)^- = \min(A^-B^-, A^-B^+, A^+B^-, A^+B^+)$  and  $(AB)^+ = \max(A^-B^-, A^-B^+, A^+B^-, A^+B^+)$ . Define  $d(A, B) = \max(|A^- - B^-|, |A^+ - B^+|)$  as the distance between  $A$  and  $B$ .

DEFINITION 1.3 ([6]). An interval-valued function  $F : [a, b] \rightarrow I_\mathbb{R}$  is Henstock integrable to  $I_0 \in I_\mathbb{R}$  on  $[a, b]$  if for every  $\epsilon > 0$  there exists a gauge  $\delta$  on  $[a, b]$  such that

$$d\left(\sum_{i=1}^n F(\xi_i)(t_i - t_{i-1}), I_0\right) < \epsilon$$

whenever  $P = \{([t_{i-1}, t_i], \xi_i) : 1 \leq i \leq n\}$  of  $[a, b]$  is a  $\delta$ -fine Henstock partition of  $[a, b]$ . We write  $(IH) \int_a^b F(x)dx = I_0$  and  $F \in IH[a, b]$ .

## 2. The interval-valued Henstock delta integral on time scales

In this section, we will define the Henstock integral of interval-valued function on time scales and investigate some properties of the integral.

DEFINITION 2.1. An interval-valued function  $F : [a, b]_T \rightarrow I_\mathbb{R}$  is Henstock delta integrable to  $I_0 \in I_\mathbb{R}$  on  $[a, b]_T$  if for every  $\epsilon > 0$  there

exists a  $\Delta$ -gauge  $\delta$  on  $[a, b]_T$  such that

$$d\left(\sum_{i=1}^n F(\xi_i)(t_i - t_{i-1}), I_0\right) < \epsilon$$

whenever  $P = \{([t_{i-1}, t_i]_T, \xi_i) : 1 \leq i \leq n\}$  is a  $\delta$ -fine Henstock partition of  $[a, b]_T$ . We write  $(IH_\Delta) \int_a^b F(x)dx = I_0$  and  $F \in IH_\Delta[a, b]_T$ .

REMARK 2.2. It is clear, if  $F(x) = F^-(x) = F^+(x)$  for all  $x \in [a, b]$ , then Definition 2.1 implies the real-valued Henstock integral on  $[a, b]$ .

REMARK 2.3. If  $F \in IH_\Delta[a, b]_T$ , then the integral is unique.

THEOREM 2.4. An interval-valued function  $F : [a, b]_T \rightarrow I_{\mathbb{R}}$  is Henstock delta integrable on  $[a, b]_T$  if and only if  $F^-, F^+ \in H_\Delta[a, b]_T$  and

$$(IH_\Delta) \int_a^b F(x)dx = \left[ (H_\Delta) \int_a^b F^-(x)dx, (H_\Delta) \int_a^b F^+(x)dx \right],$$

where  $F(x) = [F^-(x), F^+(x)]$ .

*Proof.* Let  $F \in IH_\Delta[a, b]_T$ . Then there exists an interval  $I_0 = [I_0^-, I_0^+]$  with the property that for each  $\epsilon > 0$  there exists a  $\Delta$ -gauge  $\delta$  on  $[a, b]_T$  such that

$$d\left(\sum_{i=1}^n F(\xi_i)(t_i - t_{i-1}), I_0\right) < \epsilon$$

whenever  $P = \{([t_{i-1}, t_i]_T, \xi_i) : 1 \leq i \leq n\}$  is a  $\delta$ -fine Henstock partition of  $[a, b]_T$ .

Let  $P = \{([t_{i-1}, t_i]_T, \xi_i) : 1 \leq i \leq n\}$  be a  $\delta$ -fine Henstock partition of  $[a, b]_T$ . Since

$$\begin{aligned} & d\left(\sum_{i=1}^n F(\xi_i)(t_i - t_{i-1}), I_0\right) \\ &= \max\left(\left|\sum_{i=1}^n F^-(\xi_i)(t_i - t_{i-1}) - I_0^-\right|, \left|\sum_{i=1}^n F^+(\xi_i)(t_i - t_{i-1}) - I_0^+\right|\right), \\ & \left|\sum_{i=1}^n F^-(\xi_i)(t_i - t_{i-1}) - I_0^-\right| < \epsilon, \left|\sum_{i=1}^n F^+(\xi_i)(t_i - t_{i-1}) - I_0^+\right| < \epsilon. \end{aligned}$$

Conversely, let  $F^-, F^+ \in H_\Delta[a, b]_T$ . then there exist  $H_1, H_2 \in \mathbb{R}$  with the property that given  $\Delta$ -gauge  $\delta$  on  $[a, b]_T$  such that

$$\left| \sum_{i=1}^n F^-(\xi_i)(t_i - t_{i-1}) - H_1 \right| < \epsilon, \left| \sum_{i=1}^n F^+(\xi_i)(t_i - t_{i-1}) - H_2 \right| < \epsilon.$$

whenever  $P = \{([t_{i-1}, t_i]_T, \xi_i) : 1 \leq i \leq n\}$  is a  $\delta$ -fine Henstock partition of  $[a, b]_T$ . We define  $I_0 = [H_1, H_2]$ , then if  $P = \{([t_{i-1}, t_i]_T, \xi_i) : 1 \leq i \leq n\}$  is a  $\delta$ -fine Henstock partition of  $[a, b]_T$ . We have

$$d\left(\sum_{i=1}^n F(\xi_i)(t_i - t_{i-1}), I_0\right) < \epsilon.$$

Hence  $F : [a, b]_T \rightarrow I_{\mathbb{R}}$  is Henstock delta integrable on  $[a, b]_T$ . □

From Theorem 2.4 and the properties of Henstock delta integral ([6]), we can easily obtain the following theorems.

**THEOREM 2.5.** *Let  $F, G \in IH_\Delta[a, b]_T$  and  $\beta, \gamma \in \mathbb{R}$ . Then*

- (1)  $\beta F + \gamma G \in IH_\Delta[a, b]_T$  and  $(IH_\Delta) \int_a^b (\beta F + \gamma G) dx = \beta (IH_\Delta) \int_a^b F dx + \gamma (IH_\Delta) \int_a^b G dx$
- (2) If  $F(x) \leq G(x)$  a.e. in  $[a, b]_T$ , then  $(IH_\Delta) \int_a^b F dx \leq (IH_\Delta) \int_a^b G dx$

**THEOREM 2.6.** *Let  $F \in IH_\Delta[a, c]_T$  and  $F \in IH_\Delta[c, b]_T$ . Then  $F \in IH_\Delta[a, b]_T$  and*

$$(IH_\Delta) \int_a^b F dx = (IH_\Delta) \int_a^c F dx + (IH_\Delta) \int_c^b F dx$$

**THEOREM 2.7.** *Let  $F, G \in IH_\Delta[a, b]_T$  and  $d(F, G)$  is Lebesgue delta integrable on  $[a, b]_T$ . Then*

$$d\left((IH_\Delta) \int_a^b F dx, (IH_\Delta) \int_a^b G dx\right) \leq (L_\Delta) \int_a^b d(F, G) dx$$

*Proof.* By definition of distance, we have

$$\begin{aligned}
 & d\left( (IH_\Delta) \int_a^b F dx, (IH_\Delta) \int_a^b G dx \right) \\
 &= \max \left( \left| \left( (IH_\Delta) \int_a^b F dx \right)^- - \left( (IH_\Delta) \int_a^b G dx \right)^- \right|, \right. \\
 &\quad \left. \left| \left( (IH_\Delta) \int_a^b F dx \right)^+ - \left( (IH_\Delta) \int_a^b G dx \right)^+ \right| \right) \\
 &= \max \left( \left| (IH_\Delta) \int_a^b (F^- - G^-) dx \right|, \left| (IH_\Delta) \int_a^b (F^+ - G^+) dx \right| \right) \\
 &= \max \left( (L_\Delta) \int_a^b |F^- - G^-| dx, (L_\Delta) \int_a^b |F^+ - G^+| dx \right) \\
 &\leq (L_\Delta) \int_a^b d(F, G) dx.
 \end{aligned}$$

□

### 3. The Henstock delta integral of fuzzy number valued functions

DEFINITION 3.1 ([1]). Let  $\tilde{A} \in F(\mathbb{R})$  be a fuzzy subset on  $\mathbb{R}$ . If for any  $\lambda \in [0, 1]$ ,  $A_\lambda = [A_\lambda^-, A_\lambda^+]$  and  $A_1 \neq \emptyset$ , where  $A_\lambda = \{x : \tilde{A}(x) \geq \lambda\}$ , then  $\tilde{A}$  is called a fuzzy number.

Let  $\tilde{\mathbb{R}}$  denote the set of all fuzzy numbers.

DEFINITION 3.2 ([3]). Let  $\tilde{A}, \tilde{B} \in \tilde{\mathbb{R}}$ , we define  $\tilde{A} \leq \tilde{B}$  iff  $A_\lambda \leq B_\lambda$  for all  $\lambda \in (0, 1]$ ,  $\tilde{A} + \tilde{B} = \tilde{C}$  iff  $A_\lambda + B_\lambda = C_\lambda$  for any  $\lambda \in (0, 1]$ ,  $\tilde{A} \cdot \tilde{B} = \tilde{D}$  iff  $A_\lambda \cdot B_\lambda = D_\lambda$  for any  $\lambda \in (0, 1]$ . For  $D(\tilde{A}, \tilde{B}) = \sup_{\lambda \in [0,1]} d(A_\lambda, B_\lambda)$  is called the distance between  $\tilde{A}, \tilde{B}$ .

LEMMA 3.3 ([1]). If a mapping  $H : [0, 1] \rightarrow I_{\mathbb{R}}, \lambda \mapsto H(\lambda) = [m_\lambda, n_\lambda]$ , satisfies  $[m_{\lambda_1}, n_{\lambda_1}] \supset [m_{\lambda_2}, n_{\lambda_2}]$  when  $\lambda_1 < \lambda_2$ , then

$$\tilde{A} := \bigcup_{\lambda \in (0,1]} \lambda H(\lambda) \in \tilde{\mathbb{R}}$$

and

$$A_\lambda = \bigcap_{n=1}^{\infty} H(\lambda_n),$$

where  $\lambda_n = [1 - 1/(n + 1)]\lambda$ .

DEFINITION 3.4. Let  $\tilde{F} : [a, b]_T \rightarrow \tilde{\mathbb{R}}$ . If the interval-valued function  $F_\lambda(x) = [F_\lambda^-(x), F_\lambda^+(x)]$  is Henstock delta integrable on  $[a, b]_T$  for any  $\lambda \in (0, 1]$ , then we say that  $\tilde{F}(x)$  is Henstock delta integrable on  $[a, b]_T$  and the integral is defined by Henstock delta integral is defined by

$$\begin{aligned} (FH_\Delta) \int_a^b \tilde{F}(x)dx &:= \bigcup_{\lambda \in (0,1]} \lambda (IH_\Delta) \int_a^b F_\lambda(x)dx \\ &= \bigcup_{\lambda \in (0,1]} \lambda \left[ (H_\Delta) \int_a^b F_\lambda^- dx, (H_\Delta) \int_a^b F_\lambda^+ dx \right]. \end{aligned}$$

We will write  $\tilde{F} \in FH_\Delta[a, b]_T$ .

THEOREM 3.5.  $\tilde{F} \in FH_\Delta[a, b]_T$ , then  $(FH_\Delta) \int_a^b \tilde{F}(x)dx \in \tilde{\mathbb{R}}$  and

$$\left[ (FH_\Delta) \int_a^b \tilde{F}(x)dx \right]_\lambda = \bigcap_{n=1}^\infty (IH_\Delta) \int_a^b F_{\lambda_n}(x)dx,$$

where  $\lambda_n = [1 - 1/(n + 1)]\lambda$ .

*Proof.* Let  $H : (0, 1] \rightarrow I_{\mathbb{R}}$  be defined by

$$H(\lambda) = \left[ (H_\Delta) \int_a^b F_\lambda^-(x)dx, (H_\Delta) \int_a^b F_\lambda^+(x)dx \right].$$

Since  $F_\lambda^-(x)$  and  $F_\lambda^+(x)$  are increasing and decreasing on  $\lambda$ , respectively, therefore, when  $0 < \lambda_1 \leq \lambda_2 \leq 1$  we have  $F_{\lambda_1}^-(x) \leq F_{\lambda_2}^-(x)$ ,  $F_{\lambda_1}^+(x) \geq F_{\lambda_2}^+(x)$  on  $[a, b]_T$ . Thus from Theorem 2.5, we have

$$\begin{aligned} &\left[ (H_\Delta) \int_a^b F_{\lambda_1}^-(x)dx, (H_\Delta) \int_a^b F_{\lambda_1}^+(x)dx \right] \\ &\supset \left[ (H_\Delta) \int_a^b F_{\lambda_2}^-(x)dx, (H_\Delta) \int_a^b F_{\lambda_2}^+(x)dx \right]. \end{aligned}$$

Using Theorem 2.5 and Lemma 3.3 we obtain

$$(IH_\Delta) \int_a^b \tilde{F}(x)dx := \bigcup_{\lambda \in (0,1]} \lambda \left[ (H_\Delta) \int_a^b F_\lambda^-(x)dx, (H_\Delta) \int_a^b F_\lambda^+(x)dx \right] \in \tilde{\mathbb{R}}$$

and for all  $\lambda \in (0, 1]$ ,

$$\left[ (FH_\Delta) \int_a^b \tilde{F}(x)dx \right]_\lambda = \bigcap_{n=1}^\infty (IH_\Delta) \int_a^b F_{\lambda_n}(x)dx,$$

where  $\lambda_n = [1 - 1/(n + 1)]\lambda$ .  $\square$

Using Theorem 3.5 and the properties of  $(IH)$  integral, we can obtain the properties of  $(FH_\Delta)$  integral. For examples, we get the linearity, monotonicity and interval additivity properties of  $(FH_\Delta)$  integral.

### References

- [1] L. Chengzhong, *Extension of the integral of interval-valued function and integral of fuzzy-valued functions*, Fuzzy Math. **3** (1983), 45-52.
- [2] R. Henstock, *Theory of integration*, Butterworths, London, 1963.
- [3] S. Nada, *On integration of fuzzy mapping*, Fuzzy Sets and Systems **32** (2000), 377-392.
- [4] J. M. Park, Y. K. Kim, D. H. Lee, J. H. Yoon, and J. T. Lim, *Convergence Theorems for the Henstock delta integral on time scales*, Journal of the ChungCheoug Math. Soc. **26** (2013), 880-884.
- [5] A. Peterson and B. Thompson, *Henstock-Kurzweil Delta and Nabla Integral*, J. Math. Anal. Appl. **323** (2006), 162-178.
- [6] C. Wu and Z. Gong, *On Henstock integrals of interval-valued functions and fuzzy-valued functions*, Fuzzy Set and System **115** (2000), 377-392.

\*

Department of Mathematics  
Chungbuk National University  
Cheongju 361-763, Republic of Korea  
*E-mail*: wntoh@chungbuk.ac.kr

\*\*

Department of Mathematics Education  
Chungbuk National University  
Cheongju 361-763, Republic of Korea  
*E-mail*: yoonjh@cbnu.ac.kr