

## A SEXTIC-ORDER SIMPLE-ROOT FINDER WITH RATIONAL WEIGHTING FUNCTIONS OF DERIVATIVE-TO-DERIVATIVE RATIOS

YOUNG IK KIM\*

ABSTRACT. A three-step sextic simple-root finder is constructed with the use of weighting functions of derivative-to-derivative ratios. Their convergence and computational properties are investigated along with concrete numerical examples to verify the theoretical analysis.

### 1. Introduction

High-order multipoint root-finders have been developed for a given nonlinear equation  $f(x) = 0$ , some of which can be found in [7, 11, 13]. Simple-root finders using more than two derivatives are rarely found. Jarratt[8], in 1966, suggested a multipoint iterative methods of order 4 or 5 costing three derivatives and one function below in (1.1):

$$\begin{cases} y_n = x_n + \gamma \frac{f(x_n)}{f'(x_n)}, \gamma \in \mathbb{R}, \\ z_n = x_n + \beta \frac{f(x_n)}{f'(x_n)} + \sigma \frac{f(x_n)}{f'(y_n)}, \beta, \sigma \in \mathbb{R}, \\ x_{n+1} = x_n - \frac{f(x_n)}{a_1 f'(x_n) + a_2 f'(y_n) + a_3 f'(z_n)}, a_1, a_2, a_3 \in \mathbb{R}. \end{cases} \quad (1.1)$$

DEFINITION 1.1. (Error equation, asymptotic error constant, order of convergence) Let  $x_0, x_1, \dots, x_n, \dots$  be a sequence of numbers converging to  $\alpha$ . Let  $e_n = x_n - \alpha$  for  $n = 0, 1, 2, \dots$ . If constants  $p \geq 1$ ,  $c \neq 0$  exist in such a way that  $e_{n+1} = c e_n^p + O(e_n^{p+1})$  called the *error equation*, then  $p$  and  $\eta = |c|$  are said to be the *order of convergence* and the *asymptotic error constant*, respectively. It is easy to find  $c = \lim_{n \rightarrow \infty} \frac{e_{n+1}}{e_n^p}$ . Some authors call  $c$  the asymptotic error constant.

---

Received October 22, 2014; Accepted November 10, 2014.

2010 Mathematics Subject Classification: Primary 65H05, 65H99.

Key words and phrases: Sextic-order convergence, asymptotic error constant, efficiency index, Jarratt-like.

If we select  $\alpha \neq -\frac{2}{3}, \theta \neq 0, \theta \neq \alpha$  with  $\theta = \beta + \gamma$  in (1.1), then we find a two-parameter family of fourth-order methods if  $a_1 = \frac{6\alpha\theta + 3(\alpha + \theta) + 2}{6\alpha\theta}, a_2 = \frac{3\theta + 2}{6\alpha(\alpha - \theta)}, a_3 = \frac{3\alpha + 2}{6\theta(\theta - \alpha)}, \beta = \theta - \frac{3\theta(\theta - \alpha)}{2\alpha(3\alpha + 2)}, \gamma = \frac{3\theta(\theta - \alpha)}{2\alpha(3\alpha + 2)}$  are chosen. For  $\alpha = -\frac{2}{3}$ , we have  $\theta = \alpha = -\frac{2}{3}, a_1 = \frac{1}{4}, a_2 = \frac{3}{4} + \frac{3}{8\gamma}, a_3 = -\frac{3}{8\gamma}, \beta = -\frac{2}{3} - \gamma$  and hence a one-parameter family of fourth methods are found. In particular, for  $\alpha = -1, \theta = -\frac{1}{2}, a_1 = \frac{1}{6}, a_2 = \frac{1}{6}, a_3 = \frac{2}{3}, \beta = -\frac{1}{8}, \gamma = -\frac{3}{8}$ , we obtain a fifth-order method, whose asymptotic error constant is given by:

$$c_2^4 + \frac{c_2^2 c_3}{8} - \frac{c_3^2}{4} + \frac{c_2 c_4}{2} + \frac{c_5}{24}, \quad (1.2)$$

which Jarratt did not show.

Three-point sextic-order methods with two derivatives can be found in [6, 9, 15] as well as in the works of Chun[5] and Parhi et al.[10] respectively shown in (1.3) and (1.4) below.

$$\begin{cases} y_n = x_n - \frac{2}{3} \frac{f(x_n)}{f'(x_n)}, \\ z_n = x_n - J_f(x_n) \cdot \frac{f(x_n)}{f'(x_n)}, J_f(x_n) = \frac{3f'(y_n) + f'(x_n)}{6f'(y_n) - 2f'(x_n)}, \\ x_{n+1} = z_n - \frac{f(z_n)}{a(z_n - x_n)(z_n - y_n) + \frac{3}{2}J_f(x_n)f'(y_n) + (1 - \frac{3}{2}J_f(x_n))f'(x_n)}, a \in \mathbb{R}. \end{cases} \quad (1.3)$$

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = x_n - \frac{2f(x_n)}{f'(x_n) + f'(y_n)}, \\ x_{n+1} = z_n - \frac{f'(x_n) + f'(y_n)}{3f'(y_n) - f'(x_n)} \cdot \frac{f(z_n)}{f'(x_n)}. \end{cases} \quad (1.4)$$

The main objective of this is to develop a more general class of three-point sixth-order Jarratt-like methods requiring three derivatives and one function. To this end, by modifying and extending (1.1), we consider a general family of three-point methods in the following form:

$$\begin{cases} y_n = x_n - \gamma \cdot \frac{f(x_n)}{f'(x_n)}, \gamma \in \mathbb{R}, \\ z_n = x_n - (\beta + \sigma s) \cdot \frac{f(x_n)}{f'(x_n)}, s = \frac{f'(y_n)}{f'(x_n)}, \beta, \sigma \in \mathbb{R}, \\ x_{n+1} = x_n - \frac{(1+b_1(s-1)+b_2(s-1)^2+b_3(t-1)+b_4(t-1)^2)}{(1+a_1(s-1)+a_2(s-1)^2+a_3(t-1)+a_4(t-1)^2)} \cdot \frac{f(x_n)}{f'(x_n)}, t = \frac{f'(z_n)}{f'(x_n)}, \end{cases} \quad (1.5)$$

where  $\gamma, \beta, \sigma \in \mathbb{R}, a_i, b_i (1 \leq i \leq 4)$  are parameters to be determined for sextic-order convergence. In Section 2, a successful development is described for a new family of sixth-order methods. Section 3 presents numerical experiments along with concluding remarks.

### 2. Convergence analysis

The desired conditions on parameters  $\gamma, \beta, \sigma \in \mathbb{R}$ ,  $a_i, b_i (1 \leq i \leq 4)$  will be obtained along with the proof of Theorem 2.1 describing the methodology and convergence analysis on iterative scheme (1.5).

**THEOREM 2.1.** *Assume that  $f : \mathbb{C} \rightarrow \mathbb{C}$  has a simple root  $\alpha$  and is analytic in a region containing  $\alpha$ . Let  $\Delta = f'(\alpha)$  and  $c_j = \frac{f^{(j)}(\alpha)}{j!f'(\alpha)}$  for  $j = 2, 3, \dots$ . Let  $\gamma, \beta, \sigma, a_i, b_i (1 \leq i \leq 4) \in \mathbb{R}$  be some parameters in (1.5). Let  $x_0$  be an initial guess chosen in a sufficiently small neighborhood of  $\alpha$ . Let  $\gamma = \frac{6-\sqrt{6}}{10}, \beta = \frac{6+\sqrt{6}}{10}, \sigma = -\frac{3+8\sqrt{6}}{25}$  and  $a_1 = -\frac{(41+21\sqrt{6})}{12}, a_2 = -\frac{5(38+13\sqrt{6})}{48}, a_3 = \frac{31+9\sqrt{6}}{12}, a_4 = \frac{118-23\sqrt{6}}{48}, b_1 = -\frac{(139+64\sqrt{6})}{36}, b_2 = -\frac{5(8+3\sqrt{6})}{36}, b_3 = \frac{7(11+4\sqrt{6})}{36}, b_4 = \frac{54-19\sqrt{6}}{36}$ . Then iterative scheme (1.5) defines a sextic-order simple-root finders satisfying the error equation below: for  $n = 0, 1, 2, \dots$ ,*

$$e_{n+1} = (\phi_1 c_2^5 + \phi_2 c_2^3 c_3 + \phi_3 c_2^2 c_4 + \phi_4 c_3 c_4 + \phi_5 c_2 c_3^2 + \phi_6 c_2 c_5 + \phi_7 c_6) e_n^6 + O(e_n^7), \tag{2.1}$$

where  $\phi_1 = \frac{41-19\sqrt{6}}{5}, \phi_2 = \frac{-761+454\sqrt{6}}{100}, \phi_3 = \frac{53-12\sqrt{6}}{25}, \phi_4 = -\frac{2+3\sqrt{6}}{50}, \phi_5 = -\frac{224+171\sqrt{6}}{400}, \phi_6 = \frac{-4+\sqrt{6}}{20}$  and  $\phi_7 = \frac{1}{100}$ .

*Proof.* Taylor series expansion of  $f(x_n)$  about  $\alpha$  up to 6th-order terms yields with  $f(\alpha) = 0$ :

$$f(x_n) = \Delta \{e_n + c_2 e_n^2 + c_3 e_n^3 + c_4 e_n^4 + c_5 e_n^5 + c_6 e_n^6 + O(e_n^7)\}. \tag{2.2}$$

It follows that

$$f'(x_n) = \Delta \{1 + 2c_2 e_n + 3c_3 e_n^2 + 4c_4 e_n^3 + 5c_5 e_n^4 + 6c_6 e_n^5 + 7c_7 e_n^6 + O(e_n^7)\}. \tag{2.3}$$

For brevity of notation,  $e_n$  will be denoted by  $e$  throughout the proof. With the aid of symbolic computation of Mathematica[16], we have:

$$y_n = x_n - \gamma \frac{f(x_n)}{f'(x_n)} = \alpha + (1 - \gamma)e + c_2 \gamma e^2 + 2\gamma(-c_2^2 + c_3)e^3 + Y_4 e^4 + Y_5 e^5 + Y_6 e^6 + O(e^7), \tag{2.4}$$

where  $Y_4 = \gamma(4c_2^3 - 7c_2 c_3 + 3c_4), Y_5 = -2\gamma(4c_2^4 - 10c_2^2 c_3 + 3c_3^2 + 5c_2 c_4 - 2c_5)$  and  $Y_6 = \gamma(16c_2^5 - 52c_2^3 c_3 + 33c_2 c_3^2 + 28c_2^2 c_4 - 17c_3 c_4 - 13c_2 c_5 + 5c_6)$ . In view of the fact that  $f'(y_n) = f'(x_n)|_{e_n \rightarrow (y_n - \alpha)}$ , we get:

$$f'(y_n) = \Delta [1 + (1 - \gamma)c_2 e + \{3(1 - \gamma)^2 c_3 + 2\gamma c_2^2\} e^2$$

$$+D_3e^3 + \sum_{i=4}^6 D_i e^i + O(e^7)], \tag{2.5}$$

where  $D_3 = 2\{2(1-\gamma)^3c_4 - \gamma c_2(2c_2^2 + (3\gamma - 5)c_3)\}$ ,  $D_i = D_i(\gamma, c_2, c_3, \dots, c_6)$  for  $4 \leq i \leq 6$ .

$$s = \frac{f'(y_n)}{f'(x_n)} = 1 - 2\gamma c_2 e + 3\gamma(2c_2^2 + (\gamma - 2)c_3)e^2 + \sum_{i=3}^6 E_i e^i + O(e^7), \tag{2.6a}$$

where  $E_i = E_i(\gamma, c_2, c_3, \dots, c_6)$  is a multivariate polynomial in  $c_2, c_3, \dots, c_6$  for  $3 \leq i \leq 6$ .

Hence with  $\beta = 4 - \frac{5\sqrt{6}}{3}$  and  $\sigma = -\frac{3+8\sqrt{6}}{25}$  we have:

$$z_n = x_n - (\beta + \sigma s) \cdot \frac{f(x_n)}{f'(x_n)} = (1 - \beta)e + (\beta + 2\gamma\sigma)c_2e^2 + \sum_{i=3}^6 F_i e^i + O(e^7), \tag{2.6b}$$

where  $F_i = F_i(\beta, \gamma, \sigma, c_2, c_3, \dots, c_6)$  for  $3 \leq i \leq 6$ .

In view of the fact that  $f'(z_n) = f'(x_n)|_{e_n \rightarrow (z_n - \alpha)}$ , we get:

$$f'(z_n) = \Delta[1 + 2(1 - \beta)c_2e + [3c_3(1 - \beta)^2 + 2c_2^2(\beta + 2\gamma\sigma)]e^2 + \sum_{i=3}^6 G_i e^i + O(e^7)], \tag{2.7}$$

where  $G_i = G_i(\beta, \gamma, \sigma, c_2, c_3, \dots, c_6)$  for  $3 \leq i \leq 6$ .

By direct substitution of  $z_n, f(x_n), f'(x_n), f'(y_n), f'(z_n)$  with  $s = \frac{f'(y_n)}{f'(x_n)}, t = \frac{f'(z_n)}{f'(x_n)}$  in (1.5), we find

$$x_{n+1} = x_n - \frac{(1 + b_1(s - 1) + b_2(s - 1)^2 + b_3(t - 1) + b_4(t - 1)^2) \cdot f(x_n)}{(1 + a_1(s - 1) + a_2(s - 1)^2 + a_3(t - 1) + a_4(t - 1)^2) \cdot f'(x_n)} = \alpha + (1 - 2(a_3 - b_3)\beta - 2(a_1 - b_1)\gamma)e^2 + \sum_{i=3}^6 \Gamma_i e^i + O(e^7), \tag{2.8}$$

where  $\Gamma_i = \Gamma_i(a_j, b_k, \beta, \gamma, \sigma, c_2, c_3, \dots, c_6)$  for  $3 \leq i \leq 6$  and  $1 \leq j, k \leq 4$ .

By solving  $1 - 2(a_3 - b_3)\beta - 2(a_1 - b_1)\gamma = 0$  from (2.8) for  $a_1$ , we immediately obtain

$$a_1 = b_1 - \frac{1 - 2\beta(a_3 - b_3)}{2\gamma}. \tag{2.9}$$

By substituting  $a_1$  into  $\Gamma_3 = 0$ , we find two relations independently of  $c_2$  and  $c_3$  below:

$$-2 + 6(a_3 - b_3)\beta(\beta - \gamma) + 3\gamma = 0,$$

$$1 + 4(a_4 - b_4)\beta^2 - 2(b_1\gamma + b_3\beta) + 4(a_2 - b_2)\gamma^2 + 4(a_3 - b_3)\gamma\sigma = 0.$$

As a result, we find

$$a_3 = \frac{1 + 4(a_4 - b_4)\beta^2 - 2b_1\gamma + 4(a_2 - b_2)\gamma^2}{2\beta} + \frac{(2 - 3\gamma)(\beta + 2\gamma\sigma)}{6\beta^2(\beta - \gamma)},$$

$$b_3 = \frac{1 + 4(a_4 - b_4)\beta^2 - 2b_1\gamma + 4(a_2 - b_2)\gamma^2}{2\beta} + \frac{(2 - 3\gamma)\gamma\sigma}{3\beta^2(\beta - \gamma)}. \quad (2.10)$$

By substituting  $a_1, a_3, b_3$  into  $\Gamma_4 = 0$ , we find three relations independently of  $c_2, c_3$  and  $c_4$  with  $\beta(\beta - \gamma) \neq 0$  below:

$$\begin{aligned} & 3 - 4\gamma + \beta(6\gamma - 4) = 0, \\ & \frac{2\gamma(3\gamma - 2)[-2 + (4a_2 + 3b_1 - 4b_2)\gamma^2 - 12(a_2 - b_2)\gamma^3]}{(6\gamma^2 - 8\gamma + 3)} \\ & - \frac{2\gamma(3\gamma - 2)^2(-1 - 12\gamma + 18\gamma^2)\sigma}{3(4\gamma - 3)(6\gamma^2 - 8\gamma + 3)} \\ & + \frac{(3 - 4\gamma)^2[(a_4 - b_4)(6\gamma^2 - 8\gamma + 3) - 3(2 - 3\gamma)^2(1 - 2b_1\gamma + 4(a_2 - b_2)\gamma^2)]}{4(3\gamma - 2)^3(6\gamma^2 - 8\gamma + 3)} \\ & + \frac{(4\gamma - 3)[4 + 3\gamma + (-8a_2 - 12b_1 + 8b_2)\gamma^2 + 36(a_2 - b_2)\gamma^3]}{2(6\gamma^2 - 8\gamma + 3)} = 0, \\ & - \frac{(a_4 - 2b_4)(3 - 4\gamma)^2 + (2 - 3\gamma)^2[-1 + 4(a_2 - 2b_2)\gamma^2]}{(2 - 3\gamma)^2} \\ & + \frac{32\gamma^2(-2 + 3\gamma)^4\sigma^2}{3(-3 + 4\gamma)^2(6\gamma^2 - 8\gamma + 3)} \\ & + \frac{4\gamma\sigma(2 - 3\gamma)[1 - 6\gamma(b_1 - 2a_2\gamma + 2b_2\gamma)(6\gamma^2 - 8\gamma + 3)]}{3(3 - 4\gamma)(6\gamma^2 - 8\gamma + 3)} \\ & + \frac{4\gamma\sigma(-a_4 + b_4)(3 - 4\gamma)}{(2 - 3\gamma)} = 0. \end{aligned} \quad (2.11)$$

Solving (2.11) for  $\beta, a_4, b_4$  yields

$$\begin{aligned} \beta &= \frac{3 - 4\gamma}{4 - 6\gamma}, \\ a_4 &= \frac{-6 + \beta[3 + 12\gamma(-b_1 + 3a_2\gamma - 4b_2\gamma)] + 4\gamma^2[3b_1 + 2(a_2 - 6a_2\gamma + 6b_2\gamma)]}{4\beta^2(-2 + 3\beta)} \\ & - \frac{\gamma\sigma[-4 + 12\beta + (12 + 3(-3 + 4b_1)\beta)\gamma - 6(3 + 2b_1)\gamma^2]}{6\beta^3(-2 + 3\beta)(\beta - \gamma)} \\ & - \frac{\gamma^3\sigma[4(a_2 - b_2)(3\gamma - 2) - 3b_1]}{\beta^3(-2 + 3\beta)} \\ & + \frac{\gamma^2\sigma^2(-2 + 3\gamma)(-4 + 6\beta + 3\gamma)}{3\beta^4(-2 + 3\beta)(\beta - \gamma)}, \\ b_4 &= \frac{-1 + 3b_1\gamma(-\beta + \gamma) + 2\gamma^2[6a_2(\beta - \gamma) + b_2(2 - 9\beta + 6\gamma)]}{2\beta^2(-2 + 3\beta)} \end{aligned}$$

$$\begin{aligned}
& + \frac{\gamma^2 \sigma^2 (-2 + 3\gamma)(-4 + 6\beta + 3\gamma)}{3\beta^4(-2 + 3\beta)(\beta - \gamma)} \\
& + \frac{\gamma \sigma [-2\beta(1 + 2b_1\gamma) + (3 + 4b_1)\gamma^2]}{2\beta^3(-2 + 3\beta)(\beta - \gamma)} \\
& + \frac{\gamma^3 \sigma [4(a_2 - b_2)(2 - 3\gamma) + 3b_1]}{\beta^3(-2 + \beta)}. \tag{2.12}
\end{aligned}$$

By substituting  $a_1, a_3, b_3, \beta, a_4, b_4$  into  $\Gamma_5 = 0$ , we find five relations independently of  $c_2, c_3, c_4$  and  $c_5$  below:

$$\begin{aligned}
& 3 - 12\gamma + 10\gamma^2 = 0, \quad 3 + \gamma^2(6 - 24\sigma) + 18\gamma^3\sigma + \gamma(-9 + 8\sigma) = 0, \\
& 125(137609 + 57501\sqrt{6}) + 648(28357 + 11923\sqrt{6})a_2 \\
& - 2250(4202 + 1753\sqrt{6})b_1 - 648(28357 + 11923\sqrt{6})b_2 = 0, \\
& 325(229308 + 91387\sqrt{6}) - 72(945441 + 409574\sqrt{6})a_2 \\
& + 90(647706 + 261659\sqrt{6})b_1 + 72(655416 + 304849\sqrt{6})b_2 = 0, \\
& -25(36683358 + 14824337\sqrt{6}) + 72(2673882 + 1215523\sqrt{6})a_2 \\
& - 540(627727 + 253603\sqrt{6})b_1 + 288(672867 + 227813\sqrt{6})b_2 = 0. \tag{2.13}
\end{aligned}$$

Solving (2.13) for  $\gamma, \sigma, b_1, a_2, b_2$  and simplifying, we obtain:

$$\begin{aligned}
& \sigma = -\frac{(3 + 8\sqrt{6})}{25}, \quad \gamma = \frac{6 - \sqrt{6}}{10}, \\
& a_2 = -\frac{5(38 + 13\sqrt{6})}{48}, \quad b_1 = -\frac{(139 + 64\sqrt{6})}{36}, \quad b_2 = -\frac{5(8 + 3\sqrt{6})}{36}. \tag{2.14}
\end{aligned}$$

After substituting (2.14) into (2.9), (2.10), (2.12), we further simplify:

$$\begin{aligned}
& a_1 = -\frac{(41 + 21\sqrt{6})}{12}, \quad a_3 = \frac{31 + 9\sqrt{6}}{12}, \quad b_3 = \frac{7(11 + 4\sqrt{6})}{36}, \\
& \beta = \frac{6 + \sqrt{6}}{10}, \quad a_4 = \frac{118 - 23\sqrt{6}}{48}, \quad b_4 = \frac{54 - 19\sqrt{6}}{36}. \tag{2.15}
\end{aligned}$$

We finally substitute overall relations found so far into  $\Gamma_6$  to obtain  $\Gamma_6 = (\phi_1 c_2^5 + \phi_2 c_2^3 c_3 + \phi_3 c_2^2 c_4 + \phi_4 c_3 c_4 + \phi_5 c_2 c_3^2 + \phi_6 c_2 c_5 + \phi_7 c_6)$ , where  $\phi_i$  are described in (2.1). This completes the proof.  $\square$

REMARK 2.2. From the first two equations of (2.13), the other pair of values for  $\sigma, \gamma$  is possible with  $\sigma = -\frac{(3-8\sqrt{6})}{25}, \gamma = \frac{6+\sqrt{6}}{10}$ . In this case,  $\beta = \frac{6-\sqrt{6}}{10}$  and all coefficients  $a_i, b_j$  can be obtained with  $\sqrt{6}$  replaced by  $-\sqrt{6}$ .

### 3. Numerical experiments and concluding remarks

Many numerical analysis arising in engineering problems often requires a large number of working-precision digits. Computing asymptotic error constants  $\eta = \lim_{n \rightarrow \infty} \frac{|e_n|}{|e_{n-1}|^p}$  with several significant digits of accuracy would encounter extreme calculations due to the indeterminate form of a small-number division near the root  $\alpha$ .

In the current numerical experiments, high-precision computing with programming language *Mathematica* (Version 7) has been performed with 100 working-precision digits, being large enough to minimize round-off errors as well as to clearly observe the computed asymptotic error constants requiring small-number divisions. In addition, the error bound  $\epsilon = \frac{1}{2} \times 10^{-80}$  was used. The values of initial guess  $x_0$  were selected close to  $\alpha$  to guarantee the convergence of the iterative methods. Only 15 significant digits of approximated roots  $x_n$  are displayed in Tables 1-3 due to the limited paper space, although 80 significant digits are available. Numerical experiments have been carried out on a personal computer equipped with an AMD 3.1 Ghz dual-core processor and 64-bit Windows 7 operating system.

Iterative method (1.5) was identified by **Y1** and has shown acceptable performance when applied to a test function:

$$F_1(x) = \sin(x + 1) - x + 2, \alpha \approx 2.07076672714204.$$

**DEFINITION 3.1.** (Asymptotic Convergence Order) Assume that the asymptotic error constant  $\eta = \lim_{n \rightarrow \infty} \frac{|e_n|}{|e_{n-1}|^p}$  is known. Then we can define the *asymptotic convergence order*  $p_a = \lim_{n \rightarrow \infty} \frac{\log |e_n/\eta|}{\log |e_{n-1}|}$ , being abbreviated by ACO.

Method **Y1** in Table 1 clearly confirmed hexic-order convergence. Table 1 lists iteration indexes  $n$ , approximate zeros  $x_n$ , residual errors  $|f(x_n)|$ , errors  $|e_n| = |x_n - \alpha|$  and computational asymptotic error constants  $\eta_n = \left| \frac{e_n}{e_{n-1}^6} \right|$  as well as the theoretical asymptotic error constant  $\eta$  and computational asymptotic convergence order  $p_n = \frac{\log |e_n/\eta|}{\log |e_{n-1}|}$ . The values of  $\eta_n$  agree up to 7 significant digits with  $\eta$ . As expected, the computed asymptotic order of convergence well approaches 6.

Additional functions below are tested to verify the convergence behavior of proposed scheme (1.5):

$n$	$x_n$	$ F(x_n) $	$ e_n $	$\frac{e_n}{e_{n-1}^6}$	$\eta$	$p_n$
0	1.9	0.339249	0.170767			
1	2.07076671448853	$2.527 \times 10^{-8}$	$1.265 \times 10^{-8}$	0.0005102599209	0.0002274623374	5.54288
2	2.07076672714204	$1.864 \times 10^{-51}$	$9.336 \times 10^{-52}$	0.0002274623568		6.00000
3	2.07076672714204	$0.0 \times 10^{-99}$	$0.0 \times 10^{-99}$			

$$p_n = \frac{\log |e_n/\eta|}{\log |e_{n-1}|}, * (0.95) = 0.95 + 1.5i$$

TABLE 1. Convergence for sample test functions  $F_1(x)$  with method **Y1**

$f$	$x_0$	$ x_n - \alpha $	<b>JA</b>	<b>CH</b>	<b>PG</b>	<b>Y1</b>
$f_1$	-0.31	$ x_1 - \alpha $	2.39e-8*	9.17e-10	3.67e-9	1.06e-9
		$ x_2 - \alpha $	3.95e-38	3.69e-54	6.54e-50	6.66e-54
		$ x_3 - \alpha $	0.0e-100	0.0e-100	0.0e-100	0.0e-100
$f_2$	1.2	$ x_1 - \alpha $	4.14e-5	3.76e-6	1.85e-5	3.02e-6
		$ x_2 - \alpha $	4.92e-21	3.78e-29	9.59e-27	6.06e-32
		$ x_3 - \alpha $	0.0e-99	0.0e-99	0.0e-99	0.0e-99
$f_3$	0.45 +0.85i	$ x_1 - \alpha $	1.05e-7	8.84e-9	1.28e-8	1.90e-8
		$ x_2 - \alpha $	1.99e-36	9.56e-50	1.36e-48	1.75e-47
		$ x_3 - \alpha $	0.0e-100	0.0e-99	0.0e-99	0.0e-100
$f_4$	0.1	$ x_1 - \alpha $	8.32e-7	5.00e-7	2.41e-6	3.87e-8
		$ x_2 - \alpha $	5.39e-32	1.01e-38	5.08e-34	1.91e-45
		$ x_3 - \alpha $	0.0e-130	0.0e-137	0.0e-199	0.0e-144
$f_5$	0.84	$ x_1 - \alpha $	2.55e-7	2.22e-9	8.79e-9	5.24e-9
		$ x_2 - \alpha $	2.87e-33	98.93e-53	1.04e-48	4.14e-50
		$ x_3 - \alpha $	0.0e-100	0.0e-100	0.0e-100	0.0e-100
$f_6$	1.1	$ x_1 - \alpha $	2.77e-6	1.78e-7	2.91e-7	3.91e-7
		$ x_2 - \alpha $	1.12e-27	1.09e-40	7.98e-40	9.86e-39
		$ x_3 - \alpha $	0.0e-99	0.0e-99	0.0e-99	0.0e-99
$f_7$	0.3	$ x_1 - \alpha $	5.33e-10	1.08e-11	7.57e-12	4.84e-12
		$ x_2 - \alpha $	3.31e-48	8.90e-68	6.59e-69	2.21e-70
		$ x_3 - \alpha $	0.0e-100	0.0e-100	0.0e-100	0.0e-100

\* 2.39e-8 denotes  $2.39 \times 10^{-8}$

TABLE 2. Comparison of  $|x_n - \alpha|$  for  $f_1(x) - f_7(x)$  among listed methods

$$f_1(x) = x \cos \frac{3\pi}{2}x - \log(x^2 - \frac{1}{x} - \frac{19}{9}), \alpha = -1/3, x_0 = -0.31,$$

$$f_2(x) = \sqrt{2}x \cos x^2 - \log(e + 8x^2 - 4\pi) + 1, \alpha = \sqrt{\frac{\pi}{2}}, x_0 = 1.2,$$

$$f_3(x) = \cos(x^2 - x + \frac{37}{36}) + 3x - \frac{5}{2} - i\sqrt{7}, \alpha = \frac{1}{2} + i\frac{\sqrt{7}}{3}, x_0 = 0.45 + 0.85i, i = \sqrt{-1},$$

$$f_4(x) = x^3 - 2 + (x + 2) \log(e + x^2), \alpha = 0, x_0 = 0.1,$$

$$f_5(x) = x^5 + x^3 + e^{2x} - 7, \alpha \approx 0.878720933693359, x_0 = 0.84,$$

$$f_6(x) = 4 \cos^2 x + \log(e^2 + 9x^2 - \pi^2) - 3, \alpha = \pi/3, x_0 = 1.1,$$

$$f_7(x) = 3x^2 + xe^{1-x^2} + \sin(x^3 + 2) - 2, \alpha \approx 0.323329877529435, x_0 = 0.3,$$

with  $\log z$  ( $z \in \mathbb{C}$ ) representing a principal analytic branch such that  $-\pi \leq \text{Im}(\log z) < \pi$ .



$f$	$x_0$	$p_n$	<b>JA</b>	<b>CH</b>	<b>PG</b>	<b>Y1</b>
$f_1$	-0.31	$p_1$	5.10138	6.02347	6.04048	5.90624
		$p_2$	5.00000	6.00000	6.00000	6.00000
$f_2$	1.2	$p_1$	4.70107	5.28779	5.57932	5.82983
		$p_2$	4.99994	5.99998	6.00003	6.00000
$f_3$	0.45 $+0.85i$	$p_1$	5.02463	5.99413	6.01727	5.93624
		$p_2$	5.00000	6.00000	6.00000	6.00000
$f_4$	0.1	$p_1$	5.20917	6.11028	6.03205	7.16720
		$p_2$	5.00000	6.00000	6.00000	6.00000
$f_5$	0.84	$p_1$	4.96969	6.03203	5.95361	6.07670
		$p_2$	5.00000	6.00000	6.00000	6.00000
$f_6$	1.1	$p_1$	5.00471	5.69594	5.21100	5.35830
		$p_2$	5.00000	6.00000	6.00000	6.00000
$f_7$	0.3	$p_1$	4.99851	5.93970	5.92075	5.85203
		$p_2$	5.00000	6.00000	6.00000	6.00000

TABLE 3. Comparison of computational ACO  $p_n = \frac{\log |e_n/\eta|}{\log |e_{n-1}|}$  for  $f_1(x) - f_7(x)$

For the purpose of comparison, we first identify methods (1.1), (1.3), (1.4) by **JA**, **CH**, **PG**, respectively. Table 2 displays the values of  $|x_n - \alpha|$  for methods **JA**, **CH**, **PG**, **Y1**. As a result of Table 2, proposed method shows favorable or equivalent performance as compared with existing methods **JA**, **CH** and **PG**. In Table 2, italicized numbers indicates the least errors  $|x_n - \alpha|$ . When the same order of convergence is known, one should note that the local convergence of  $|x_n - \alpha|$  behaves differently depending on  $c_j$ , namely  $f(x)$  and  $\alpha$ . Table 3 well exhibits computational convergence orders of the listed methods, among which method **Y1** clearly shows the convergence order of 6.

During the current numerical experiments, **CH** has shown best accuracy for  $f_3, f_5$ , while **Y1** for  $f_2, f_4, f_7$ . On the other hand, two methods **CH**, **Y1** have shown similar performance for  $f_1$  and three methods **CH**, **PG**, **Y1** for  $f_6$ . Computational accuracy is sensitively dependent on the structures of the iterative methods, the sought zeros and the test functions as well as good initial approximations. We should be aware that no iterative method always shows best accuracy for all the test functions. The corresponding efficiency index for the proposed method (1.5) is found to be  $6^{1/4}$ , which is same as those of listed sextic-order methods but better than that of fourth-order Jarratt's method [2]. A new development of a family of higher-order iterative methods will be pursued in the near future using the current approach employing three derivatives and one function.

### References

- [1] L. V. Ahlfors, *Complex Analysis*, McGraw-Hill Book, Inc, 1979.
- [2] I. K. Argyros, D. Chen, and Q. Qian, *The Jarratt method in Banach space setting*, J. Comput. Appl. Math. **51** (1994), 103-106.
- [3] W. Bi, Q. Wu, and H. Ren, *A new family of eighth-order iterative methods for solving nonlinear equations*, Appl. Math. Comput. **214** (2009), no. 4, 236-245.
- [4] C. Chun, *Certain improvements of Chebyshev-Halley methods with accelerated fourth-order convergence*, Appl. Math. Comput. **189** (2007), 597-601.
- [5] C. Chun, *Some improvements of Jarratts method with sixth-order convergence*, Appl. Math. Comput. **190** (2007), 1432-1437.
- [6] L. Fanga, T. Chen, L. Tian, L. Sun, and B. Chen, *A modified Newton-type method with sixth-order convergence for solving nonlinear equations*, Procedia Engineering **15** (2011), 3124-3128.
- [7] Y. H. Geum and Y. I. Kim, *A biparametric family of four-step sixteenth-order root-finding methods with the optimal efficiency index*, Appl. Math. Lett. **24** (2011), 1336-1342.
- [8] P. Jarratt, *Multipoint iterative methods for solving certain equations*, The Computer Journal **8** (1966), no. 4, 398-400.
- [9] Y. I. Kim, *A new two-step biparametric family of sixth-order iterative methods free from second derivatives for solving nonlinear algebraic equations*, Appl. Math. Comput. **215** (2010), 3418-3424.
- [10] S. K. Parhi and D. K. Gupta, *A sixth order method for nonlinear equations*, Appl. Math. Comput. **203** (2008), 50-55.
- [11] Y. Peng, H. Feng, Q. Li, and X. Zhang, *A fourth-order derivative-free algorithm for nonlinear equations*, J. Comput. Appl. Math. **235** (2011), 2551-2559.
- [12] B. V. Shabat, *Introduction to Complex Analysis PART II, Functions of Several Variables*, American Mathematical Society (1992).
- [13] F. Soleymani, *Regarding the accuracy of optimal eighth-order methods*, Math. Comput. Model. **53** (2011), 5-6, 1351-1357.
- [14] J. F. Traub, *Iterative Methods for the Solution of Equations*, Chelsea Publishing Company (1982).
- [15] X. Wang, J. Kou, and Y. Li, *A variant of Jarratt method with sixth-order convergence*, Appl. Math. Comput. **204** (2008), 14-19.
- [16] S. Wolfram, *The Mathematica Book*(5th ed.), Wolfram Media, 2003.

\*

Department of Applied Mathematics  
Dankook University  
Cheonan 330-714, Republic of Korea  
*E-mail*: yikbell@dankook.ac.kr