

## UNIQUE POINT OF COINCIDENCE FOR TWO MAPPINGS WITH $\varphi$ - OR $\psi\text{-}\phi$ -CONTRACTIVE CONDITIONS ON 2-METRIC SPACES

MING-XING XU\*, XIN HUANG\*\*, AND YONG-JIE PIAO\*\*\*

**ABSTRACT.** We discuss and obtain some existence theorems of unique point of coincidence for two mappings satisfying  $\varphi$ -contractive conditions or  $\psi\text{-}\phi$ -contractive conditions determined by semi-continuous functions on non-complete 2-metric spaces, in which the mappings do not satisfy commutativity and uniform boundedness. The obtained results generalize and improve many well-known and corresponding conclusions.

### 1. Introduction and preliminaries

There have appeared many common fixed point theorems of mappings with some contractive conditions on 2-metric spaces. But most of them held under subsidiary conditions ([9, 11]), for example; commutativity of mappings or uniform boundness of mappings at some point, and so on. The authors in ([1, 2, 3, 4, 5, 6, 7, 8, 10]) obtained generalized results of coincidence points and common fixed points for infinite or finite family of mappings satisfying generalized linear or non-linear contractive or quasi-contractive conditions and expansive conditions under removing the above subsidiary conditions. These obtained results greatly generalize and improve the corresponding conclusions.

In this paper, we will introduce three real functions with some kind of properties to establish contractive conditions of two self-mappings on 2-metric spaces, and construct convergent sequences to discuss the existence problems of unique points of coincidence of the given mappings.

---

Received January 11, 2016; Accepted July 15, 2016.

2010 Mathematics Subject Classification: Primary 47H05, 47H10, 54E40.

Key words and phrases: point of coincidence,  $\varphi$ -contractive condition,  $\psi\text{-}\phi$ -contractive condition, altering distance function, weakly compatible.

Correspondence should be addressed to Yong-Jie Piao, [sxpyj@ybu.edu.cn](mailto:sxpyj@ybu.edu.cn).

This work was Supported by the National Natural Science Foundation of China (No. 11361064) and YDBKSKY of China(No. 2016088).

**DEFINITION 1.1.** ([5, 6, 9]) A 2-metric space  $(X, d)$  consists of a nonempty set  $X$  and a function  $d : X \times X \times X \rightarrow [0, +\infty)$  such that

- (i) for distant elements  $x, y \in X$ , there exists an  $u \in X$  such that  $d(x, y, u) \neq 0$ ;
- (ii)  $d(x, y, z) = 0 \iff$  at least two elements in  $\{x, y, z\}$  are equal;
- (iii)  $d(x, y, z) = d(u, v, w)$ , where  $\{u, v, w\}$  is any permutation of  $\{x, y, z\}$ ;
- (iv)  $d(x, y, z) \leq d(x, y, u) + d(x, u, z) + d(u, y, z)$  for all  $x, y, z, u \in X$ .

**DEFINITION 1.2.** ([5, 6, 9]) A sequence  $\{x_n\}_{n \in \mathbb{N}}$  in 2-metric space  $(X, d)$  is said to be a Cauchy sequence, if for each  $\varepsilon > 0$  there exists  $N \in \mathbb{N}$  such that  $d(x_n, x_m, a) < \varepsilon$  for all  $a \in X$  and  $n, m > N$ .  $\{x_n\}_{n \in \mathbb{N}}$  is said to be convergent to  $x \in X$ , if for each  $a \in X$ ,  $\lim_{n \rightarrow +\infty} d(x_n, x, a) = 0$ . And we write that  $x_n \rightarrow x$  and call  $x$  the limit of  $\{x_n\}_{n \in \mathbb{N}}$ .

**DEFINITION 1.3.** ([5, 6, 9]) A 2-metric space  $(X, d)$  is said to be complete, if every cauchy sequence in  $X$  is convergent.

**LEMMA 1.4.** ([12]) Let  $\{x_n\}$  be a sequence in 2-metric space  $(X, d)$  such that  $\lim_{n \rightarrow \infty} d(x_n, x_{n+1}, a) = 0$  for all  $a \in X$ . If  $\{x_n\}$  is not a Cauchy sequence, then there exist  $a \in X$  and  $\epsilon > 0$  such that for each  $i \in \mathbb{N}$  there exist  $m(i), n(i) \in \mathbb{N}$  with  $m(i), n(i) > i$  such that

- (i)  $m(i) > n(i)$  and  $n(i) \rightarrow \infty$  as  $i \rightarrow \infty$ ;
- (ii)  $d(x_{m(i)}, x_{n(i)}, a) > \epsilon$ , but  $d(x_{m(i)-1}, x_{n(i)}, a) \leq \epsilon$ .

**LEMMA 1.5.** ([5, 6, 7]) If a sequence  $\{x_n\}$  in a 2-metric space  $(X, d)$  converges to  $x \in X$ , then

$$\lim_{n \rightarrow \infty} d(x_n, b, c) = d(x, b, c), \forall b, c \in X.$$

**DEFINITION 1.6.** ([5, 6, 7]) Let  $f, g : X \rightarrow X$  be two mappings. If  $w = fx = gx$  for some  $w, x \in X$ , then  $x$  is called a coincidence point of  $f$  and  $g$ ,  $w$  is called a point of coincidence of  $f$  and  $g$ .

**DEFINITION 1.7.** ([5, 6, 7]) Two mappings  $f, g : X \rightarrow X$  are called be weakly compatible if  $fgx = gfx$  whenever  $fx = gx$  for  $x \in X$

**LEMMA 1.8.** ([5, 6, 7]) If  $f, g : X \rightarrow X$  are weakly compatible and have an unique point of coincidence  $w$ , then  $w$  is the unique common fixed point of  $f$  and  $g$ .

## 2. Unique point of coincidence and common fixed point

Let  $\varphi : [0, \infty) \rightarrow [0, \infty)$  be a non-decreasing function satisfying the following conditions:

$(\varphi_1)$ :  $\varphi(0) = 0$ ;  $(\varphi_2)$ :  $0 < \varphi(t) < t$  for all  $t > 0$ ;  $(\varphi_3)$ :  $\sum_{n=1}^{\infty} \varphi^n(t) < \infty$  for each  $t \in (0, \infty)$ .

**THEOREM 2.1.** Let  $(X, d)$  be a 2-metric space,  $f, g : X \rightarrow X$  two mappings such that  $fX \subset gX$ . Suppose that

$$(2.1) \quad d(fx, fy, a) \leq \varphi(M(x, y, a)), \quad \forall x, y, a \in X,$$

where  $M(x, y, a) = \max\{d(gx, gy, a), d(gx, fx, a), d(gy, fy, a), \frac{1}{2}[d(gx, fy, a) + d(gy, fx, a)]\}$  and  $\varphi$  is upper semi-continuous. If  $fX$  or  $gX$  is complete, and  $X$  is bounded (i.e.,  $\sup_{x,y,z \in X} d(x, y, z) < +\infty$ ), then  $f$  and  $g$  have a unique point of coincidence. Furthermore, if  $f$  and  $g$  are weakly compatible, then  $f$  and  $g$  have a unique common fixed point.

*Proof.* Take  $x_0 \in X$  and construct sequences  $\{x_n\}$  and  $\{y_n\}$  satisfying

$$y_n = fx_n = gx_{n+1}, \quad n = 0, 1, 2, \dots.$$

For any fixed  $n \geq 1$ , in view of (2.1),

$$\begin{aligned} & d(y_n, y_{n+1}, y_{n+2}) \\ &= d(fx_{n+2}, fx_{n+1}, y_n) \\ &\leq \varphi(M(x_{n+2}, x_{n+1}, y_n)), \\ &= \varphi\left(\max\left\{d(gx_{n+2}, gx_{n+1}, y_n), d(gx_{n+2}, fx_{n+2}, y_n), d(gx_{n+1}, fx_{n+1}, y_n), \right.\right. \\ &\quad \left.\left.\frac{d(gx_{n+2}, fx_{n+1}, y_n) + d(gx_{n+1}, fx_{n+2}, y_n)}{2}\right\}\right) \\ &= \varphi(d(y_{n+2}, y_{n+1}, y_n)). \end{aligned}$$

Hence by  $(\varphi_2)$ ,

$$d(y_n, y_{n+1}, y_{n+2}) = 0, \quad \forall n = 1, 2, \dots.$$

Suppose that  $d(y_k, y_n, y_{n+1}) = 0$  for  $n - k \geq 1$ , then by (2.1),

$$d(y_{n+1}, y_{n+2}, y_k) = d(fx_{n+1}, fx_{n+2}, y_k) \leq \varphi(M(x_{n+1}, x_{n+2}, y_k)),$$

where

$$\begin{aligned} & M(x_{n+1}, x_{n+2}, y_k) \\ &= \max\left\{d(gx_{n+1}, gx_{n+2}, y_k), d(gx_{n+1}, fx_{n+1}, y_k), d(gx_{n+2}, fx_{n+2}, y_k), \right. \\ &\quad \left.\frac{d(gx_{n+1}, fx_{n+2}, y_k) + d(gx_{n+2}, fx_{n+1}, y_k)}{2}\right\} \\ &= \max\left\{d(y_{n+1}, y_{n+2}, y_k), \frac{d(y_n, y_{n+2}, y_k)}{2}\right\} \end{aligned}$$

But  $d(y_n, y_{n+2}, y_k) \leq d(y_n, y_{n+1}, y_k) + d(y_{n+1}, y_{n+2}, y_k) + d(y_n, y_{n+1}, y_{n+2}) = d(y_{n+1}, y_{n+2}, y_k)$ , hence

$$M(x_{n+1}, x_{n+2}, y_k) = d(y_{n+1}, y_{n+2}, y_k),$$

therefore

$$d(y_k, y_{n+1}, y_{n+2}) \leq \varphi(d(y_{n+1}, y_{n+2}, y_k)),$$

which implies that  $d(y_k, y_{n+1}, y_{n+2}) = 0$  by  $(\varphi_2)$ . Hence

$$(2.2) \quad d(y_k, y_n, y_{n+1}) = 0, \quad \forall k, n \in \mathbb{N}, k \leq n.$$

For every  $m, n, l \in \mathbb{N}$  with  $m \leq n \leq l$ , using (2.2), we obtain

$$\begin{aligned} & d(y_m, y_n, y_l) \\ & \leq d(y_m, y_n, y_{l-1}) + d(y_m, y_{l-1}, y_l) + d(y_n, y_{l-1}, y_l) = d(y_m, y_n, y_{l-1}) \\ & \leq \dots \\ & \leq d(y_m, y_n, y_{n+1}) \\ & = 0. \end{aligned}$$

Hence we have

$$(2.3) \quad d(y_m, y_n, y_l) = 0, \quad \forall m, n, l \in \mathbb{N}.$$

For any fixed  $n \geq 1$  and  $a \in X$ ,

$$d(y_n, y_{n+1}, a) = d(fx_n, fx_{n+1}, a) \leq \varphi(M(x_n, x_{n+1}, a)),$$

where

$$\begin{aligned} & M(x_n, x_{n+1}, a) \\ & = \max \left\{ d(gx_n, gx_{n+1}, a), d(gx_n, fx_n, a), d(gx_{n+1}, fx_{n+1}, a), \right. \\ & \quad \left. \frac{d(gx_n, fx_{n+1}, a) + d(gx_{n+1}, fx_n, a)}{2} \right\} \\ & = \max \left\{ d(y_{n-1}, y_n, a), d(y_n, y_{n+1}, a), \frac{d(y_{n-1}, y_{n+1}, a)}{2} \right\}. \end{aligned}$$

But using Definition 1.1(iv) and (2.3), we obtain

$$d(y_{n-1}, y_{n+1}, a) \leq d(y_{n-1}, y_n, a) + d(y_n, y_{n+1}, a),$$

hence

$$M(x_n, x_{n+1}, a) = \max \{ d(y_{n-1}, y_n, a), d(y_n, y_{n+1}, a) \}.$$

If there exists  $a \in X$  such that  $d(y_{n-1}, y_n, a) < d(y_n, y_{n+1}, a)$ , then  $M(x_n, x_{n+1}, a) = d(y_n, y_{n+1}, a) > 0$ , hence by  $(\varphi_2)$ , we have

$$d(y_n, y_{n+1}, a) \leq \varphi(M(x_n, x_{n+1}, a)) = \varphi(d(y_n, y_{n+1}, a)) < d(y_n, y_{n+1}, a),$$

which is a contradiction. So  $M(x_n, x_{n+1}, a) = d(y_{n-1}, y_n, a)$  for all  $n \in \mathbb{N}$  and  $a \in X$ , therefore, for all  $n \in \mathbb{N}$  and  $a \in X$ ,

$$d(y_n, y_{n+1}, a) \leq \varphi(M(x_n, x_{n+1}, a)) = \varphi(d(y_{n-1}, y_n, a)).$$

Continuing this process, we obtain that for all  $n \in \mathbb{N}$  and  $a \in X$ ,

$$d(y_n, y_{n+1}, a) \leq \varphi^n(d(y_0, y_1, a)).$$

Since  $X$  is bounded, there exists  $M > 0$  such that  $d(y_0, y_1, a) < M$  for all  $a \in X$ . So we have

$$d(y_n, y_{n+1}, a) \leq \varphi^n(M), \quad \forall n \in \mathbb{N}, a \in X.$$

For any  $m, n \in \mathbb{N}$  with  $n > m$  and  $a \in X$ ,

$$\begin{aligned} & d(y_m, y_n, a) \\ & \leq d(y_m, y_{m+1}, a) + d(y_{m+1}, y_n, a) + d(y_m, y_n, y_{m+1}) \\ & = d(y_m, y_{m+1}, a) + d(y_{m+1}, y_n, a) \\ & \leq d(y_m, y_{m+1}, a) + d(y_{m+1}, y_{m+2}, a) + d(y_{m+2}, y_n, a) + d(y_{m+1}, y_n, y_{m+2}) \\ & = d(y_m, y_{m+1}, a) + d(y_{m+1}, y_{m+2}, a) + d(y_{m+2}, y_n, a) \\ & \leq \dots \\ & \leq d(y_m, y_{m+1}, a) + d(y_{m+1}, y_{m+2}, a) + \dots + d(y_{n-1}, y_n, a) \\ & \leq \sum_{k=m}^{n-1} \varphi^k(M). \end{aligned}$$

Hence by  $(\varphi_3)$ , we know that  $\{y_n\}$  is a Cauchy sequence in  $X$ .

Suppose that  $gX$  is complete, then there exist  $u, v \in X$  such that  $y_n = fx_n = gx_{n+1} \rightarrow u = gv$  as  $n \rightarrow \infty$ .

For any  $n$  and  $a \in X$ ,

$$\begin{aligned} & d(y_n, fv, a) = d(fx_n, fv, a) \\ & \leq \varphi(M(x_n, v, a)) \\ & = \varphi(\max\{d(gx_n, gv, a), d(gx_n, fx_n, a), d(gv, fv, a), \frac{d(gx_n, fv, a) + d(gv, fx_n, a)}{2}\}) \\ & = \varphi(\max\{d(y_{n-1}, gv, a), d(y_{n-1}, y_n, a), d(gv, fv, a), \frac{d(y_{n-1}, fv, a) + d(gv, y_n, a)}{2}\}). \end{aligned}$$

Letting  $n \rightarrow \infty$ , then we obtain from the above and Lemma 1.5 that

$$\begin{aligned}
d(gv, fv, a) &= \lim_{n \rightarrow \infty} d(y_n, fv, a) \\
&\leq \limsup_{n \rightarrow \infty} \varphi(\max\{d(y_{n-1}, gv, a), d(y_{n-1}, y_n, a), d(gv, fv, a), \\
&\quad \frac{d(y_{n-1}, fv, a) + d(gv, y_n, a)}{2}\}) \\
&\leq \varphi(\lim_{n \rightarrow \infty} \max\{d(y_{n-1}, gv, a), d(y_{n-1}, y_n, a), d(gv, fv, a), \\
&\quad \frac{d(y_{n-1}, fv, a) + d(gv, y_n, a)}{2}\}) \\
&= \varphi(d(gv, fv, a)).
\end{aligned}$$

This implies that

$$d(gv, fv, a) = 0, \forall a \in X,$$

hence

$$fv = gv = u.$$

So  $u$  is a point of coincidence of  $f$  and  $g$ .

Suppose that  $u_1$  is another point of coincidence of  $f$  and  $g$ , then there exists  $v_1$  satisfying  $u_1 = fv_1 = gv_1$ , and there exists  $a \in X$  satisfying  $d(u, u_1, a) > 0$ . By (2.1) and  $(\varphi_2)$ , we obtain the following contradiction

$$\begin{aligned}
d(u, u_1, a) &= d(fv, fv_1, a) \\
&\leq \varphi(M(v, v_1, a)) = \varphi(\max\{d(gv, gv_1, a), d(gv, fv, a), d(gv_1, fv_1, a), \\
&\quad \frac{d(gv, fv_1, a) + d(gv_1, fv, a)}{2}\}) \\
&= \varphi(d(u, u_1, a)) < d(u, u_1, a).
\end{aligned}$$

Hence  $u$  is the unique point of coincidence of  $f$  and  $g$ .

Suppose that  $fX$  is complete. Then there exist  $u, v, w \in X$  such that  $y_n = fx_n \rightarrow u = fw = gv$  since  $fX \subset gX$ , hence the corresponding conclusion follows from the similar discussion, and the rest proof follows from Lemma 1.8.  $\square$

A mapping  $\psi : [0, \infty) \rightarrow [0, \infty)$  is called an altering distance function if  $\psi$  is continuous and non-decreasing and  $\psi(t) = 0 \Leftrightarrow t = 0$ .

**THEOREM 2.2.** *Let  $(X, d)$  be a 2-metric space,  $f, g : X \rightarrow X$  two mappings such that  $fX \subset gX$ . Suppose that*

$$(2.4) \quad \psi(d(fx, fy, a)) \leq \psi(M(x, y, a)) - \phi(M(x, y, a)), \forall x, y, a \in X,$$

where,  $M(x, y, a)$  is that in Theorem 2.1,  $\psi$  is an altering distance function,  $\phi : [0, \infty) \rightarrow [0, \infty)$  is a lower semi-continuous function such that

$\phi(t) = 0 \Leftrightarrow t = 0$ . If  $fX$  or  $gX$  is complete, then  $f$  and  $g$  have a unique point of coincidence. Furthermore, if  $f$  and  $g$  are weakly compatible, then  $f$  and  $g$  have a unique common fixed point.

*Proof.* Take  $x_0 \in X$  and construct sequences  $\{x_n\}$  and  $\{y_n\}$  satisfying

$$y_n = fx_n = gx_{n+1}, \quad \forall n = 0, 1, 2, 3, \dots$$

For any  $n \in \mathbb{N}$ ,

$\psi(d(fx_{n+1}, fx_{n+2}, y_n)) \leq \psi(M(x_{n+1}, x_{n+2}, y_n)) - \phi(M(x_{n+1}, x_{n+2}, y_n))$ , where  $M(x_{n+1}, x_{n+2}, y_n) = d(y_{n+1}, y_{n+2}, y_n)$  (see Theorem 2.1). Hence

$$\psi(d(y_{n+1}, y_{n+2}, y_n)) \leq \psi(d(y_{n+1}, y_{n+2}, y_n)) - \phi(d(y_{n+1}, y_{n+2}, y_n)),$$

so

$$\phi(d(y_{n+1}, y_{n+2}, y_n)) = 0, \quad \forall n \in \mathbb{N}.$$

By the property of  $\phi$ ,

$$(2.5) \quad d(y_n, y_{n+1}, y_{n+2}) = 0, \quad \forall n \in \mathbb{N}.$$

Suppose that  $d(y_k, y_n, y_{n+1}) = 0$ , where  $n \geq k + 1$ . Using (2.4), we have

$\psi(d(fx_{n+1}, fx_{n+2}, y_k)) \leq \psi(M(x_{n+1}, x_{n+2}, y_k)) - \phi(M(x_{n+1}, x_{n+2}, y_k))$ , where  $M(x_{n+1}, x_{n+2}, y_k) = \max\{d(y_{n+1}, y_{n+2}, y_k), \frac{d(y_n, y_{n+2}, y_k)}{2}\}$  (see Theorem 2.1). By (2.5) and the assumption,

$$\begin{aligned} d(y_n, y_{n+2}, y_k) &\leq d(y_n, y_{n+1}, y_{n+2}) + d(y_{n+1}, y_{n+2}, y_k) + d(y_n, y_{n+1}, y_k) \\ &= d(y_{n+1}, y_{n+2}, y_k), \end{aligned}$$

so

$$M(x_{n+1}, x_{n+2}, y_k) = d(y_{n+1}, y_{n+2}, y_k).$$

Hence

$$\psi(d(y_k, y_{n+1}, y_{n+2})) \leq \psi(d(y_k, y_{n+1}, y_{n+2})) - \phi(d(y_k, y_{n+1}, y_{n+2})),$$

which implies that

$$(2.6) \quad \phi(d(y_k, y_{n+1}, y_{n+2})) = 0 \implies d(y_k, y_{n+1}, y_{n+2}) = 0.$$

Therefore, in view of (2.5) and (2.6), we have the next fact:

$$(2.7) \quad d(y_k, y_n, y_{n+1}) = 0, \quad \forall n, k \in \mathbb{N}, n \geq k \geq 1.$$

For all  $m, n, k \in \mathbb{N}$  with  $k \geq n \geq m$ , using (2.7), we have

$$\begin{aligned} d(y_m, y_n, y_k) &\leq d(y_m, y_n, y_{k-1}) + d(y_m, y_{k-1}, y_k) + d(y_n, y_{k-1}, y_k) \\ &= d(y_m, y_n, y_{k-1}). \end{aligned}$$

Continuing this process, we obtain the following fact: for all  $k \geq n \geq m$ ,

$$(2.8) \quad d(y_m, y_n, y_k) \leq d(y_m, y_n, y_{k-1}) \leq \dots \leq d(y_m, y_n, y_{n+1}) = 0.$$

For any fixed  $n \in \mathbb{N}$  and any  $a \in X$ ,

$$\begin{aligned}\psi(d(y_{n+1}, y_{n+2}, a)) &= \psi(d(fx_{n+1}, fx_{n+2}, a)) \\ &\leq \psi(M(x_{n+1}, x_{n+2}, a)) - \phi(M(x_{n+1}, x_{n+2}, a)),\end{aligned}$$

where  $M(x_{n+1}, x_{n+2}, a) = \max\{d(y_n, y_{n+1}, a), (y_{n+1}, y_{n+2}, a)\}$  (see Theorem 2.1).

If  $d(y_n, y_{n+1}, a) < (y_{n+1}, y_{n+2}, a)$  for some  $a \in X$ , then  $M(x_{n+1}, x_{n+2}, a) = d(y_{n+1}, y_{n+2}, a) > 0$ . Hence using the property of  $\phi$ , we have

$$\begin{aligned}\psi(d(y_{n+1}, y_{n+2}, a)) &\leq \psi(d(y_{n+1}, y_{n+2}, a)) - \phi(d(y_{n+1}, y_{n+2}, a)) \\ &< \psi(d(y_{n+1}, y_{n+2}, a)),\end{aligned}$$

which is a contradiction. Hence

$$M(x_{n+1}, x_{n+2}, a) = d(y_n, y_{n+1}, a), \quad \forall a, n$$

and we have

$$\begin{aligned}(2.9) \quad \psi(d(y_{n+1}, y_{n+2}, a)) &\leq \psi(d(y_n, y_{n+1}, a)) - \phi(d(y_n, y_{n+1}, a)) \\ &\leq \psi(d(y_n, y_{n+1}, a)), \quad \forall a, n.\end{aligned}$$

By the property of  $\psi$ , we obtain that

$$d(y_{n+1}, y_{n+2}, a) \leq d(y_n, y_{n+1}, a), \quad \forall a, n.$$

So for any fixed  $a \in X$ ,  $\{d(y_n, y_{n-1}, a)\}$  is a non-increasing and non-negative real sequence, hence there exists  $r(a) \geq 0$  such that

$$\lim_{n \rightarrow \infty} d(y_n, y_{n-1}, a) = r(a).$$

Letting  $n \rightarrow \infty$  in the both sides of the first inequality in (2.9), we obtain

$$\begin{aligned}\psi(r(a)) &\leq \psi(r(a)) - \liminf_{n \rightarrow \infty} \phi(d(y_n, y_{n+1}, a)) \\ &\leq \psi(r(a)) - \phi(\lim_{n \rightarrow \infty} d(y_n, y_{n+1}, a)) \\ &= \psi(r(a)) - \phi(r(a)),\end{aligned}$$

hence  $\phi(r(a)) = 0$ , which implies that  $r(a) = 0$ . Therefore, we have

$$(2.10) \quad \lim_{n \rightarrow \infty} d(y_n, y_{n+1}, a) = 0, \quad \forall a \in X.$$

If  $\{y_n\}$  is not Cauchy, then by Lemma 1.4, there exist  $a \in X$  and  $\epsilon > 0$  such that for any  $i \in \mathbb{N}$  there exist  $m(i) > n(i) \in \mathbb{N}$  satisfying

- (i)  $m(i), n(i) > i$ ,  $m(i) > n(i) + 1$  and  $n(i) \rightarrow \infty$  as  $i \rightarrow \infty$ ;
- (ii)  $d(y_{m(i)}, y_{n(i)}, a) > \epsilon$ , but  $d(y_{m(i)-1}, y_{n(i)}, a) \leq \epsilon$ ,  $i = 1, 2, \dots$ .

Using (2.8) and (2.10) and the following result

$$\begin{aligned} & d(y_{m(i)}, y_{n(i)}, a) \\ & \leq d(y_{m(i)}, y_{m(i)-1}, a) + d(y_{m(i)-1}, y_{n(i)}, a) + d(y_{m(i)}, y_{n(i)}, y_{m(i)-1}), \end{aligned}$$

we obtain

$$(2.11) \quad \lim_{i \rightarrow \infty} d(y_{m(i)}, y_{n(i)}, a) = \lim_{i \rightarrow \infty} d(y_{m(i)-1}, y_{n(i)}, a) = \epsilon.$$

The following two inequalities hold

$$\begin{aligned} & |d(y_{m(i)}, y_{n(i)}, a) - d(y_{m(i)}, y_{n(i)-1}, a)| \\ & \leq d(y_{n(i)-1}, y_{n(i)}, a) + d(y_{m(i)}, y_{n(i)}, y_{n(i)-1}) \end{aligned}$$

and

$$\begin{aligned} & |d(y_{m(i)-1}, y_{n(i)-1}, a) - d(y_{m(i)}, y_{n(i)-1}, a)| \\ & \leq d(y_{m(i)-1}, y_{m(i)}, a) + d(y_{m(i)}, y_{m(i)-1}, y_{n(i)-1}), \end{aligned}$$

hence using (2.8), (2.10) and (2.11), we have

$$\begin{aligned} (2.12) \quad & \lim_{n \rightarrow \infty} d(y_{m(i)}, y_{n(i)}, a) = \lim_{n \rightarrow \infty} d(y_{m(i)-1}, y_{n(i)}, a) \\ & = \lim_{i \rightarrow \infty} d(y_{m(i)}, y_{n(i)-1}, a) = \lim_{i \rightarrow \infty} d(y_{m(i)-1}, y_{n(i)-1}, a) = \epsilon. \end{aligned}$$

Since

$$\begin{aligned} (2.13) \quad & \psi(d(y_{m(i)}, y_{n(i)}, a)) = \psi(d(fx_{m(i)}, fx_{n(i)}, a)) \\ & \leq \psi(M(x_{m(i)}, x_{n(i)}, a)) - \phi(M(x_{m(i)}, x_{n(i)}, a)), \end{aligned}$$

where

$$\begin{aligned} & M(x_{m(i)}, x_{n(i)}, a) \\ & = \max \left\{ d(gx_{m(i)}, gx_{n(i)}, a), d(gx_{m(i)}, fx_{m(i)}, a), d(gx_{n(i)}, fx_{n(i)}, a), \right. \\ & \quad \left. \frac{d(gx_{m(i)}, fx_{n(i)}, a) + d(gx_{n(i)}, fx_{m(i)}, a)}{2} \right\} \\ & = \max \left\{ d(y_{m(i)-1}, y_{n(i)-1}, a), d(y_{m(i)-1}, y_{m(i)}, a), d(y_{n(i)-1}, y_{n(i)}, a), \right. \\ & \quad \left. \frac{d(y_{m(i)-1}, y_{n(i)}, a) + d(y_{n(i)-1}, y_{m(i)}, a)}{2} \right\}. \end{aligned}$$

By (2.10) and (2.12), we know

$$(2.14) \quad \lim_{i \rightarrow \infty} M(x_{m(i)}, x_{n(i)}, a) = \epsilon.$$

Letting  $i \rightarrow \infty$  in (2.13) and using (2.12) and (2.14), we obtain

$$\begin{aligned}\psi(\epsilon) &\leq \psi(\epsilon) - \liminf_{n \rightarrow \infty} \phi(M(x_{m(i)}, x_{n(i)}, a)) \\ &\leq \psi(\epsilon) - \phi\left(\lim_{n \rightarrow \infty} M(x_{m(i)}, x_{n(i)}, a)\right) = \psi(\epsilon) - \phi(\epsilon),\end{aligned}$$

which implies that  $\phi(\epsilon) = 0$ , i.e.,  $\epsilon = 0$ . This is a contradiction, hence  $\{y_n\}$  is a Cauchy sequence.

Suppose that  $gX$  is complete. Then there exist  $u, v \in X$  such that  $y_n = fx_n = gx_{n+1} \rightarrow u = gv$ . For any  $n$  and  $a \in X$ , we have that

$$\psi(y_n, fv, a) = d(fx_n, fv, a) \leq \psi(M(x_n, v, a) - \phi(M(x_n, v, a))), \quad (2.15)$$

where

$$\begin{aligned}&M(x_n, v, a) \\ &= \max\left\{d(gx_n, gv, a), d(gx_n, fx_n, a), d(gv, fv, a), \frac{d(gx_n, fv, a) + d(gv, fx_n, a)}{2}\right\} \\ &= \max\left\{d(y_{n-1}, gv, a), d(y_{n-1}, y_n, a), d(gv, fv, a), \frac{d(y_{n-1}, fv, a) + d(gv, y_n, a)}{2}\right\}.\end{aligned}$$

Let  $n \rightarrow \infty$ , then

$$\lim_{n \rightarrow \infty} M(x_n, v, a) = d(gv, gv, a), \quad \forall a \in X.$$

Hence Letting  $n \rightarrow \infty$  in (2.15), we obtain that

$$\begin{aligned}&\psi(fv, gv, a) \\ &\leq \psi(d(fv, gv, a)) - \liminf_{n \rightarrow \infty} \phi(M(x_n, v, a)) \\ &\leq \psi(d(fv, gv, a)) - \phi\left(\lim_{n \rightarrow \infty} M(x_n, v, a)\right) \\ &= \psi(d(fv, gv, a)) - \phi(d(fv, gv, a)), \quad \forall a \in X.\end{aligned}$$

This implies that  $\phi(d(fv, gv, a)) = 0$ ,  $\forall a \in X \implies d(fv, gv, a) = 0$ ,  $\forall a \in X$ , hence

$$fv = gv = u.$$

Suppose that  $u_1$  is another point of coincidence of  $f$  and  $g$ , then there exists  $v_1$  satisfying  $u_1 = fv_1 = gv_1$ , and there exists  $a \in X$  satisfying  $d(u, u_1, a) > 0$ . By (2.4)

$$\psi(d(u, u_1, a)) = \psi(d(fv, fv_1, a)) \leq \psi(M(v, v_1, a)) - \phi(M(v, v_1, a)),$$

where  $M(v, v_1, a) = d(u, u_1, a)$ , hence

$$\psi(d(u, u_1, a)) \leq \psi(d(u, u_1, a)) - \phi(d(u, u_1, a)),$$

which implies  $d(u, u_1, a) = 0$ , a contradiction. So  $u$  is the unique pint of coincidence of  $f$  and  $g$ .

Suppose that  $fX$  is complete. Then there exist  $u, v, w \in X$  such that  $y_n = fx_n \rightarrow u = fw = gv$  since  $fX \subset gX$ . The rest proof follows from the similar discussion and Lemma 1.8.  $\square$

REMARK 2.3. (1)  $\varphi$  in Theorem 2.1 need not be a strictly increasing function.

(2) We find that the condition  $\psi(t) = 0 \iff t = 0$  in Theorem 2.2 is superfluous. Hence we only need that  $\psi$  is a continuous and non-decreasing function.

### References

- [1] N. V. Dung, N. T. Hieu, N. T. Thanh Ly, and V. D. Thinh, *Remarks on the fixed point problem of 2-metric spaces*, Fixed Point Theory and Applications 2013: 167. Doi:10.1186/1687-1812-2013-167.
- [2] N. V. Dung and V. T. L. Hang, *Fixed point theorems for weak C-contractions in partially ordered 2-metric spaces*, Fixed Point Theory and Applications 2013, 161. Doi:10.1186/1687-1812-2013-161.
- [3] B. K. Lahiri, Pratulananda Das and Lakshmi Kanta Dey, *Cantor's theorem in 2-metric spaces and its applications to fixed point theorems*, Taiwanese J. Math. **15** (2011), no. 1, 337-352.
- [4] T. Phaneendra and K. K. Swamy, *A unique common fixed point of a pair of self-maps on a 2-metric space*, Mathematica Aeterna. **3** (2013), no. 4, 271-277.
- [5] Y. J. Piao, *Uniqueness of common fixed points for a family of maps with  $\phi_j$ -quasi-contractive type in 2-metric space*, Acta Mathematica Scientia **32** (2012), no. 6, 1079-1085.(In Chinese)
- [6] Y. J. Piao, *Uniqueness of common fixed point for a family of mappings with  $\phi$ -contractive condition in 2-metric space*, Applied Mathematics **3** (2012), 73-77.
- [7] Y. J. Piao, *New unique common fixed point theorems for a infinite family of mappings with  $\phi$ - $\psi$ - $\varphi$ -contractive conditions on 2-metric spaces*, Advances in Fixed Point Theory **5** (2015), no. 4, 420-432.
- [8] Y. J. Piao, *Fixed point theorems for contractive and expansive mappings of Geraghty type on 2-metric spaces*, Advances in Fixed Point Theory **6** (2016), no. 2, 123-135.
- [9] S. L. Singh, *Some contractive type principles on 2-metric spaces and applications*, Mathematics Seminar Notes(Kobe University) **7** (1979), no. 1, 1-11.
- [10] S. L. Singh, S. N. Sishira, and S. Stofile, *Suzuki contraction theorem on a 2-metric space*, J. Adv. Math. Stud. **5** (2012), no. 1, 71-76.
- [11] H. S. Yang and D. S. Xiong, *A common fixed point theorem on p-metric spaces*, Journal of Yunnan Normal University(Science Edition) **21** (2001), no. 1, 9-12.
- [12] D. Zhang and F. Gu, *The common fixed point theorems for a class of  $\Phi$ -contraction conditions mappings in 2-metric spaces*, Journal of Jiangxi Normal University(Natural Science) **35** (2011), no. 6, 595-600.(In Chinese)

\*

Department of Mathematics  
Yanbian University  
Yanji 133002, P. R. China  
*E-mail:* 2799031582@qq.com

\*\*

Department of Mathematics  
Yanbian University  
Yanji 133002, P. R. China  
*E-mail:* 916554437@qq.com

\*\*\*

Department of Mathematics  
Yanbian University  
Yanji 133002, P. R. China  
*E-mail:* sxyj@ybu.edu.cn