

## SYMMETRIC BI- $(f, g)$ -DERIVATIONS IN LATTICES

KYUNG HO KIM\* AND YONG HOON LEE\*\*

ABSTRACT. In this paper, as a generalization of symmetric bi-derivations and symmetric bi- $f$ -derivations of a lattice, we introduce the notion of symmetric bi- $(f, g)$ -derivations of a lattice. Also, we define the isotone symmetric bi- $(f, g)$ -derivation and obtain some interesting results about isotone. Using the notion of  $Fix_a(L)$  and  $KerD$ , we give some characterization of symmetric bi- $(f, g)$ -derivations in a lattice.

### 1. Introduction

Lattices play an important role in many fields such as information theory, information retrieval, information access controls and cryptanalysis ([2], [6], [20]). Recently the properties of lattices were widely researched ([1], [2], [5], [10], [12], [20], [22]). In the theory of rings and near rings, the properties of derivations are an important topic to study ([3], [4], [19]). In [21], G. Szász introduced the notion of derivation on a lattice and discussed some related properties. Y. B. Jun and X. L. Xin [13] applied the notion of derivation in ring, near ring and lattice theory to BCI-algebras. In [24], J. Zhan and Y. L. Liu introduced the notion of left-right (or right-left)  $f$ -derivation of a BCI algebra and investigated some properties.

Recently, the notion of  $f$ -derivation, symmetric bi-derivations and permuting tri-derivations in lattices are introduced and proved some results ([8], [9] and [18]). In this paper, as a generalization of symmetric bi-derivations and symmetric bi- $f$ -derivations of a lattice, we introduce the notion of symmetric bi- $(f, g)$ -derivations of a lattice. Also, we define the isotone symmetric bi- $(f, g)$ -derivation and obtain some interesting

---

Received June 15, 2016; Accepted July 15, 2016.

2010 Mathematics Subject Classification: Primary 06F35, 03G25.

Key words and phrases: (semi)lattice, symmetric bi-derivation, symmetric bi- $(f, g)$ -derivation, isotone,  $Fix_a(L)$ ,  $KerD$ .

Correspondence should be addressed to Yong Hoon Lee, [yonghoon@dankook.ac.kr](mailto:yonghoon@dankook.ac.kr).

results about isotone. Using the notion of  $Fix_a(L)$  and  $KerD$ , we give some characterization of symmetric bi- $(f, g)$ -derivations in a lattice.

## 2. Preliminaries

DEFINITION 2.1. Let  $L$  be a nonempty set endowed with operations  $\wedge$  and  $\vee$ . By a *lattice*  $(L, \wedge, \vee)$ , we mean a set  $L$  satisfying the following conditions:

- (1)  $x \wedge x = x, x \vee x = x,$
- (2)  $x \wedge y = y \wedge x, x \vee y = y \vee x,$
- (3)  $(x \wedge y) \wedge z = x \wedge (y \wedge z), (x \vee y) \vee z = x \vee (y \vee z),$
- (4)  $(x \wedge y) \vee x = x, (x \vee y) \wedge x = x,$  for all  $x, y, z \in L.$

DEFINITION 2.2. Let  $(L, \wedge, \vee)$  be a lattice. A binary relation  $\leq$  is defined by  $x \leq y$  if and only if  $x \wedge y = x$  and  $x \vee y = y.$

LEMMA 2.1. Let  $(L, \wedge, \vee)$  be a lattice. Define the binary relation  $\leq$  as the Definition 2.2. Then  $(L, \leq)$  is a poset and for any  $x, y \in L,$   $x \wedge y$  is the greatest lower bound of  $\{x, y\}$  and  $x \vee y$  is the least upper bound of  $\{x, y\}.$

DEFINITION 2.3. A lattice  $L$  is *distributive* if the identity (1) or (2) holds:

- (1)  $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z),$
- (2)  $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z).$

In any lattice, the conditions (1) and (2) are equivalent.

DEFINITION 2.4. A lattice  $L$  is *modular* if the following identity holds: If  $x \leq z,$  then  $x \vee (y \wedge z) = (x \vee y) \wedge z.$

DEFINITION 2.5. A non-empty subset  $I$  of  $L$  is called an *ideal* if the following conditions hold:

- (1) If  $x \leq y$  and  $y \in I,$  then  $x \in I$  for all  $x, y \in L.$
- (2) If  $x, y \in I$  then  $x \vee y \in I.$

DEFINITION 2.6. Let  $(L, \wedge, \vee)$  be a lattice. Let  $f : L \rightarrow M$  be a function from a lattice  $L$  to a lattice  $M.$

- (1)  $f$  is called a *meet-homomorphism* if  $f(x \wedge y) = f(x) \wedge f(y)$  for all  $x, y \in L.$
- (2)  $f$  is called a *join-homomorphism* if  $f(x \vee y) = f(x) \vee f(y)$  for all  $x, y \in L.$

- (3)  $f$  is called a *lattice-homomorphism* if  $f$  is both a join-homomorphism and a meet-homomorphism.

DEFINITION 2.7. Let  $L$  be a lattice. A mapping  $D(.,.) : L \times L \rightarrow L$  is said to be *symmetric* if  $D(x, y) = D(y, x)$  holds for all  $x, y \in L$ .

DEFINITION 2.8. Let  $L$  be a lattice. A mapping  $d(x) = D(x, x)$  is called a *trace* of  $D(.,.)$ , where  $D(.,.) : L \times L \rightarrow L$  is a symmetric mapping.

DEFINITION 2.9. Let  $L$  be a lattice and let  $D(.,.) : L \times L \rightarrow L$  be a symmetric mapping. We call  $D$  a *symmetric bi-derivation* on  $L$  if it satisfies the following condition

$$D(x \wedge y, z) = (D(x, z) \wedge y) \vee (x \wedge D(y, z))$$

for all  $x, y, z \in L$ .

Obviously, a symmetric bi-derivation  $D$  on  $L$  satisfies the relation

$$D(x, y \wedge z) = (D(x, y) \wedge z) \vee (y \wedge D(x, z))$$

for all  $x, y, z \in L$ .

DEFINITION 2.10. Let  $L$  be a lattice and let  $D(.,.) : L \times L \rightarrow L$  be a symmetric mapping.  $D$  is called a *symmetric bi- $f$ -derivation* on  $L$  if there exists a function  $f : L \rightarrow L$  such that

$$D(x \wedge y, z) = (D(x, z) \wedge f(y)) \vee (f(x) \wedge D(y, z))$$

for all  $x, y, z \in L$ .

### 3. Symmetric bi-( $f, g$ )-derivations

DEFINITION 3.1. Let  $L$  be a lattice and let  $D(.,.) : L \times L \rightarrow L$  be a symmetric mapping.  $D$  is called a *symmetric bi-( $f, g$ )-derivation* on  $L$  if there exist two functions  $f, g : L \rightarrow L$  such that

$$D(x \wedge y, z) = (D(x, z) \wedge f(y)) \vee (g(x) \wedge D(y, z))$$

for all  $x, y, z \in L$ .

Obviously, a symmetric bi-( $f, g$ )-derivation  $D$  on  $L$  satisfies the relation

$$D(x, y \wedge z) = (D(x, y) \wedge f(z)) \vee (g(y) \wedge D(x, z))$$

for all  $x, y, z \in L$ .

EXAMPLE 3.1. Let  $L = \{0, 1, 2\}$  be a lattice of following Figure 1 and define mappings  $D$  and  $f, g$  on  $L$  by

$$D(x, y) = \begin{cases} 1 & \text{if } (x, y) = (0, 0) \\ 1 & \text{if } (x, y) = (0, 1) \\ 1 & \text{if } (x, y) = (1, 0) \\ 0 & \text{if } (x, y) = (0, 2) \\ 0 & \text{if } (x, y) = (2, 0) \\ 0 & \text{if } (x, y) = (1, 1) \\ 0 & \text{if } (x, y) = (2, 2) \\ 0 & \text{if } (x, y) = (1, 2) \\ 0 & \text{if } (x, y) = (2, 1) \end{cases}$$

and

$$f(x) = \begin{cases} 1 & \text{if } x = 0 \\ 2 & \text{if } x = 1 \\ 2 & \text{if } x = 2, \end{cases} \quad g(x) = \begin{cases} 0 & \text{if } x = 0 \\ 1 & \text{if } x = 1 \\ 1 & \text{if } x = 2 \end{cases}$$



FIGURE 1

Then it is easily checked that  $D$  is a symmetric bi- $(f, g)$ -derivation of a lattice  $L$ . But  $D$  is not a symmetric bi-derivation since

$$1 = D(0 \wedge 0, 0) \neq (D(0, 0) \wedge 0) \vee (0 \wedge D(0, 0)) = (1 \wedge 0) \vee (0 \wedge 1) = 0.$$

PROPOSITION 3.1. Let  $L$  be a lattice and  $d$  a trace of a symmetric bi- $(f, g)$ -derivation  $D$ . Then

$$d(x) \leq f(x) \vee g(x)$$

for all  $x \in L$ .

*Proof.* Since  $x \wedge x = x$  for all  $x \in L$ , we have

$$d(x) = D(x, x) = D(x \wedge x, x) = (D(x, x) \wedge f(x)) \vee (g(x) \wedge D(x, x)).$$

Since  $D(x, x) \wedge f(x) \leq f(x)$  and  $D(x, x) \wedge g(x) \leq g(x)$ , we get  $d(x) \leq f(x) \vee g(x)$ .  $\square$

PROPOSITION 3.2. *Let  $L$  be a lattice and let  $D$  be a symmetric bi-( $f, g$ )-derivation on  $L$ . Then  $D(x, y) \leq f(x) \vee g(x)$  and  $D(x, y) \leq f(y) \vee g(y)$  for all  $x, y \in L$ .*

*Proof.* Since  $x \wedge x = x$  for all  $x \in L$ , we have for all  $y \in L$ ,

$$D(x, y) = D(x \wedge x, y) = (D(x, y) \wedge f(x)) \vee (g(x) \wedge D(x, y)).$$

Since  $D(x, y) \wedge f(x) \leq f(x)$  and  $D(x, y) \wedge g(x) \leq g(x)$ , we have  $D(x, y) \leq f(x) \vee g(x)$ . Similarly,  $D(x, y) \leq f(y) \vee g(y)$  for all  $x, y \in L$ .  $\square$

COROLLARY 3.1. *Let  $L$  be a lattice and let  $D$  be a symmetric bi-( $f, g$ )-derivation on  $L$ . If  $g(x) \leq f(x)$  for all  $x \in L$ , then  $D(x, y) \leq f(x)$  and  $D(x, y) \leq f(y)$  for all  $x, y \in L$ .*

PROPOSITION 3.3. *Let  $L$  be a lattice and let  $D$  be a symmetric bi-( $f, g$ )-derivation on  $L$ . If  $L$  has a least element  $0$  such that  $f(0) = 0$  and  $g(0) = 0$ , we have  $D(0, y) = 0$ .*

*Proof.* For all  $x, y \in L$ , we have  $D(x, y) \leq f(x) \vee g(x)$  from Proposition 3.4. Since  $0$  is the least element of a lattice  $L$ , we get

$$0 \leq D(0, y) \leq f(0) \vee g(0) = 0,$$

which implies  $D(0, y) = 0$ .  $\square$

PROPOSITION 3.4. *Let  $L$  be a lattice and let  $D$  be a symmetric bi-( $f, g$ )-derivation on  $L$  where  $g(x) \leq f(x)$  for all  $x \in L$ . Then the following identities hold for all  $x, y, w \in L$ :*

- (1)  $D(x, y) \wedge D(w, y) \leq D(x \wedge w, y) \leq D(x, y) \vee D(w, y)$ .
- (2)  $D(x \wedge w, y) \leq f(x) \vee f(w)$ .

*Proof.* (1) For all  $x, y, w \in L$ , we have

$$D(x \wedge w, y) = (D(x, y) \wedge f(w)) \vee (g(x) \wedge D(w, y)),$$

which implies  $D(x, y) \wedge f(w) \leq D(x \wedge w, y)$ . Since  $D(w, y) \leq f(w)$  for all  $y \in L$ , we have  $D(x, y) \wedge D(w, y) \leq D(x, y) \wedge f(w)$ . Hence we get  $D(x, y) \wedge D(w, y) \leq D(x \wedge w, y)$ . Since  $D(x, y) \wedge f(w) \leq D(x, y)$  and  $g(x) \wedge D(w, y) \leq D(w, y)$ , we have  $D(x \wedge w, y) \leq D(x, y) \vee D(w, y)$ , which implies  $D(x, y) \wedge D(w, y) \leq D(x \wedge w, y) \leq D(x, y) \vee D(w, y)$ .

(2) Since  $D(x, y) \wedge f(w) \leq f(w)$  and  $g(x) \wedge D(w, y) \leq f(x) \wedge D(w, y) \leq f(x)$ , we get

$$(D(x, y) \wedge f(w)) \vee (g(x) \wedge D(w, y)) \leq f(x) \vee f(w).$$

$\square$

PROPOSITION 3.5. *Let  $L$  be a lattice with a greatest element  $1$  and let  $D$  be a symmetric bi- $(f, g)$ -derivation on  $L$  such that  $f(1) = g(1) = 1$ . Then the following properties hold for all  $x, y \in L$ :*

- (1) *If  $f(x) \leq D(1, y)$  and  $g(x) \leq D(1, y)$ , then  $D(x, y) = f(x) \vee g(x)$ .*
- (2) *If  $g(x) \geq D(1, y)$ , then  $D(x, y) \geq D(1, y)$ .*

*Proof.* (1) For all  $x, y \in L$ , we have

$$\begin{aligned} D(x, y) &= D(x \wedge 1, y) \\ &= (D(x, y) \wedge f(1)) \vee (g(x) \wedge D(1, y)) \\ &= D(x, y) \vee g(x). \end{aligned}$$

Hence we have  $g(x) \leq D(x, y)$ .

Similarly, since  $x \wedge 1 = x$ , we obtain

$$\begin{aligned} D(x, y) &= D(1 \wedge x, y) \\ &= (D(1, y) \wedge f(x)) \vee (g(1) \wedge D(x, y)) \\ &= D(x, y) \vee f(x). \end{aligned}$$

Thus we get  $f(x) \leq D(x, y)$ .

From (1) and (2), we have

$$f(x) \vee g(x) \leq D(x, y).$$

From Proposition 3.4, we have  $D(x, y) \leq f(x) \vee g(x)$ . Finally, we have

$$f(x) \vee g(x) \leq D(x, y) \leq f(x) \vee g(x),$$

which implies  $D(x, y) = f(x) \vee g(x)$ .

(2) For all  $x, y \in L$ ,

$$\begin{aligned} D(x, y) &= D(x \wedge 1, y) \\ &= (D(x, y) \wedge f(1)) \vee (g(x) \wedge D(1, y)) \\ &= D(x, y) \vee D(1, y). \end{aligned}$$

Hence we have  $D(x, y) \geq D(1, y)$ . □

THEOREM 3.1. *Let  $L$  be a distribute lattice and let  $D$  be a symmetric bi- $(f, g)$ -derivation on  $L$  with the trace  $d$ . Then*

$$d(x \wedge y) = (d(x) \wedge (f(y)) \vee (g(x) \wedge d(y)) \vee ((g(x) \wedge f(y)) \wedge D(x, y))$$

for all  $x, y \in L$ .

*Proof.* For all  $x, y \in L$ , we have

$$\begin{aligned}
d(x \wedge y) &= D(x \wedge y, x \wedge y) \\
&= (D(x, x \wedge y) \wedge f(y)) \vee (g(x) \wedge D(y, x \wedge y)) \\
&= (D(x \wedge y, x) \wedge f(y)) \vee (g(x) \wedge (D(x \wedge y, y))) \\
&= \{[(D(x, x) \wedge f(y)) \vee (g(x) \wedge D(x, y))] \wedge f(y)\} \\
&\vee \{g(x) \wedge [(D(x, y) \wedge f(y)) \vee (g(x) \wedge D(y, y))]\} \\
&= \{((d(x) \wedge f(y)) \wedge f(y)) \vee ((g(x) \wedge f(y)) \wedge D(x, y))\} \\
&\vee \{((g(x) \wedge f(y)) \wedge D(x, y)) \vee ((g(x) \wedge (g(x) \wedge d(y))))\} \\
&= (d(x) \wedge f(y)) \vee (g(x) \wedge d(y)) \vee ((f(y) \wedge g(x)) \wedge D(x, y)).
\end{aligned}$$

□

**COROLLARY 3.2.** *Let  $L$  be a distribute lattice and let  $D$  be a symmetric bi-( $f, g$ )-derivation with the trace  $d$ . Then for all  $x, y \in L$ ,*

- (1)  $(g(x) \wedge f(y)) \wedge D(x, y) \leq d(x \wedge y)$ .
- (2)  $g(x) \wedge d(y) \leq d(x \wedge y)$ .
- (3)  $d(x) \wedge f(y) \leq d(x \wedge y)$ .

*Proof.* (1), (2) and (3) are easily seen from the above theorem respectively. □

**COROLLARY 3.3.** *Let  $L$  be a distribute lattice and let  $D$  be a symmetric bi-( $f, g$ )-derivation with the trace  $d$ . If  $1$  is the greatest element of  $L$ , we have  $(g(x) \wedge f(1)) \wedge D(x, 1) \leq d(x \wedge 1) = d(x)$  for all  $x \in L$  and  $g(x) \wedge d(1) \leq d(x \wedge 1) = d(x)$  for all  $x \in L$ .*

**DEFINITION 3.2.** Let  $L$  be a lattice and let  $D$  be a symmetric bi-( $f, g$ )-derivation on  $L$ .

- (1) If  $x \leq w$  implies  $D(x, y) \leq D(w, y)$ , then  $D$  is called an *isotone symmetric bi-( $f, g$ )-derivation*.
- (2) If  $D$  is one-to-one, then  $D$  is called a *monomorphic symmetric bi-( $f, g$ )-derivation*.
- (3) If  $D$  is onto, then  $D$  is called an *epic symmetric bi-( $f, g$ )-derivation*.

**THEOREM 3.2.** *Let  $L$  be a lattice and let  $D$  be a symmetric bi-( $f, g$ )-derivation on  $L$ . The following conditions are equivalent.*

- (1)  $D$  is an isotone symmetric bi-( $f, g$ )-derivation.
- (2)  $D(x, y) \vee D(w, y) \leq D(x \vee w, y)$  for all  $x, y, w \in L$ .

*Proof.* (1)  $\Rightarrow$  (2). Suppose that  $D$  is an isotone symmetric bi-( $f, g$ )-derivation on  $L$ . Since  $x \leq x \vee w$  and  $w \leq x \vee w$ , we obtain  $D(x, y) \leq$

$D(x \vee w, y)$  and  $D(w, y) \leq D(x \vee w, y)$ . Therefore,  $D(x, y) \vee D(w, y) \leq D(x \vee w, y)$ .

(2)  $\Rightarrow$  (1). Suppose that  $D(x, y) \vee D(w, y) \leq D(x \vee w, y)$  and  $x \leq w$ . Then we have

$$\begin{aligned} D(x, y) &\leq D(x, y) \vee D(w, y) \leq D(x \vee w, y) \\ &= D(w, y). \end{aligned}$$

Hence  $D$  is an isotone symmetric bi- $(f, g)$ -derivation on  $L$ . □

Let  $L$  be a lattice and let  $D$  be a symmetric bi- $(f, g)$ -derivation on  $L$ . For each  $a \in L$  and define a set  $Fix_a(L)$  by

$$Fix_a(L) = \{x \in L \mid D(x, a) = f(x)\}.$$

**PROPOSITION 3.6.** *Let  $L$  be a lattice and let  $D$  an isotone symmetric bi- $(f, g)$ -derivation on  $L$ . If  $f : L \rightarrow L$  is a lattice homomorphism and  $g(x) \leq f(x)$  for all  $x \in L$ , then  $Fix_a(L)$  is a sublattice of  $L$ .*

*Proof.* Let  $x, y \in Fix_a(L)$ . Then  $D(x, a) = f(x)$  and  $D(y, a) = f(y)$ . Then  $f(x \wedge y) = f(x) \wedge f(y) = D(x, a) \wedge D(y, a) \leq D(x \wedge y, a)$ . Hence  $D(x \wedge y, a) = f(x \wedge y)$ , that is,  $x \wedge y \in Fix_a(L)$ . Moreover, we have  $f(x \vee y) = f(x) \vee f(y) = D(x, a) \vee D(y, a) \leq D(x \vee y, a)$  by Theorem 3.2. Thus  $D(x \vee y, a) = f(x \vee y)$ , which implies  $x \vee y \in Fix_a(L)$ . □

**PROPOSITION 3.7.** *Let  $L$  be a lattice and let  $D$  be a symmetric bi- $(f, g)$ -derivation on  $L$  where  $g(x) \leq f(x)$  for all  $x \in L$ . If  $f$  is an increasing function,  $x \leq y$  and  $y \in Fix_a(L)$  imply  $D(x, a) = D(x, a) \vee g(x)$ .*

*Proof.* Let  $x \leq y$  and  $y \in Fix_a(L)$ . Then we have  $D(x, a) \leq f(x) \leq f(y)$  and  $g(x) \leq f(x) \leq f(y)$ . Hence we obtain

$$\begin{aligned} D(x, a) &= D(x \wedge y, a) \\ &= (D(x, a) \wedge f(y)) \vee (g(x) \wedge D(y, a)) \\ &= (D(x, a) \wedge f(y)) \vee (g(x) \wedge f(y)) \\ &= D(x, a) \vee g(x). \end{aligned}$$

This completes the proof. □

**PROPOSITION 3.8.** *Let  $L$  be a distributive lattice and let  $D$  be a symmetric bi- $(f, g)$ -derivation of  $L$  where  $g(x) \leq f(x)$  for all  $x, y \in L$ . If  $f$  is a meet-homomorphism and  $x, y \in Fix_a(L)$ , we have  $x \wedge y \in Fix_a(L)$  for all  $x, y \in L$ .*



*Proof.* Let  $x, y \in \text{Fix}_a(L)$ . Then  $f(x) = D(x, a)$  and  $f(y) = D(y, a)$ . Hence we have

$$\begin{aligned} D(x \wedge y, a) &= (D(x, a) \wedge f(y)) \vee (g(x) \wedge D(y, a)) \\ &= (f(x) \wedge f(y)) \vee (g(x) \wedge f(y)) \\ &= (f(x) \vee g(x)) \wedge f(y) \\ &= f(x) \wedge f(y) \\ &= f(x \wedge y), \end{aligned}$$

which implies  $x \wedge y \in \text{Fix}_a(L)$ .  $\square$

**PROPOSITION 3.9.** *Let  $L$  be a lattice and let  $D$  be an isotone symmetric bi-( $f, g$ )-derivation on  $L$  where  $g(x) \leq f(x)$  for all  $x \in L$ . If  $x, y \in \text{Fix}_a(L)$  and  $f$  is an increasing function, then  $x \vee y \in \text{Fix}_a(L)$ .*

*Proof.* Since  $x \leq x \vee y$  and  $y \leq x \vee y$ , we have  $f(x \vee y) \leq f(x)$  and  $f(x \vee y) \leq f(y)$  respectively. Hence we obtain  $f(x \vee y) \leq f(x) \vee f(y) = D(x, a) \vee D(y, a) \leq D(x \vee y, a)$  since  $D$  is an isotone symmetric bi-( $f, g$ )-derivation. From Proposition 3.4 (2), we have  $D(x \vee y, a) \leq f(x \vee y)$ , which implies  $D(x \vee y, a) = f(x \vee y)$ . Hence  $x \vee y \in \text{Fix}_a(L)$ .  $\square$

**PROPOSITION 3.10.** *Let  $L$  be a lattice,  $D$  a symmetric bi-( $f, g$ )-derivation on  $L$  where  $f(x) \leq g(x)$  and  $1$  the greatest element of  $L$ . Then the following identities hold.*

- (1) *If  $g(x) \leq D(1, y)$  and  $f(1) = 1$ , then  $D(x, y) = g(x)$ .*
- (2) *If  $g(x) \geq D(1, y)$  and  $f(1) = 1$ , then  $D(x, y) \geq D(1, y)$ .*

*Proof.* (1) Let  $g(x) \leq D(1, y)$ . Then we have  $D(x, y) \leq f(x) \vee g(x) = g(x)$ , and so

$$\begin{aligned} D(x, y) &= D(x \wedge 1, y) \\ &= (D(x, y) \wedge f(1)) \vee (g(x) \wedge D(1, y)) \\ &= D(x, y) \vee g(x) \\ &= g(x). \end{aligned}$$

(2) Let  $g(x) \geq D(1, y)$ . Then we have

$$\begin{aligned} D(x, y) &= D(x \wedge 1, y) \\ &= (D(x, y) \wedge f(1)) \vee (g(x) \wedge D(1, y)) \\ &= D(x, y) \vee D(1, y). \end{aligned}$$

Hence we obtain  $D(1, y) \leq D(x, y)$  for all  $x, y \in L$ .  $\square$

**THEOREM 3.3.** *Let  $L$  be a lattice with the greatest element 1 and let  $D$  be an isotone symmetric bi- $(f, g)$ -derivation on  $L$ . Let  $f(1) = g(1) = 1$  and either  $f(x) \geq g(x)$  or  $f(x) \leq g(x)$  for all  $x \in L$ . Then*

$$D(x, y) = (f(x) \vee g(x)) \wedge D(1, y)$$

for all  $x, y, z \in L$ .

*Proof.* Suppose that  $D$  is an isotone symmetric bi- $(f, g)$ -derivation on  $L$ . Then  $D(x, y) \leq D(1, y)$  for all  $x, y \in L$ . Now let  $g(x) \leq f(x)$  for  $x \in L$ . Then we have  $D(x, y) \leq g(x) \vee f(x) = f(x)$ . From this, we get  $D(x, y) \leq f(x) \wedge D(1, y)$ . Also, we obtain

$$\begin{aligned} D(x, y) &= D((x \vee 1) \wedge x, y) \\ &= [(D(x \vee 1), y) \wedge f(x)] \vee [g(x \vee 1) \wedge D(x, y)] \\ &= [D(1, y) \wedge f(x)] \vee [g(1) \wedge D(x, y)] \\ &= [D(D(1, y) \wedge f(x)) \vee [1 \wedge D(x, y)]] \\ &= [D(1, y) \wedge f(x)] \vee D(x, y) \\ &= D(1, y) \wedge f(x). \end{aligned}$$

Since  $f(x) \vee g(x) = f(x)$ , we have

$$D(x, y) = (f(x) \vee g(x)) \wedge D(1, y).$$

Now suppose that  $f(x) \leq g(x)$  for  $x \in L$ . Similarly, we have  $D(x, y) \leq f(x) \vee g(x) = g(x)$ . From this, we have  $D(x, y) \leq g(x) \wedge D(1, y)$ . Also, we obtain

$$\begin{aligned} D(x, y) &= D(x \wedge (x \vee 1), y) \\ &= [(D(x, y) \wedge f(x \vee 1))] \vee [g(x) \wedge D((x \vee 1), y)] \\ &= [D(x, y) \wedge f(1)] \vee [g(x) \wedge D(1, y)] \\ &= [D(D(x, y) \wedge 1)] \vee [g(x) \wedge D(1, y)] \\ &= D(x, y) \vee [g(x) \wedge D(1, y)] \\ &= g(x) \wedge D(1, y). \end{aligned}$$

Since  $f(x) \vee g(x) = g(x)$ , we have

$$D(x, y) = (f(x) \vee g(x)) \wedge D(1, y).$$

This completes the proof.  $\square$

Let  $D$  be a symmetric bi- $(f, g)$ -derivation of  $L$  and let 0 be a least element of  $L$ . Define a set  $KerD$  by

$$KerD = \{x \in L \mid D(x, 0) = 0\}.$$

PROPOSITION 3.11. *Let  $L$  be a lattice with a least element  $0$  and let  $D$  be a symmetric bi-( $f, g$ )-derivation on  $L$ . If  $x, y \in \text{Ker}D$ , then  $x \wedge y \in \text{Ker}D$ .*

*Proof.* Let  $x, y \in \text{Ker}D$ . Then  $D(x, 0) = D(y, 0) = 0$ . Hence we have

$$\begin{aligned} D(x \wedge y, 0) &= (D(x, 0) \wedge f(y)) \vee (g(x) \wedge D(y, 0)) \\ &= (0 \wedge f(x)) \vee (g(x) \wedge 0) \\ &= 0 \vee 0 = 0, \end{aligned}$$

which implies  $x \wedge y \in \text{Ker}D$ .  $\square$

PROPOSITION 3.12. *Let  $L$  be a lattice with a least element  $0$  and let  $D$  be an isotone symmetric bi-( $f, g$ )-derivation on  $L$ . If  $x \leq y$  and  $y \in \text{Ker}D$ , then  $x \in \text{Ker}D$ .*

*Proof.* Let  $y \in \text{Ker}D$ . Then  $D(y, 0) = 0$  and  $D(x, 0) \leq D(y, 0) = 0$  since  $D$  is isotone. Hence we have  $D(x, 0) = 0$ , and so

$$\begin{aligned} D(x, 0) &= D(x \wedge y, 0) = (D(x, 0) \wedge f(y)) \vee (g(x) \wedge D(y, 0)) \\ &= (0 \wedge f(x)) \vee (g(x) \wedge 0) \\ &= 0 \vee 0 = 0, \end{aligned}$$

which implies  $x \in \text{Ker}D$ .  $\square$

## References

- [1] R. Balbes and P. Dwinger, *Distributive Lattices*, University of Missouri Press, Columbia, United States, 1974.
- [2] A. J. Bell, The co-information lattice, in: 4th International Symposium on Independent Component Analysis and Blind Signal Separation (ICA2003), Nara, Japan, 2003, 921-926.
- [3] H. E. Bell and L. C. Kappe, *Rings in wich derivations satisfy certain algebraic conditions*, Acta Math. Hungar. **53** (1989), no. 3-4, 339-346.
- [4] H. E. Bell and G. Mason, *On derivations in near-rings and near-fields*, North-Holland Math. Studies **137** (1987), 31-35.
- [5] G. Birkhoof, *Lattice Theory*, American Mathematical Society Colloquium, 1940.
- [6] C. Carpineto and G. Romano, *Information retrieval through hybrid navigation of lattice representations*, Int. J. Human-Computers Studies **45** (1996), 553-558.
- [7] Y. Çeven and M. A. Öztürk, *On the trace of a permuting tri-additive mapping in left  $s$ -unital rings*, International Journal of Pure and Applied Mathematics **23** (2005), no. 4, 465-474.
- [8] Y. Çeven and M. A. Öztürk, *On  $f$ -derivations of lattices*, Bull. Korean Math. Soc. **45** (2008), no. 4, 701-707.
- [9] Y. Çeven, *Symmetric bi-derivations of Lattices*, Quaestiones Mathematicae **32** (2009), no. 2, 1-5.

- [10] C. Degang, Z. Wenxiu, D. Yeung, and E. C. C. Tsang, *Rough approximations on a complete distributive lattice with applications to generalized rough sets*, Informat. Sci. **176** (2006), 1829-1848.
- [11] L. Ferrari, *On derivations of lattices*, Pure Math. Appl. **12** (2001), no. 4, 365-382.
- [12] A. Honda and M. Grabish, *Entropy of capacities on lattices and set systems*, Inform. Sci. **176** (2006), 3472-3489.
- [13] Y. B. Jun and X. L. Xin, *On derivations of BCI-algebras*, Inform. Sci. **159** (2004), 167-176.
- [14] F. Karacal, *On the direct decomposability of strong negations and S-implication operators on product lattices*, Informat. Sci. **176** (2006), 3011-3025.
- [15] D. Özden and M. A. Öztürk, *Permuting tri-derivations in prime and semi-prime gamma rings*, Kyungpook Math. J. **46** (2006), 153-167.
- [16] M. A. Öztürk, *Permuting Tri-derivations in Prime and Semi-prime Rings*, East Asian Math. J. **15** (1999), no. 2, 177-190.
- [17] M. A. Öztürk, Y. Çeven, and Y. B. Jun, *Generalized derivations of BCI-algebras*, Honam Math. J. **31** (2009), no. 4, 601-609.
- [18] M. A. Öztürk, H. Yazarlı and K. H. Kim, *Permuting tri-derivations in lattices*, Quaestiones Mathematicae **32** (2009), no. 3, 415-425.
- [19] E. Posner, *Derivations in prime rings*, Proc. Am. Math. Soc. **8** (1957), 1093-1100.
- [20] R. S. Sandhu, *Role hierarchies and constraints for lattice-based access controls in: Proceedings of the 4th European Symposium on Research in Computer Security*, Rome, Italy, 1996, 65-79.
- [21] G. Szász, *Derivations of lattices*, Acta Sci. Math. (Szeged) **37** (1975), 149-154.
- [22] X. L. Xin, T. Y. Li, and J. H. Lu, *On derivations of lattices*, Information Sciences **178** (2008), no. 2, 307-316.
- [23] H. Yazarlı, M. A. Öztürk, and Y. B. Jun, *Tri-additive maps and permuting tri-derivations*, Commun. Fac. Sci. Univ. Ank. Series A1 **54** (2005), no. 1, 1-8.
- [24] J. Zhan and Y. L. Liu, *On f-derivations of BCI-algebras*, Int. J. of Mathematics and Mathematical Sciences, 2005, 1675-1684.

\*

Department of Mathematics  
Korea National University of transportation  
Chungju 380-702, Republic of Korea  
*E-mail*: ghkim@ut.ac.kr

\*\*

Department of Mathematics  
Dankook University  
Cheonan 330-714, Republic of Korea  
*E-mail*: yonghoon@dankook.ac.kr