

PRINCIPAL COFIBRATIONS AND GENERALIZED CO- H -SPACES

YEON SOO YOON*

ABSTRACT. For a map $p : X \rightarrow A$, there are concepts of co- H^p -spaces, co- T^p -spaces, which are generalized ones of co- H -spaces [17,18]. For a principal cofibration $i_r : X \rightarrow C_r$ induced by $r : X' \rightarrow X$ from $\iota : X' \rightarrow cX'$, we obtain some sufficient conditions to having extensions co- $H^{\bar{p}}$ -structures and co- $T^{\bar{p}}$ -structures on C_r of co- H^p -structures and co- T^p -structures on X respectively. We can also obtain some results about co- H^p -spaces and co- T^p -spaces in homology decompositions for spaces, which are generalizations of Golasinski and Klein's result about co- H -spaces.

1. Introduction

A map $f : X \rightarrow B$ is *cocyclic* [13] if there is a map $\theta : X \rightarrow X \vee B$ such that $j\theta \sim (1 \times f)\Delta$, where $j : X \vee B \rightarrow X \times B$ is the inclusion and $\Delta : X \rightarrow X \times X$ is the diagonal. It is clear that a space X is a co- H -space if and only if the identity map 1_X of X is cocyclic. We called a space X as a *co- H^p -space for a map $p : X \rightarrow A$* [17] if there is a cocyclic map $p : X \rightarrow A$, that is, there is a co- H^p -structure $\theta : X \rightarrow X \vee A$ such that $j\theta \sim (1 \times p)\Delta$, where $j : X \vee A \rightarrow X \times A$ is the inclusion and $\Delta : X \rightarrow X \times X$ is the diagonal. It is clear that if X is a co- H -space, then for any map $p : X \rightarrow A$, X is a co- H^p -space. In Example 2.4, there is a space Q_p which is a co- H^δ -space, but not a co- H -space. Let τ be the adjoint functor from the group $[\Sigma X, Y]$ to the group $[X, \Omega Y]$. The symbols e and e' denote $\tau^{-1}(1_{\Omega X})$ and $\tau(1_{\Sigma X})$ respectively. In [1], Aguade introduced a T -space as a space X having the property that the evaluation fibration $\Omega X \rightarrow X^{S^1} \rightarrow X$ is fibre homotopically trivial. It is well known [1] that a space X is a T -space if and only if the evaluating

Received January 11, 2017; Accepted January 20, 2017.

2010 Mathematics Subject Classification: 55P45, 55P35.

Key words and phrases: co- H^p -spaces, co- T^p -spaces for maps, Homology decompositions.

This work was supported by Hannam University Research Fund, 2016.

map $e : \Sigma\Omega X \rightarrow X$ is cyclic. As a dual space of T -space, we introduced [14] that a space X is a co- T -space if $e' : X \rightarrow \Omega\Sigma X$ is cocyclic. A space X is called [5] a G' -space if $G^n(X) = H^n(X)$ for all n . It is clear that any co- H -space is a co- T -space, and any co- T -space is a G' -space. It is known [14] that $\mathbb{R}P^2$ is a G' -space, but not a co- T -space. We called a space X as a *co- T^p -space for a map $p : X \rightarrow A$* [18] if $e' : X \rightarrow \Omega\Sigma X$ is p -cocyclic, that is, there is a co- T^p -structure $\theta : X \rightarrow \Omega\Sigma X \vee A$ such that $j\theta \sim (e' \times p)\Delta$, where $j : \Omega\Sigma X \vee A \rightarrow \Omega\Sigma X \times A$ is the inclusion and $\Delta : X \rightarrow X \times X$ is the diagonal map. It is shown [18] that X is a co- T -space if and only if for any space A and any map $p : X \rightarrow A$, X is a co- T^p -space for a map $p : X \rightarrow A$. We called a space X as a *G'_p -space for a map $p : X \rightarrow A$* [19] if $e' : X \rightarrow \Omega\Sigma X$ is weakly p -cocyclic, that is, $e^*(H^n(\Omega\Sigma X)) \subset G^n(X, p, A)$ for all n . For a map $p : X \rightarrow A$, there are concepts of co- H^p -spaces, co- T^p -spaces and G'_p -spaces which are generalized ones of co- H -spaces. In general, any co- H -space is a co- H^p -space, any co- H^p -space is a co- T^p -space and any co- T^p -space is a G'_p -space. In [19], we already studied about some properties of G'_p -spaces for maps and their homology decompositions.

In this paper, we study about relationships between co- H^p -spaces, co- T^p -spaces and their homology decompositions respectively. For a principal cofibration $i_r : X \rightarrow C_r$ induced by $r : X' \rightarrow X$ from $\iota_{X'} : X' \rightarrow cX'$, we obtain some sufficient conditions to having extendings co- $H^{\bar{p}}$ -structures and co- $T^{\bar{p}}$ -structures on C_r of H^p -structures and T^p -structures on X respectively. Let X and A be rational spaces and $p : X \rightarrow A$ a map, and $\{X_n, q_n, i_n\}$ and $\{A_n, q'_n, i'_n\}$ homology decompositions for X and A respectively. Then we can obtain that X is a co- H^p -space for a map $p : X \rightarrow A$ if and only if for each n , X_n is a co- H^{p_n} -space and the all pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- H^{p_n} -primitive. Thus we have, as a corollary, that Golasinski and Klein's result about co- H -spaces. We can also obtain that X is a co- T^p -space for a map $p : X \rightarrow A$ if and only if for each n , X_n is a co- T^{p_n} -space and the all pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- T^{p_n} -primitive.

2. Dual Gottlieb sets for maps and generalized co- H -spaces

Let $p : X \rightarrow A$ be a map. A based map $f : X \rightarrow B$ is called *p -cocyclic* [10] if there is a map $\theta : X \rightarrow A \vee B$ such that the following diagram is

homotopy commutative;

$$\begin{array}{ccc} X & \xrightarrow{\theta} & A \vee B \\ \Delta \downarrow & & j \downarrow \\ X \times X & \xrightarrow{(p \times f)} & A \times B, \end{array}$$

where $j : A \vee B \rightarrow A \times B$ is the inclusion and $\Delta : X \rightarrow X \times X$ is the diagonal map. We call such a map θ a *coassociated map* of a p -cocyclic map f .

In the case $p = 1_X : X \rightarrow X$, $f : X \rightarrow B$ is called *cocyclic* [13]. Clearly any cocyclic map is a p -cocyclic map and also $f : X \rightarrow B$ is p -cocyclic iff $p : X \rightarrow A$ is f -cocyclic. The *dual Gottlieb set* $DG(X, p, A; B)$ for a map $p : X \rightarrow A$ [16] is the set of all homotopy classes of p -cocyclic maps from X to B . In the case $p = 1_X : X \rightarrow X$, we called such a set $DG(X, 1, X; B)$ the *dual Gottlieb set* [13] denoted $DG(X; B)$, that is, the dual Gottlieb set is exactly same with the dual Gottlieb set for the identity map. We denote $DG(X, p, A; K(\pi, n))$ by $G^n(X, p, A; \pi)$ and $DG(X, p, A; K(\mathbb{Z}, n))$ by $G^n(X, p, A)$, $DG(X; K(\mathbb{Z}, n))$ by $G^n(X)$. Haslam [5] introduced and studied the *coevaluation subgroups* $G^n(X; \pi)$ of $H^n(X; \pi)$. $G^n(X; \pi)$ is defined to be the set of all homotopy classes of cocyclic maps from X to $K(\pi, n)$. A space X is called [5] a G^l -space if $G^n(X) = H^n(X)$ for all n .

In general, $DG(X; B) \subset DG(X, p, A; B) \subset [X, B]$ for any map $p : X \rightarrow B$ and any space B . It is known [16] that for any n , $G^n(S^n \times S^n; \mathbb{Z}) \neq G^n(S^n \times S^n, p_1, S^n; \mathbb{Z}) \neq H^n(S^n \times S^n; \mathbb{Z})$.

The next proposition is an immediate consequence from the definition.

PROPOSITION 2.1. [17]

- (1) For any maps $g : X \rightarrow A$, $h : A \rightarrow B$ and any space C , $DG(X, g, A; C) \subset DG(X, hg, B; C)$.
- (2) $DG(X, B) = DG(X, 1_X, X; B) \subset DG(X, g, A; B) \subset DG(X, *, A; B) = [X, B]$ for any spaces X, A and B .
- (3) $DG(X, B) = \cap \{DG(X, g, A; B) | g : X \rightarrow A \text{ is a map and } A \text{ is a space}\}$.
- (4) If $h : A \rightarrow B$ is a homotopy equivalence, then $DG(X, g, A; C) = DG(X, hg, B; c)$.
- (5) For any map $k : Y \rightarrow X$, $k^*(DG(X, g, A; B)) \subset DG(Y, gk, A; B)$.
- (6) For any map $k : Y \rightarrow X$, $k^*(DG(X; B)) \subset DG(Y, k, X; B)$.
- (7) For any map $s : B \rightarrow C$, $s_*(DG(X, g, A; B)) \subset DG(X, g, A; C)$.

PROPOSITION 2.2.

- (1) [9] X is a co- H -space $\iff DG(X, B) = [X, B]$ for any space B .
- (2) [14] X is a co- T -space $\iff DG(X, \Omega C) = [X, \Omega C]$ for any space C .
- (3) [5] X is a G' -space $\iff G^n(X) = H^n(X)$ for all n .

It is clear that any co- H -space is a co- T -space and any co- T -space is a G' -space. It is known [14] that $\mathbb{R}P^2$ is a G' -space, but not a co- H -space and co- T -space.

PROPOSITION 2.3. *Let $p : X \rightarrow A$ be a map. Then*

- (1) [17] X is a co- H^p -space $\iff DG(X, p, A; B) = [X, B]$ for any space B .
- (2) [18] X is a co- T^p -space $\iff DG(X, p, A; \Omega C) = [X, \Omega C]$ for any space C .
- (3) [19] X is a G'_p -space $\iff G^n(X, p, A) = H^n(X)$ for all n .

Thus we know that any co- H -space is a co- H^p -space, any co- H^p -space is a co- T^p -space and any co- T^p -space is a G'_p -space for any map $p : X \rightarrow A$.

The following example says that there is a space which is a co- H^p -space, but not a co- H -space.

EXAMPLE 2.4. *For any odd prime p , let $[f]$ be the generator of p -primary summand of $\pi_{4p-3}(S^2)$ which is isomorphic $\mathbb{Z}/p\mathbb{Z}$. Then it is known [7] that for $Q_p = S^2 \cup_f e^{4p-2}$, $\text{cat } Q_p = 2$. It is also well known fact that a space X is a co- H -space if and only if $\text{cat } X \leq 1$. Thus we know that Q_p is not a co- H -space. It is also known [6, Proposition 15.8] that for a cofibration sequence $S^{4p-3} \xrightarrow{f} S^2 \xrightarrow{i} Q_p \xrightarrow{\delta} S^{4p-2} \rightarrow \dots$, $\delta : Q_p \rightarrow S^{4p-2}$ is a cocyclic map. Moreover, it is known [16] that $p : X \rightarrow A$ is a cocyclic map if and only if $DG(X, p, A; B) = [X, B]$ for any space B . Thus we know that Q_p is a co- H^δ -space.*

3. Principal cofibrations and generalized co- H -spaces

Given maps $p : X \rightarrow A$, $p' : X' \rightarrow A'$, let $(s, r) : p' \rightarrow p$ be a map from p' to p , that is, the following diagram is commutative;

$$\begin{array}{ccc} X' & \xrightarrow{p'} & A' \\ r \downarrow & & s \downarrow \\ X & \xrightarrow{p} & A. \end{array}$$

It is a well known fact that $Y \xrightarrow{\iota} cY \rightarrow \Sigma Y$ is a cofibration, where $\iota(y) = [y, 1]$. Let $i_r : X \rightarrow C_r$ be the cofibration induced by $r : X' \rightarrow X$ from $\iota_{X'} : X' \rightarrow cX'$. Let $i_s : A \rightarrow C_s$ be the cofibration induced by $s : A' \rightarrow A$ from $\iota_{A'} : A' \rightarrow cA'$. Then there is a map $\bar{p} : C_t \rightarrow C_s$ such that the following diagram is commutative

$$\begin{array}{ccc} X & \xrightarrow{p} & A \\ i_r \downarrow & & i_s \downarrow \\ C_r & \xrightarrow{\bar{p}} & C_s, \end{array}$$

where $C_r = cX' \amalg X/[x', 1] \sim r(x')$, and $C_s = cA' \amalg A/[a', 1] \sim s(a')$, $\bar{p} : C_r \rightarrow C_s$ is given by $\bar{p}([x', t]) = [p'(x'), t]$ if $[x', t] \in cX'$ and $\bar{p}(x) = p(x)$ if $x \in X$, $i_r(x) = x$, $i_s(a) = a$.

DEFINITION 3.1. Let X be a co- H^p -space for a map $p : X \rightarrow A$. A map $(s, r) : p' \rightarrow p$ is called a *co- H^p -primitive* if there is a map $\theta : X \rightarrow A \vee X$ such that $j\theta \sim (p \times 1)\Delta$ and $(i_s \vee i_r)\theta r \sim * : X' \rightarrow C_s \vee C_r$, where $j : A \vee X \rightarrow A \times X$ is the inclusion.

DEFINITION 3.2. Let X be a co- T^p -space for a map $p : X \rightarrow A$. A map $(s, r) : p' \rightarrow p$ is called a *co- T^p -primitive* if there is a map $\theta : X \rightarrow A \vee \Omega\Sigma X$ such that $j\theta \sim (p \times e')\Delta$ and $(i_s \vee \Omega\Sigma i_r)\theta r \sim * : X' \rightarrow C_s \vee \Omega\Sigma C_r$, where $j : A \vee \Omega\Sigma X \rightarrow A \times \Omega\Sigma X$ is the inclusion.

DEFINITION 3.3. [19] Let X be a G'_p -space for a map $p : X \rightarrow A$. A map $(s, r) : p' \rightarrow p$ is called a *G'_p -primitive* if for each map $g : \Omega\Sigma X \rightarrow K(\mathbb{Z}, m)$, m arbitrary, there is a map $G : X \rightarrow A \vee K(\mathbb{Z}, m)$ such that $jG \sim (p \times g \circ e'_X)\Delta$ and $(i_s \vee 1)Gr \sim * : X' \rightarrow C_s \vee K(\mathbb{Z}, m)$, where $j : A \vee K(\mathbb{Z}, m) \rightarrow A \times K(\mathbb{Z}, m)$ is the inclusion and $e'_X : X \rightarrow \Omega\Sigma X$ is the adjoint functor image, $\tau(1_{\Sigma X})$, of $1_{\Sigma X}$.

PROPOSITION 3.4.

- (1) If X is a co- H^p -space for a map $p : X \rightarrow A$ and $(s, r) : p' \rightarrow p$ is a co- H^p -primitive, then $(s, r) : p' \rightarrow p$ is a co- T^p -primitive.
- (2) If X is a co- T^p -space for a map $p : X \rightarrow A$ and $(s, r) : p' \rightarrow p$ is a co- T^p -primitive, then $(s, r) : p' \rightarrow p$ is a G'_p -primitive.

Proof.

- (1) Since $(s, r) : p' \rightarrow p$ is a co- H^p -primitive, there is a map $\theta : X \rightarrow A \vee X$ such that $j\theta \sim (p \times 1)\Delta$ and $(i_s \vee i_r)\theta r \sim * : X' \rightarrow C_s \vee C_r$, where $j : A \vee X \rightarrow A \times X$ is the inclusion. Let $\theta' = (1 \vee e')\theta : X \rightarrow A \vee \Omega\Sigma X$. Then $j'\theta' \sim (1 \times e')j\theta \sim (1 \times e')(p \times 1)\Delta = (p \times e')\Delta$, where $j' : A \vee \Omega\Sigma X \rightarrow A \times \Omega\Sigma X$ is the inclusion. Moreover, since

$(i_s \vee \Omega\Sigma i_r)(1 \vee e'_X) \sim (1 \vee e'_{C_r})(i_s \vee i_r) : A \vee X \rightarrow C_s \vee \Omega\Sigma C_r$, we have that $(i_s \vee \Omega\Sigma i_r)\theta' r \sim (i_s \vee \Omega\Sigma i_r)(1 \vee e'_X)\theta r \sim (1 \vee e'_{C_r})(i_s \vee i_r)\theta r \sim (1 \vee e'_{C_r}) * \sim *$. Thus $(s, r) : p' \rightarrow p$ is a $\text{co-}T^p$ -primitive.

(2) Since $(s, r) : p' \rightarrow p$ is a $\text{co-}T^p$ -primitive, there is a map $\theta : X \rightarrow A \vee \Omega\Sigma X$ such that $j\theta \sim (p \times e')\Delta$ and $(i_s \vee \Omega\Sigma i_r)\theta r \sim * : X' \rightarrow C_s \vee \Omega\Sigma C_r$, where $j : A \vee \Omega\Sigma X \rightarrow A \times \Omega\Sigma X$ is the inclusion. For any m , each $g : \Omega\Sigma X \rightarrow K(\mathbb{Z}, m)$, let $\theta' = (1 \vee g)\theta : X \rightarrow A \vee K(\mathbb{Z}, m)$. Then $j'\theta' \sim (1 \times g)j\theta \sim (1 \times g)(p \times e')\Delta = (p \times ge')\Delta$, where $j' : A \vee K(\mathbb{Z}, m) \rightarrow A \times K(\mathbb{Z}, m)$ is the inclusion. Moreover, since $(1 \vee \Omega\Sigma g)(i_s \vee \Omega\Sigma i_r) \sim (i_s \vee 1)(1 \vee g) : A \vee \Omega\Sigma X \rightarrow C_s \vee \Omega\Sigma K(\mathbb{Z}, m)$, we have that $(i_s \vee 1)\theta' r = (i_s \vee 1)(1 \vee g)\theta r \sim (1 \vee \Omega\Sigma g)(i_s \vee \Omega\Sigma i_r)\theta r \sim (1 \vee \Omega\Sigma g) * \sim *$. Thus $(s, r) : p' \rightarrow p$ is a G'_p -primitive. \square

LEMMA 3.5.

- (1) A map $f : X \rightarrow B$ can be extended to a map $h : C_r \rightarrow B$ with $hi_r = f$ if and only if $fr \sim *$.
- (2) [15] Given maps $g_t : C_r \rightarrow B_t (t = 1, 2)$ and $g : C_r \rightarrow B_1 \vee B_2$ satisfying $p_t j g_i r \sim g_t i_r (t = 1, 2)$, then there is a map $h : C_r \rightarrow B_1 \vee B_2$ such that $g_i r = hi_r$ and $p_t j h \sim g_t (t = 1, 2)$, where $j : B_1 \vee B_2 \rightarrow B_1 \times B_2$ is the inclusion and $p_t : B_1 \times B_2 \rightarrow B_t, t = 1, 2$ are projections.

THEOREM 3.6.

- (1) If X is a $\text{co-}H^p$ -space for a map $p : X \rightarrow A$ and $(s, r) : p' \rightarrow p$ is $\text{co-}H^p$ -primitive, then C_r is a $\text{co-}H^{\bar{p}}$ -space for a map $\bar{p} : C_r \rightarrow C_s$.
- (2) If X is a $\text{co-}T^p$ -space for a map $p : X \rightarrow A$ and $(s, r) : p' \rightarrow p$ is $\text{co-}T^p$ -primitive, then C_r is a $\text{co-}T^{\bar{p}}$ -space for a map $\bar{p} : C_r \rightarrow C_s$.

Proof.

- (1) Since $(s, r) : p' \rightarrow p$ is a $\text{co-}H^p$ -primitive, there is a map $\theta : X \rightarrow A \vee X$ such that $j\theta \sim (p \times 1)\Delta$ and $(i_s \vee i_r)\theta r \sim * : X' \rightarrow C_s \vee C_r$, where $j : A \vee X \rightarrow A \times X$ is the inclusion. From Lemma 3.5(1), there is an extending $\theta' : C_r \rightarrow C_s \vee C_r$ of $(i_s \vee i_r) \circ \theta : X \rightarrow C_s \vee C_r$, that is, $\theta' \circ i_r = (i_s \vee i_r) \circ \theta$. Then we have that $p_1 j' \theta' i_r = p_1 j' (i_s \vee i_r) \theta = p_1 (i_s \times i_r) j \theta \sim p_1 (i_s \times i_r) (p \times 1) \Delta = i_s \circ p \sim \bar{p} \circ i_r$ and $p_2 j' \theta' i_r = p_2 j' (i_s \vee i_r) \theta = p_2 (i_s \times i_r) j \theta \sim p_2 (i_s \times i_r) (p \times 1) \Delta \sim i_r \sim 1_{C_r} \circ i_r$. Thus we have, from Lemma 3.5(2), that there is a map $\bar{\theta} : C_r \rightarrow C_s \vee C_r$ such that $\bar{\theta} i_r = \theta' i_r = (i_s \vee i_r) \theta$ and $p_1 j' \bar{\theta} \sim \bar{p}$ and $p_2 j' \bar{\theta} \sim 1_{C_r}$. Thus we know that $1 : C_r \rightarrow C_r$ is \bar{p} -cocyclic and C_r is a $\text{co-}H^{\bar{p}}$ -space for a map $\bar{p} : C_r \rightarrow C_s$. This proves the theorem.
- (2) Since $(s, r) : p' \rightarrow p$ is a $\text{co-}T^p$ -primitive, there is a map $\theta : X \rightarrow A \vee \Omega\Sigma X$ such that $j\theta \sim (p \times e')\Delta$ and $(i_s \vee \Omega\Sigma i_r)\theta r \sim * : X' \rightarrow C_s \vee \Omega\Sigma C_r$, where $j : A \vee \Omega\Sigma X \rightarrow A \times \Omega\Sigma X$ is the inclusion. From Lemma

3.5(1), there is an extending $\theta' : C_r \rightarrow C_s \vee \Omega\Sigma C_r$ of $(i_s \vee \Omega\Sigma i_r) \circ \theta : X \rightarrow C_s \vee \Omega\Sigma C_r$, that is, $\theta' \circ i_r = (i_s \vee \Omega\Sigma i_r) \circ \theta$. Then we have that $p_1 j' \theta' i_r = p_1 j' (i_s \vee \Omega\Sigma i_r) \theta = p_1 (i_s \times \Omega\Sigma i_r) j \theta \sim p_1 (i_s \times \Omega\Sigma i_r) (p \times e') \Delta = i_s \circ p \sim \bar{p} \circ i_r$ and $p_2 j' \theta' i_r = p_2 j' (i_s \vee \Omega\Sigma i_r) \theta = p_2 (i_s \times \Omega\Sigma i_r) j \theta \sim p_2 (i_s \times \Omega\Sigma i_r) (p \times e') \Delta \sim \Omega\Sigma i_r e'_X \sim e'_{C_r} \circ i_r$. Thus we have, from Lemma 3.5(2), that there is a map $\bar{\theta} : C_r \rightarrow C_s \vee \Omega\Sigma C_r$ such that $\bar{\theta} i_r = \theta' i_r = (i_s \vee \Omega\Sigma i_r) \theta$ and $p_1 j' \bar{\theta} \sim \bar{p}$ and $p_2 j' \bar{\theta} \sim e'_{C_r}$. Thus we know that $e'_{C_r} : C_r \rightarrow \Omega\Sigma C_r$ is \bar{p} -cocyclic and C_r is a co- $T^{\bar{p}}$ -space for a map $\bar{p} : C_r \rightarrow C_s$. This proves the theorem. \square

In 1959, Eckmann and Hilton [2] introduced a dual concept of Postnikov system as follows; A *homology decomposition* of X consists of a sequence of spaces and maps $\{X_n, q_n, i_n\}$ satisfying (1) $q_n : X_n \rightarrow X$ induces an isomorphism $(q_n)_* : H_i(X_n) \rightarrow H_i(X)$ for $i \leq n$ and $H_i(X_n) = 0$ for $i > n$, (2) $i_n : X_n \rightarrow X_{n+1}$ is a cofibration with cofiber $M(H_{n+1}(X), n)$ (a Moore space of type $(H_{n+1}(X), n)$), (3) $q_n \sim q_{n+1} \circ i_n$. It is known by [6] that if X be a 1-connected space having the homotopy type of CW complex, then there is a homology decomposition $\{X_n, q_n, i_n\}$ of X such that $i_n : X_n \rightarrow X_{n+1}$ is the principal cofibration induced from $\iota : M(H_{n+1}(X), n) \rightarrow cM(H_{n+1}(X), n)$ by a map $\kappa'_n : M(H_{n+1}(X), n) \rightarrow X_n$ which is called the dual Postnikov invariants. A space X is called a *rational space* [11] if X is a 1-connected space having homotopy type of a CW -complex such that for each $n > 0$, $H_n(X, \mathbb{Z})$ is a finite dimensional vector space over \mathbb{Q} . It is well known [11] that if X and A are rational spaces and $p : X \rightarrow A$ is a based map, then there exist homology decompositions $\{X_n, q_n, i_n\}$ and $\{A_n, q'_n, i'_n\}$ for X and A respectively and induced maps $\{p_n : X_n \rightarrow A_n\}$ satisfying

(1) for each n , the following diagram is homotopy commutative

$$\begin{array}{ccc} M(H_{n+1}(X), n) & \xrightarrow{\tilde{p}_*} & M(H_{n+1}(A), n) \\ k'_n(X) \downarrow & & k'_n(A) \downarrow \\ X_n & \xrightarrow{p_n} & A_n \end{array}$$

, that is, $(k'_n(A), k'_n(X)) : \tilde{p}_\# \rightarrow p_n$ is a map,

(2) $p_{n+1} : X_{n+1} \rightarrow A_{n+1}$ given by $p_{n+1} = \bar{p}_n$ satisfying commute diagram

$$\begin{array}{ccc} X_n & \xrightarrow{p_n} & A_n \\ i_n (= \iota_{k'_n(X)}) \downarrow & & i'_n (= \iota_{k'_n(A)}) \downarrow \\ X_{n+1} & \xrightarrow{p_{n+1}} & A_{n+1}, \end{array}$$

(3) for each n , the following diagram is homotopy commutative

$$\begin{array}{ccc} X_n & \xrightarrow{p_n} & A_n \\ q_n \downarrow & & q'_n \downarrow \\ X & \xrightarrow{p} & A. \end{array}$$

THEOREM 3.7. *Let X and A be rational spaces and $p : X \rightarrow A$ a map, and $\{X_n, q_n, i_n\}$ and $\{A_n, q'_n, i'_n\}$ homology decompositions for X and A respectively.*

- (1) *If X is a co- H^p -space for a map $p : X \rightarrow A$, then each X_n is a co- H^{p_n} -space and the all pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- H^{p_n} -primitive.*
- (2) *If X_{n-1} is a co- $H^{p_{n-1}}$ -space and the pair of k' -invariants $(k'_{n-1}(A), k'_{n-1}(X)) : \tilde{p}_* \rightarrow p_{n-1}$ is co- $H^{p_{n-1}}$ -primitive, then X_n is a co- H^{p_n} -space.*

Proof. (1) Since X is a co- H^p -space for a map $p : X \rightarrow A$, there is a map $\theta : X \rightarrow A \vee X$ such that $j\theta \sim (p \times 1)\Delta$, where $j : A \vee X \rightarrow A \times X$ is the inclusion. Then $\{A_n \vee X_n, q'_n \vee q_n, i'_n \vee i_n\}$ is a homology decomposition for $A \vee X$. Then we have, by Toomer's result [12, Theorem 4], that there are families of maps $p_n : X_n \rightarrow A_n$ and $\theta_n : X_n \rightarrow A_n \vee X_n$ such that $i'_n p_n = p_{n+1} i_n$ and $q'_n p_n \sim p q_n$, and $(i'_n \vee i_n)\theta_n = \theta_{n+1} i_n$ and $(q'_n \vee q_n)\theta_n \sim \theta q_n$ for $n = 2, 3, \dots$ respectively, and $k'_n(A)\tilde{p}_* \sim p_n k'_n(X) : M(H_{n+1}(X), n) \rightarrow A_n$ and $(k'_n(A) \vee k'_n(X))\tilde{\theta}_* \sim \theta_n k'_n(X) : M(H_{n+1}(X), n) \rightarrow A_n \vee X_n$, where $k'_n(A) : M(H_{n+1}(A), n) \rightarrow A_n$, $k'_n(X) : M(H_{n+1}(X), n) \rightarrow X_n$ are k' -invariants of A and X respectively, $\tilde{p}_* : M(H_{n+1}(X), n) \rightarrow M(H_{n+1}(A), n)$ and $\tilde{\theta}_* : M(H_{n+1}(X), n) \rightarrow M(H_{n+1}(A \vee X), n) \approx M(H_{n+1}(A) \oplus H_{n+1}(X), n) \approx M(H_{n+1}(A), n) \vee M(H_{n+1}(X), n)$ are the induced maps by $p : X \rightarrow A$ and $\theta : X \rightarrow A \vee X$ respectively. It is known [12] that the homology decomposition of a rational space is well defined up to homotopy type. Thus we know that if $f \sim g : X \rightarrow A$, then $f_n \sim g_n : X_n \rightarrow A_n$. Since $p_1 j\theta \sim p$ and $p_2 j\theta \sim 1$, we know that $p_1 j_n \theta_n \sim p_n$ and $p_2 j_n \theta_n \sim 1$. Thus for each n , there exists a co- H^p -structure $\theta_n : X_n \rightarrow A_n \vee X_n$ such that $j_n \theta_n \sim (p_n \times 1)\Delta$, where $j_n : A_n \vee X_n \rightarrow A_n \times X_n$ is the inclusion and $p_n : X_n \rightarrow A_n$ is an induced map from $p : X \rightarrow A$, and X_n is a co- H^{p_n} -space. Moreover, since there is an extension $\theta_{n+1} : X_{n+1} \rightarrow A_{n+1} \vee X_{n+1}$ of θ_n such that $\theta_{n+1} i_n = (i'_n \vee i_n)\theta_n$, we know, from Lemma , that $(i'_n \vee i_n)\theta_n k'_n(X) \sim *$ and all the pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- H^{p_n} -primitive.

(2) It follows from Theorem 3.6 (1). □

Observe that X and A are homotopy types of the direct limits $\varinjlim X_n$ and $\varinjlim A_n$ respectively. Moreover, since each X_n a co- H^{p_n} -space and all pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- H^{p_n} -primitive, we see that X admit a co- H^p -structure. Thus we have the following corollary.

COROLLARY 3.8. *Let X and A be rational spaces and $p : X \rightarrow A$ a map, and $\{X_n, q_n, i_n\}$ and $\{A_n, q'_n, i'_n\}$ homology decompositions for X and A respectively. Then X is a co- H^p -space for a map $p : X \rightarrow A$ if and only if for each n , X_n is a co- H^{p_n} -space and the all pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- H^{p_n} -primitive.*

Taking $p = 1_{X_n}$, $p' = 1_{M(H_{n+1}(X), n)}$, $r = s = k'_n(X)$, we can obtain, from the fact that $i_n : X_n \rightarrow X_{n+1}$ is a co- H -map if and only if $(k'_n(X), k'_n(X)) : 1 \rightarrow 1_{X_n}$ is co- H -primitive and the above corollary, the following corollary given by Golasinski and Klein [3] for rational spaces.

COROLLARY 3.9. [3] *Let X be a rational space and $\{X_n, q_n, i_n\}$ a homology decomposition for X . Then X is n co- H -space if and only if for each X_n there exists such a co- H -structure that $i_n : X_n \rightarrow X_{n+1}$ is a co- H -map.*

THEOREM 3.10. *Let X and A be rational spaces and $p : X \rightarrow A$ a map, and $\{X_n, q_n, i_n\}$ and $\{A_n, q'_n, i'_n\}$ homology decompositions for X and A respectively.*

- (1) *If X is a co- T^p -space for a map $p : X \rightarrow A$, then each X_n is a co- T^{p_n} -space and the all pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- T^{p_n} -primitive.*
- (2) *If X_{n-1} is a co- $T^{p_{n-1}}$ -space and the pair of k' -invariants $(k'_{n-1}(A), k'_{n-1}(X)) : \tilde{p}_* \rightarrow p_{n-1}$ is co- $T^{p_{n-1}}$ -primitive, then X_n is a co- T^{p_n} -space.*

Proof. (1) Since X is a co- T^p -space for a map $p : X \rightarrow A$, there is a map $\theta : X \rightarrow A \vee \Omega\Sigma X$ such that $j\theta \sim (p \times e')\Delta$, where $j : A \vee \Omega\Sigma X \rightarrow A \times \Omega\Sigma X$ is the inclusion. Then $\{A_n \vee \Omega\Sigma X_n, q'_n \vee \Omega\Sigma q_n, i'_n \vee \Omega\Sigma i_n\}$ is a homology decomposition for $A \vee \Omega\Sigma X$. Then we have, by Toomer's result [15, Theorem 4], that there are families of maps $p_n : X_n \rightarrow A_n$ and $\theta_n : X_n \rightarrow A_n \vee \Omega\Sigma X_n$ such that $i'_n p_n = p_{n+1} i_n$ and $q'_n p_n \sim p q_n$, and $(i'_n \vee \Omega\Sigma i_n)\theta_n = \theta_{n+1} i_n$ and $(q'_n \vee \Omega\Sigma q_n)\theta_n \sim \theta q_n$ for $n = 2, 3, \dots$ respectively, and $k'_n(A)\tilde{p}_* \sim p_n k'_n(X) : M(H_{n+1}(X), n) \rightarrow A_n$ and $(k'_n(A) \vee$

$k'_n(\Omega\Sigma X)\tilde{\theta}_* \sim \theta_n k'_n(X) : M(H_{n+1}(X), n) \rightarrow A_n \vee \Omega\Sigma X_n$, where $k'_n(A) : M(H_{n+1}(A), n) \rightarrow A_n$, $k'_n(X) : M(H_{n+1}(X), n) \rightarrow X_n$ are k' -invariants of A and X respectively, $\tilde{p}_* : M(H_{n+1}(X), n) \rightarrow M(H_{n+1}(A), n)$ and $\tilde{\theta}_* : M(H_{n+1}(X), n) \rightarrow M(H_{n+1}(A \vee \Omega\Sigma X), n) \approx M(H_{n+1}(A) \oplus H_{n+1}(\Omega\Sigma X), n) \approx M(H_{n+1}(A), n) \vee M(H_{n+1}(\Omega\Sigma X), n)$ are the induced maps by $p : X \rightarrow A$ and $\theta : X \rightarrow A \vee \Omega\Sigma X$ respectively. It is known [12] that the homology decomposition of a rational space is well defined up to homotopy type. Thus we know that if $f \sim g : X \rightarrow A$, then $f_n \sim g_n : X_n \rightarrow A_n$. Since $p_1 j \theta \sim p$ and $p_2 j \theta \sim e'$, we know that $p_1 j_n \theta_n \sim p_n$ and $p_2 j_n \theta_n \sim e'_{X_n}$. Thus for each n , there exists a co- T^p -structure $\theta_n : X_n \rightarrow A_n \vee \Omega\Sigma X_n$ such that $j_n \theta_n \sim (p_n \times e'_{X_n}) \Delta$, where $j_n : A_n \vee \Omega\Sigma X_n \rightarrow A_n \times \Omega\Sigma X_n$ is the inclusion and $p_n : X_n \rightarrow A_n$ is an induced map from $p : X \rightarrow A$, and X_n is a co- T^{p_n} -space. Moreover, since there is an extension $\theta_{n+1} : X_{n+1} \rightarrow A_{n+1} \vee \Omega\Sigma X_{n+1}$ of θ_n such that $\theta_{n+1} i_n = (i'_n \vee \Omega\Sigma i_n) \theta_n$, we know, from Lemma , that $(i'_n \vee \Omega\Sigma i_n) \theta_n k'_n(X) \sim *$ and all the pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- T^{p_n} -primitive.

(2) It follows from Theorem 3.6(2). □

Observe that X and A are homotopy types of the direct limits $\varinjlim X_n$ and $\varinjlim A_n$ respectively. Moreover, since each X_n a co- T^{p_n} -space and all pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- T^{p_n} -primitive, we see that X admit a co- T^p -structure. Thus we have the following corollary.

COROLLARY 3.11. *Let X and A be rational spaces and $p : X \rightarrow A$ a map, and $\{X_n, q_n, i_n\}$ and $\{A_n, q'_n, i'_n\}$ homology decompositions for X and A respectively. Then X is a co- T^p -space for a map $p : X \rightarrow A$ if and only if for each n , X_n is a co- T^{p_n} -space and the all pair of k' invariants $(k'_n(A), k'_n(X)) : \tilde{p}_* \rightarrow p_n$ are co- T^{p_n} -primitive.*

References

- [1] J. Aguade, *Decomposable free loop spaces*, Canad. J. Math. **39** (1987), 938-955.
- [2] B. Eckmann and P. Hilton, *Decomposition homologique d'un polyedre simplement connexe*, ibid **248** (1959), 2054-2558.
- [3] M. Golasinski and J. R. Klein, *On maps into a co-H-spaces*, Hiroshima Math. J. **28** (1998), 321-327.
- [4] D. H. Gottlieb, *A certain subgroup of the fundamental group*, Amer. J. Math. **87** (1965), 840-856.

- [5] H. B. Haslam, *G-spaces and H-spaces*, Ph. D. Thesis, Univ. of California, Irvine, 1969.
- [6] P. Hilton, *Homotopy Theory and Duality*, Gordon and Breach Science Pub. 1965.
- [7] N. Iwase, *Ganea's conjecture on LS-category*, Bull. London Math. Soc. **30** (1998), no. 6, 623-634.
- [8] D. W. Kahn, *A note on H-spaces and Postnikov systems of spheres*, Proc. Amer. Math. Soc. **15** (1964), 300-307.
- [9] K. L. Lim, *Cocyclic maps and coevaluation subgroups*, Canad. Math. Bull. **30** (1987), 63-71.
- [10] N. Oda, *The homotopy of the axes of pairings*, Canad. J. Math. **17** (1990), 856-868.
- [11] G. H. Toomer, *Liusternik-Schnirelmann Category and the Moore spectral sequence*, Ph. D. Thesis, Cornell Univ. 1974.
- [12] G. H. Toomer, *Two Applications of Homology Decompositions*, Can. J. Math. **27** (1975), no. 2, 323-329.
- [13] K. Varadarajan, *Generalized Gottlieb groups*, J. Indian Math. Soc. **33** (1969), 141-164.
- [14] M. H. Woo and Y. S. Yoon, *T-spaces by the Gottlieb groups and duality*, J. Austral. Math. Soc. (Series A) **59** (1995), 193-203.
- [15] Y. S. Yoon, *Lifting Gottlieb sets and duality*, Proc. Amer. Math. Soc. **119** (1993), no. 4, 1315-1321.
- [16] Y. S. Yoon, *The generalized dual Gottlieb sets*, Topology Appl. **109** (2001), 173-181.
- [17] Y. S. Yoon, *H^f -spaces for maps and their duals*, J. Korea Soc. Math. Edu. Series B **14** (2007), no. 4, 287-306.
- [18] Y. S. Yoon, *Lifting T-structures and their duals*, J. Chungcheong Math. Soc. **20** (2007), no. 3, 245-259.
- [19] Y. S. Yoon, *G'_p -spaces for maps and Homology Decompositions*, J. Chungcheong Math. Soc. **28** (2015), no. 4, 603-614.
- [20] Y. S. Yoon, *Principal Fibrations and Generalized H-spaces*, J. Chungcheong Math. Soc. **29** (2016), no. 1, 177-186.

*

Department of Mathematics Education
Hannam University
Daejeon 306-791, Korea
E-mail: yoon@hannam.ac.kr