

SOME SHADOWING PROPERTIES OF THE SHIFTS ON THE INVERSE LIMIT SPACES

NYAMDAAA TSEG MID*

ABSTRACT. Let $f : X \rightarrow X$ be a continuous surjection of a compact metric space X and let $\sigma_f : X_f \rightarrow X_f$ be the shift map on the inverse limit space X_f constructed by f . We show that if a continuous surjective map f has some shadowing properties: the asymptotic average shadowing property, the average shadowing property, the two side limit shadowing property, then σ_f also has the same properties.

1. Introduction and preliminaries

Inverse limit is a useful tool to study the dynamical properties of smooth systems; some dynamical properties of can be interpreted by the topological structures of the inverse limit dynamical system; for example, Chen and Li [4] proved that (X, f) has the shadowing property if and only if $(\lim_{\leftarrow}(X, f), \sigma_f)$ has so. Li [8] proved that some dynamical properties hold simultaneously for both f and σ_f . Also A. Barzanouni [2] show that the relationship between ergodic shadowing property and inverse shadowing property for a surjective continuous map on a compact metric space and shift map on the inverse limit space. M. Lee [7] show that f has the asymptotic average shadowing, the average shadowing, the ergodic shadowing property then σ_f is topologically transitive.

In this paper, we discuss (X, f) has the two-sided limit shadowing property, the average shadowing property and the asymptotic average shadowing property then shift map on the inverse limit space has so.

Let X be a compact metric space with metric d and $X^{\mathbb{Z}}$ denote the product topological space $X^{\mathbb{Z}} = \{(x_i) : x_i \in X, i \in \mathbb{Z}\}$. Then $X^{\mathbb{Z}}$ is compact. We define a compatible metric \tilde{d} for $X^{\mathbb{Z}}$ by

$$\tilde{d}((x_i)(y_i)) = \sum_{i=-\infty}^{\infty} \frac{d(x_i, y_i)}{2^{|i|}}.$$

Received August 15, 2018; Accepted October 24, 2018.

2010 Mathematics Subject Classification: Primary 37C50, 54H20 .

Key words and phrases: inverse limit space, shift map, two-sided limit, average shadowing, asymptotic average shadowing property.

*The author was supported by the second stage of Brain Korea 21 project.

A homeomorphism $\sigma_f : X^{\mathbb{Z}} \rightarrow X^{\mathbb{Z}}$, which is defined by

$$\sigma_f((x_i)) = (y_i) \text{ and } y_i = x_{i+1} \text{ for all } i \in \mathbb{Z},$$

is called the shift map.

For $f : X \rightarrow X$ a continuous surjection, we let,

$$X_f = \{(x_i) : x_i \in X, \text{ and } f(x_i) = x_{i+1}, i \in \mathbb{Z}\}.$$

Then X_f is a closed subset of $X^{\mathbb{Z}}$. Moreover we have $\sigma_f((x_i)) = (f(x_i))$ for all $(x_i) \in X_f$, and so X_f is σ_f -invariant, i.e. $\sigma_f(X_f) = X_f$.

The space X_f is called the inverse limit space constructed by f . The restriction $\sigma_f = \sigma_f|_{X_f} : X_f \rightarrow X_f$ is called the shift map determined by f .

A sequence $(x_i)_{i \in \mathbb{Z}}$ of points X is a *two-sided limit pseudo orbit* for f if it satisfying

$$d(f(x_i), x_{i+1}) \rightarrow 0, \text{ as } |i| \rightarrow \infty.$$

A sequence $(x_i)_{i \in \mathbb{Z}}$ of points X is *two-sided limit shadowed* by $y \in X$ if it satisfying

$$d(f^i(y), x_i) \rightarrow 0, \text{ as } |i| \rightarrow \infty.$$

We say that f has the *two-sided limit shadowing property* if every two-sided limit pseudo-orbit is two-sided limit shadowed.

For $\delta > 0$, a sequence $(x_i)_{i \in \mathbb{Z}}$ of points in X is a δ -*average pseudo orbit* for f if there is an integer $N = N(\delta) > 0$ such that:

$$\frac{1}{n} \sum_{i=1}^n d(f(x_{i+k}), x_{i+k+1}) < \delta, \text{ for all } n \geq N, k \in \mathbb{Z}.$$

A sequence $(x_i)_{i \in \mathbb{Z}}$ of points in X is ε -*shadowed in average* by $y \in X$ if

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n d(f^i(y), x_i) < \varepsilon.$$

We say that f has the *average shadowing property* if for every $\varepsilon > 0$ there is $\delta > 0$ such that every δ -average pseudo orbit of f is ε -shadowed on average by some point in X .

A sequence $(x_i)_{i \in \mathbb{Z}}$ of points in X is a δ -*asymptotic average pseudo orbit* for f if there is an integer $N = N(\delta) > 0$ such that:

$$\frac{1}{n} \sum_{i=1}^n d(f(x_{i+k}), x_{i+k+1}) \rightarrow 0, \text{ for all } n \geq N, k \in \mathbb{Z}.$$

A sequence $(x_i)_{i \in \mathbb{Z}}$ of points X is ε -*asymptotically shadowed in average* by $y \in X$ if

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n d(f^i(y), x_i) \rightarrow 0.$$

We say that f has the *asymptotic average shadowing property* provided that every asymptotic average pseudo-orbit of f is asymptotically shadowed in average by some point in X . We say that a finite δ -pseudo orbit $\{x_i\}_{i=0}^k$ of f is

a δ -chain from x_0 to x_k with length $k + 1$. A non-empty subset A of X is said to be *chain transitive* if for any $x, y \in A$ and any $\delta > 0$ there is a δ -chain of f from x to y . A map f is said to be *chain transitive* if X is a chain transitive set.

THEOREM 1.1. [5, Theorem 3.1] *Let X be a compact metric space and f be a continuous map from X onto itself. If f has the asymptotic average shadowing property, then f is chain transitive.*

2. Main results

THEOREM 2.1. *Let $f : X \rightarrow X$ be a continuous surjective map on a compact metric space X .*

1. *If f has average shadowing property, then the shift map σ_f on the inverse limit space X_f has average shadowing property.*
2. *If f has asymptotic average shadowing property, then the shift map σ_f on the inverse limit space X_f has asymptotic average shadowing property.*
3. *If f has two-side limit shadowing property, then the shift map σ_f on the inverse limit space X_f has two-side limit shadowing property.*

Proof. Proof of 1. Let $\varepsilon > 0$ and $D = \text{diam } X$. Choose $N > 0$ with $D/2^{N-2} < \varepsilon$, and let $\gamma > 0$ be a number such that

$$d(x, y) \leq \gamma \Rightarrow \max_{0 \leq i \leq 2N} d(f^i(x), f^i(y)) \leq \frac{\varepsilon}{8}.$$

By average shadowing property of f there is $\delta_1 > 0$ such that any δ_1 -average pseudo orbit of f is γ -average shadowed. Choose $\delta_2 > 0$ with $0 < 2^N \delta_2 < \delta_1$. Define a sequence $\{w_i^n\}_{n,i \in \mathbb{Z}}$ is a δ_2 -average pseudo orbit of σ_f in X_f such that $\{w_i^n\}_{n \in \mathbb{Z}}$ is a periodic δ_1 -average pseudo orbit of f for each $i \in \mathbb{Z}$.

Then we have,

$$\begin{aligned} \delta_2 > \frac{1}{n} \sum_{i=0}^{n-1} \tilde{d}(\sigma_f(w_j^{k+i}), w_j^{k+i+1}) &= \frac{1}{n} \sum_{i=0}^{n-1} \left(\sum_{j=-\infty}^{\infty} \frac{d(f(w_j^{k+i}), w_j^{k+i+1})}{2^{|j|}} \right) \\ &\geq \frac{1}{n} \sum_{i=0}^{n-1} \frac{d(f(w_{-N}^{k+i}), w_{-N}^{k+i+1})}{2^{|N|}}, \quad n \geq N. \end{aligned}$$

Since $\{w_{-N}^n\}_{n \in \mathbb{Z}}$ is a periodic δ_1 -average pseudo orbit of f , $\{w_i^n\}_{n,i \in \mathbb{Z}}$ δ_2 -average pseudo orbit of σ_f . Also we can find $z \in X$, such that $n \geq N$

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} d(f^i(z), w_{-N}^i) < \gamma.$$

We put $z_{i-N} = f^i(z)$ for $i \geq 0$ and take $z_{i-N} \in f^{-1}(z_{i+1-N})$ for $i < 0$. Then $\tilde{z} = \{z_i\}_{i \in \mathbb{Z}} \in X_f$ and

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \tilde{d}(\sigma_f^i(z_j), w_j^i) &= \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left(\sum_{j=-\infty}^{\infty} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} \right) \\ &= \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left(\sum_{j=-N}^N \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} + \sum_{j=-\infty}^{-N-1} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} + \sum_{j=N+1}^{\infty} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} \right) \\ &\leq \frac{3\varepsilon}{8} + \frac{\varepsilon}{8} + \frac{\varepsilon}{8} < \varepsilon. \end{aligned}$$

Hence $\{w_i^n\}_{n,i \in \mathbb{Z}}$ is a δ_2 -average pseudo orbit of σ_f is ε -shadowed in average by $\tilde{z} \in X_f$.

Proof of 2. Let $D = \text{diam}(X)$ and $\varepsilon > 0$. Choose $N > 0$ with $D/2^{N-2} < \varepsilon$, and let $\gamma > 0$ be a number such that

$$d(x, y) < \gamma \rightarrow 0 \quad \Rightarrow \quad \max_{0 \leq i \leq 2N} d(f^i(x), f^i(y)) \leq \frac{\varepsilon}{8} \rightarrow 0.$$

Since asymptotic average shadowing property of f , there is $\delta_1 > 0$ such that any δ_1 -asymptotic average pseudo orbit of f is γ -asymptotically shadowed. Choose $\delta_2 > 0$ with $0 < 2^N \delta_2 < \delta_1$. Define a sequence $\{w_i^n\}_{n,i \in \mathbb{Z}}$ is a δ_2 -asymptotic average pseudo orbit of σ_f in X_f such that $\{w_i^n\}_{n \in \mathbb{Z}}$ is δ_1 -asymptotic average pseudo orbit of f for all $i \in \mathbb{Z}$. Then

$$\begin{aligned} \delta_2 > \frac{1}{n} \sum_{i=0}^{n-1} \tilde{d}(\sigma_f(w_j^{k+i}), w_j^{k+i+1}) &= \frac{1}{n} \sum_{i=0}^{n-1} \left(\sum_{j=-\infty}^{\infty} \frac{d(f(w_j^{k+i}), w_j^{k+i+1})}{2^{|j|}} \right) \\ &\geq \frac{1}{n} \sum_{i=0}^{n-1} \frac{d(f(w_{-N}^{k+i}), w_{-N}^{k+i+1})}{2^{|N|}}. \\ \Rightarrow \delta_1 > 2^{|N|} \delta_2 &\geq \frac{1}{n} \sum_{i=0}^{n-1} d(f(w_{-N}^{k+i}), w_{-N}^{k+i+1}) \rightarrow 0. \end{aligned}$$

Since $\{w_{-N}^n\}_{n \in \mathbb{Z}}$ is a δ_1 -asymptotic average pseudo orbit of f , $\{w_i^n\}_{n,i \in \mathbb{Z}}$ is a δ_2 -asymptotic average pseudo orbit of σ_f . Thus we can find $z \in X$, such that $n \geq N$

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} d(f^i(z), w_{-N}^i) \rightarrow 0.$$

We put $z_{i-N} = f^i(z)$ for $i \geq 0$ and take $z_{i-N} \in f^{-1}(z_{i+1-N})$ for $i < 0$. Then $\tilde{z} = \{z_i\}_{i \in \mathbb{Z}} \in X_f$ and

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \tilde{d}(\sigma_f^i(z_j), w_j^i) &= \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left(\sum_{j=-\infty}^{\infty} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} \right) \\ &= \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left(\sum_{j=-N}^N \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} + \sum_{j=-\infty}^{-N-1} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} + \sum_{j=N+1}^{\infty} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} \right) \\ &\leq \frac{3\varepsilon}{8} + \frac{\varepsilon}{8} + \frac{\varepsilon}{8} \rightarrow 0. \end{aligned}$$

Hence $\{w_i^n\}_{n,i \in \mathbb{Z}}$ is a δ_2 -asymptotic average pseudo orbit of σ_f is ε - asymptotically shadowed in average by $\tilde{z} \in X_f$.

Proof of 3. Let $D = \text{diam}(X)$ and $\varepsilon > 0$. let $|k| > N$ and $\gamma > 0$ be a number such that

$$d(x, y) < \gamma = \frac{D}{2^{|k|}} \Rightarrow \max_{0 \leq i \leq 2N} d(f^i(x), f^i(y)) \leq \frac{\varepsilon}{8} \rightarrow 0.$$

Define a sequence $\{w_i^n\}_{n,i \in \mathbb{Z}}$ is a two-sided limit pseudo-orbit of σ_f in X_f such that $\{w_i^n\}_{n \in \mathbb{Z}}$ is a two-sided limit pseudo-orbit of f for all $i \in \mathbb{Z}$. Then we have

$$\tilde{d}(\sigma_f(x_i^n, x_i^{n+1})) \geq \frac{d(f(x_{-N}^n), x_{-N}^{n+1})}{2^N}.$$

Since $\{x_{-N}^n\}_{n \in \mathbb{Z}}$ is a two-sided limit pseudo orbit of f , $\{w_i^n\}_{n \in \mathbb{Z}}$ is a two-sided limit pseudo-orbit of σ_f . Then we can find $z \in X$ such that $d(f^n(z), x_{-N}^n) = 0$ when $|n| \rightarrow \infty$.

We put $z_{i-N} = f^i(z)$ for $i \geq 0$ and take $z_{i-N} \in f^{-1}(z_{i+1-N})$ for $i < 0$. Then $\tilde{z} = \{z_i\}_{i \in \mathbb{Z}} \in X_f$ and

$$\begin{aligned} \tilde{d}(\sigma_f^i(z_j), w_j^i) &= \sum_{j=-\infty}^{\infty} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} \\ &= \sum_{j=-N}^N \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} + \sum_{j=-\infty}^{-N-1} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} + \sum_{j=N+1}^{\infty} \frac{d(f^i(z_j), w_j^i)}{2^{|j|}} \\ &\leq \frac{3\varepsilon}{8} + \frac{\varepsilon}{8} + \frac{\varepsilon}{8} \rightarrow 0. \end{aligned}$$

Hence $\{w_i^n\}_{n,i \in \mathbb{Z}}$ two-sided limit pseudo-orbit of σ_f is two-sided limit shadowed by $\tilde{z} \in X_f$. □

COROLLARY 2.2. *If a continuous map f has asymptotic average shadowing property, then the shift map σ_f on the inverse limit space X_f is chain transitive.*

Proof. By 2. of Theorem 2.1, f has asymptotic average shadowing property, then σ_f has asymptotic average shadowing property, so by Theorem 1.1, σ_f is chain transitive. □

COROLLARY 2.3. [7, Remark 2.3] *If surjective continuous map f has average shadowing property, then the shift map σ_f on the inverse limit space X_f is chain transitive.*

COROLLARY 2.4. *If homeomorphism f of a compact metric space has two-sided limit shadowing property, then the shift map σ_f on the inverse limit space X_f has average shadowing and asymptotic average shadowing property. Moreover, σ_f is chain transitive.*

Proof. By [3, Theorem B], f has two-sided limit shadowing property, then f has average shadowing and asymptotic average shadowing property, so by Theorem 2.1, σ_f has average shadowing and asymptotic average shadowing property. Then by Corollary 2.2 and Corollary 2.3, σ_f is chain transitive. □

References

- [1] N. Aoki and K. Hiriade, *Topological Theory of Dynamical Systems*, Recent Advances, North-Holland, 1994.
- [2] A. Barzanouni, *Inverse Limit Spaces with Various Shadowing Property*, J. Math. **2014** (2014), 4pages. <http://dx.doi.org/10.1155/2014/169183>
- [3] B. Carvalho and D. Kwietniak, *On homeomorphisms with the two-sided limit shadowing property*, J. Math. Anal. Appl. **420** (2014), no. 1, 801-813
- [4] L. Chen and S. Li, *Shadowing Property for Inverse Limit Spaces*, Proc. Amer. Math. Soc. **115** (1992), no. 1, 573-580.
- [5] R. Gu, *The asymptotic average shadowing property and transitivity*, Nonlinear Anal. **67** (2007), no. 6, 1680-1689
- [6] J. Park and Y. Zhang, *Average shadowing properties on compact metric spaces*, Commun. Korean Math. Soc. **21** (2006), no. 2, 355-361.
- [7] M. Lee, *Various shadowing properties for inverse limit systems*, J. Chungcheong. Math. Soc. **29** (2016), no. 4, 657-661
- [8] S. Li, *Dynamical properties of the shift maps on the inverse limit spaces*, Ergodic Theory and Dynam. Systems, **12** (1992), no. 1, 95-108.
- [9] E. Shah and T. K. Das, *On shadowing property for inverse limit spaces*, Math. Slovaca, **58** (2008),no. 1, 95-100.
- [10] Y. Zhang, *On the average-shadowing property*, Acta Sci. Natur. Univ. Pekinensis, **37** (2001), no. 5, 648-651.

*

Department of Mathematics
Chungnam National University
Daejeon 34134, Republic of Korea
E-mail: ts.nyamdavaa@yahoo.com