

CLOSED AND DENSE ELEMENTS OF BE -ALGEBRAS

M.BALA PRABHAKAR*, S.KALESHA VALI**, AND M. SAMBASIVA
RAO.***

ABSTRACT. The notions of closed elements and dense elements are introduced in BE -algebras. Characterization theorems of closed elements and closed filters are obtained. The notion of dense elements is introduced in BE -algebras. Dense BE -algebras are characterized with the help of maximal filters and congruences. The concept of D -filters is introduced in BE -algebras. A set of equivalent conditions is derived for every D -filter to become a closed filter.

1. Introduction

The notion of BE -algebras was introduced and extensively studied by H.S. Kim and Y.H. Kim in [5]. These classes of BE -algebras were introduced as a generalization of the class of BCK -algebras by K. Iseki and S. Tanaka [4]. Some properties of filters of BE -algebras were studied by S.S. Ahn and Y.H. Kim in [1] and by J.L. Meng in [6]. In [10], A. Walendziak discussed some relationships between congruence relations and normal filters of a BE -algebra. In [3], Gispert and Torrens defined the Boolean center and the Boolean skeleton of a bounded BCK -algebra and they used the Boolean skeleton to obtain a representation of bounded BCK -algebras. In [7], C. Muresan studied some properties of dense elements and the radical of residuated lattices. Later in 2011, D. Piciu and D. Tascu [8] developed a theory of localization for bounded commutative BCK -algebras.

In this paper, the notions of closed elements is introduced in BE -algebras. A set of equivalent conditions is derived for every element of a BE -algebra to become closed. The notion of closed filters is introduced in BE -algebras. Closed filters are characterized in terms of closed elements

Received Oct 22, 2018; Accepted Jan 10, 2019.

2010 Mathematics Subject Classification: 06F35, 03G25, 08A30.

Key words and phrases: BE -algebra; maximal filter; closed element; closed filter; dense element; dense BE -algebra; D -filter.

*** The corresponding author.

of BE -algebras. The notions of dense elements and dense BE -algebras are introduced. Some characterization theorems of dense BE -algebras are derived in terms of maximal filters and congruences. The concept of D -filters is introduced in BE -algebras. A set of equivalent conditions is obtained for every D -filter of a BE -algebra to become a closed filter.

2. Preliminaries

In this section, we present certain definitions and results which are taken mostly from the papers [1], [2], [3] and [5] for the ready reference of the reader.

DEFINITION 2.1. [5] An algebra $(X, *, 1)$ of type $(2, 0)$ is called a BE -algebra if it satisfies the following properties:

- (1) $x * x = 1$,
- (2) $x * 1 = 1$,
- (3) $1 * x = x$,
- (4) $x * (y * z) = y * (x * z)$ for all $x, y, z \in X$.

A BE -algebra X is called self-distributive if $x * (y * z) = (x * y) * (x * z)$ for all $x, y, z \in X$. A BE -algebra X is called transitive if $y * z \leq (x * y) * (x * z)$ for all $x, y, z \in X$. Every self-distributive BE -algebra is transitive. A BE -algebra X is called implicative if $(x * y) * x = x$ for all $x, y \in X$. A BE -algebra X is called commutative if $(x * y) * y = (y * x) * x$ for all $x, y \in X$. We introduce a relation \leq on a BE -algebra X by $x \leq y$ if and only if $x * y = 1$ for all $x, y \in X$. Clearly \leq is reflexive and symmetric. If X is commutative, then \leq is anti-symmetric and hence a partial order on X . Throughout this article, X stands for a partially ordered set.

THEOREM 2.2. [6] Let X be a transitive BE -algebra and $x, y, z \in X$. Then

- (1) $1 \leq x$ implies $x = 1$,
- (2) $y \leq z$ implies $x * y \leq x * z$ and $z * x \leq y * x$.

DEFINITION 2.3. [1] A non-empty subset F of a BE -algebra X is called a filter of X if, for all $x, y \in X$, it satisfies the following properties:

- (1) $1 \in F$,
- (2) $x \in F$ and $x * y \in F$ imply that $y \in F$.

For any $a \in X$, $\langle a \rangle = \{x \in X \mid a^n * x = 1 \text{ for some } n \in \mathbb{N}\}$ is called the principal filter generated a . If X is self-distributive, then $\langle a \rangle = \{x \in X \mid a * x = 1\}$. Let $(X, *, 0, 1)$ be a bounded BE -algebra, where 0 is the smallest element of X with respect to the ordering \leq . Hence $0 * x = 1$ for all $x \in X$. For any $x \in X$, define a unary operation N on X as $xN = x * 0$, where xN is called the pseudo-complement of x . It is easily seen that $0N = 1$ and $1N = 0$.

PROPOSITION 2.4. [2] *Let X be a transitive BE -algebra and $x, y \in X$. Then the following properties hold.*

- (1) $x \leq xNN$,
- (2) $x \leq y$ implies $yN \leq xN$,
- (3) $xNNN = xN$,
- (4) $x * yN = xNN * yN$,
- (5) $(x * yNN)NN = x * yNN$,
- (6) $(x * y)NN \leq xNN * yNN$.

THEOREM 2.5. [3] *Let X be a BE -algebra and $a, b \in X$. Then $a * c = 1$ and $b * c = 1$ imply $c = 1$ for all $c \in X$ if and only if $\langle a \rangle \cap \langle b \rangle = \{1\}$.*

An element x of X is called Boolean [3] if $\langle x \rangle \cap \langle xN \rangle = \{1\}$. Let us denote the set of all Boolean elements of a bounded BE -algebra X by $\mathcal{B}(X)$. Clearly $0, 1 \in \mathcal{B}(X)$.

PROPOSITION 2.6. [3] *Let X be a transitive BE -algebra. Then for every $a \in \mathcal{B}(X)$ and $x, y \in X$, the following conditions hold.*

- (1) $aNN = a$,
- (2) $a * (a * x) = a * x$,
- (3) $a * (x * y) = (a * x) * (a * y)$.

3. Closed elements of BE -algebras

In this section, the notion of closed elements is introduced and studied their properties. A set of equivalent conditions is established for every element of a BE -algebra to become a Boolean element. A set of equivalent conditions is derived for every element of a BE -algebra to become a closed element.

DEFINITION 3.1. An element a of a BE -algebra is a *closed element* if $aNN = a$.

We denote by $\mathcal{C}(X)$ the set of all closed elements of a BE -algebra X . Obviously $0, 1 \in \mathcal{C}(X)$. In the following, a characterization of closed elements is derived.

THEOREM 3.2. *The following are equivalent in a transitive BE -algebra X :*

- (1) *Every element is closed;*
- (2) *for $x, y \in X$, $xN = yN$ implies $x = y$;*
- (3) *for $x, y \in X$, $xN * yN = y * x$.*

Proof. (1) \Leftrightarrow (2): Assume that every element of X is closed. Let $x, y \in X$ be such that $xN = yN$. Hence, it implies that $x = xNN = yNN = y$. Conversely, assume the condition (2). Let $x \in X$. By Proposition 2.4 (3), we have $xNNN = xN$. From the condition (2), we get $xNN = x$. Hence x is closed.

(2) \Rightarrow (3): Assume that the condition (2) holds. Let $x, y \in X$ be two arbitrary elements. Hence $xN * yN = (x * 0) * (y * 0) = y * ((x * 0) * 0) = y * xNN = y * x$.

(3) \Rightarrow (1): Assume that the condition (3) holds. Let $a \in X$. Then, we get $aNN = aN * 0 = aN * 1N = 1 * a = a$. Therefore a is closed. \square

It is observed from Proposition 2.6 (1) that every Boolean element of a BE -algebra is a closed element. It is evident from the following example that every closed element of a BE -algebra need not be Boolean.

EXAMPLE 3.3. *Let $X = \{1, a, b, c, d, 0\}$ be a non-empty set. Define a binary operation $*$ on X as follows:*

$*$	1	a	b	c	d	0
1	1	a	b	c	d	0
a	1	1	a	c	c	d
b	1	1	1	c	c	c
c	1	a	b	1	a	b
d	1	1	a	1	1	a
0	1	1	1	1	1	1

Clearly $(X, *, 0, 1)$ is a bounded BE -algebra with smallest element 0. Observe that $aNN = dN = a; bNN = cN = b; cNN = bN = c$ and $dNN = aN = d$. Therefore $\mathcal{C}(X) = X$. But the elements a and d of the BE -algebra X are not Boolean, because of $\langle a \rangle \cap \langle aN \rangle = \langle a \rangle \cap \langle d \rangle = \{1, a\} \cap \{1, d, a, c\} = \{1, a\} \neq \{1\}$ also $\langle d \rangle \cap \langle dN \rangle \neq \{1\}$.

THEOREM 3.4. *In an implicative BE -algebra, every closed element is Boolean.*

Proof. Let X be an implicative BE -algebra. Let $a \in \mathcal{C}(X)$. Let $x \in X$ and suppose $a * x = 1$ and $aN * x = 1$. Then, it infers that $xN \leq aNN = a \leq x$. Since X is implicative, we get $x = (x * 0) * x = xN * x = 1$. Hence by Theorem 2.5, we get that $\langle a \rangle \cap \langle aN \rangle = \{1\}$. Therefore a is a Boolean element. \square

REMARK 3.5. For any transitive BE -algebra X , define the set $X^* = \{x \in X \mid x = aN \text{ for some } a \in X\}$. Since $xNNN = xN$ for all $x \in X$, it is clear that $X^* = \mathcal{C}(X)$.

THEOREM 3.6. *Let X be a transitive BE -algebra. Then $\mathcal{C}(X)$ is closed under $*$.*

Proof. From Proposition 2.4 (5), it is clear. \square

For any two BE -algebras $(X_1, *, 0, 1)$ and $(X_2, *, 0', 1')$, it is clear that their product $X_1 \times X_2$ is a BE -algebra in which the pseudo-complement of any $(a, b) \in X_1 \times X_2$ is defined as (aN, bN) . Then the following result is an easy consequence.

THEOREM 3.7. *Let $(X_1, *, 0, 1)$ and $(X_2, *, 0, 1)$ be two bounded BE -algebras. Then a_1 and a_2 are closed elements of X_1 and X_2 respectively if and only if (a_1, a_2) is a closed element of $X_1 \times X_2$.*

Proof. Let $a_1 \in X_1$ and $a_2 \in X_2$. Assume that a_1 and a_2 are closed elements of X_1 and X_2 respectively. Clearly (a_1, a_2) is a closed element of $X_1 \times X_2$. Conversely assume that (a_1, a_2) is a closed element of $X_1 \times X_2$. Consider the projections $\Pi_i : X_1 \times X_2 \rightarrow X_i$ for $i = 1, 2$. Let $\Pi_i(a_1, a_2) = a_i$ for $i = 1, 2$. We now prove that a_1 and a_2 are closed elements of X_1 and X_2 respectively. Since (a_1, a_2) is a closed element of $X_1 \times X_2$, we get the following:

$$\begin{aligned} a_1NN &= \Pi_1(a_1NN, 1) = \Pi_1(a_1NN, 1NN) \\ &= \Pi_1(a_1, 1)NN = \Pi_1(a_1, 1) = a_1. \end{aligned}$$

Therefore a_1 is a closed element of X_1 . Similarly a_2 is a closed element of X_2 . \square

The following corollary is a direct consequence of the above theorem.

COROLLARY 3.8. *For any two BE -algebras X_1 and X_2 , $\mathcal{C}(X_1 \times X_2) = \mathcal{C}(X_1) \times \mathcal{C}(X_2)$.*

THEOREM 3.9. *Let $(X, *, 0, 1)$ and $(Y, *, 0', 1')$ be two BE -algebras and $\alpha : X \rightarrow Y$ a BE -morphism. If a is a closed element of X , then $\alpha(a)$ is a closed element of Y .*

Proof. Let $a \in \mathcal{C}(X)$. Then $\alpha(a)NN = \alpha(aNN) = \alpha(a)$. Therefore $\alpha(a) \in \mathcal{C}(Y)$. \square

In the following, the notion of *closed filters* of BE -algebras is introduced.

DEFINITION 3.10. A filter F of a BE -algebra X is called a closed filter if $xNN \in F$ implies $x \in F$ for any $x \in X$.

If every element of a BE -algebra is closed, then clearly every filter is a closed filter. However, in the following, closed filters of BE -algebras are characterized.

THEOREM 3.11. A filter F of a transitive BE -algebra X is closed if and only if for any $x, y \in X$, $xN = yN$ and $x \in F$ imply $y \in F$.

Proof. Assume that F is a closed filter of X . Let $x, y \in X$ be such that $xN = yN$ and $x \in F$. Since $x \in F$ and $x \leq xNN$, we get $yNN = xNN \in F$. Since F is closed, it yields that $y \in F$. Conversely, assume that the condition holds. Let $xNN \in F$ for $x \in X$. Since $xNNN = xN$, we get $x \in F$. Therefore F is a closed filter of X . \square

PROPOSITION 3.12. Every maximal filter of a transitive BE -algebra is closed.

Proof. Let F be a maximal filter of a transitive BE -algebra X . Let $x, y \in X$ be such that $xN = yN$ and $x \in F$. Suppose $y \notin F$. Then $\langle F \cup \{y\} \rangle = X$. Hence $0 \in \langle F \cup \{y\} \rangle$, which implies that $y^n * 0 \in F$ for some positive integer n . Hence

$$\begin{aligned}
y^n * 0 \in F &\Rightarrow \underbrace{y * (y * (\dots (y * 0) \dots))}_{n \text{ times}} \in F \\
&\Rightarrow \underbrace{y * (y * (\dots (y * (y * 0)) \dots))}_{n-1 \text{ times}} \in F \\
&\Rightarrow \underbrace{y * (y * (\dots (y * (x * 0)) \dots))}_{n-1 \text{ times}} \in F \\
&\Rightarrow x * \underbrace{(y * (y * (\dots (y * 0)) \dots))}_{n-1 \text{ times}} \in F \\
&\dots \\
&\dots \\
&\Rightarrow x^n * 0 \in F.
\end{aligned}$$

Since $x \in F$, we get that $0 \in F$, which is a contradiction. Hence, it infers $y \in F$. Therefore F is closed. \square

For any filter F of a self-distributive BE -algebra X , it was observed in [9] that θ_F defined by $(x, y) \in \theta_F \Leftrightarrow x * y \in F$ and $y * x \in F$ is the unique congruence whose kernel is F . If X is bounded, then the quotient algebra $X/F = \{F_x \mid x \in X\}$ (where F_x is the congruence class of x modulo θ_F) is also a bounded BE -algebra with smallest element F_0 in which $F_x * F_y = F_{x*y}$ and $F_x N = F_{xN}$ for all $x, y \in X$.

THEOREM 3.13. *The following are equivalent in a self-distributive BE -algebra X :*

- (1) *Every element of X is closed;*
- (2) *for any filter F , $\mathcal{C}(X/F) = X/F$;*
- (3) *every filter is closed.*

Proof. (1) \Rightarrow (2): It is obvious.

(2) \Rightarrow (3): Assume that the condition (2) holds. Let F be a filter of X and $x \in X$. Suppose $xNN \in F$. For this $x \in X$, we get $F_x \in X/F$. By the condition (2), we get $F_{xNN} = F_x NN = F_x$. Hence $(xNN, x) \in \theta_F$, which gives $xNN * x \in F$. Since $xNN \in F$, it yields that $x \in F$. Therefore F is closed.

(3) \Rightarrow (1): Assume that every filter is closed. Let $x \in X$. By (3), we get $\langle xNN \rangle$ is closed and $xNN \in \langle xNN \rangle$. Hence $x \in \langle xNN \rangle$. Thus $xNN \leq x$. Therefore $xNN = x$. \square

In the following theorem, properties of the homomorphic images and inverse images of closed filters of BE -algebras are studied.

THEOREM 3.14. *Let X and Y be two BE -algebras and $\psi : X \rightarrow Y$ a bounded BE -morphism. If F is a closed filter of Y , then $\psi^{-1}(F)$ is a closed filter of X . Moreover, if ψ is onto, then $\psi(F)$ is a closed filter for any closed filter F of X .*

Proof. Let F be a closed filter of Y . Clearly $\psi^{-1}(F)$ is a filter of X . Let $xNN \in \psi^{-1}(F)$. Then $\psi(x)NN = \psi(xNN) \in F$. Since F is a closed filter of Y , we get that $\psi(x) \in F$. Hence $x \in \psi^{-1}(F)$. Therefore $\psi^{-1}(F)$ is a closed filter of X . Suppose ψ is onto. Let F be a closed filter of X . Clearly $\psi(F)$ is a filter of Y . Let $xNN \in \psi(F)$ where $x \in Y$. Since ψ is onto, there exists $a \in X$ such that $\psi(a) = x$. Hence $\psi(aNN) = \psi(a)NN = xNN$. Hence $aNN \in F$. Since F is closed, it yields $a \in F$. Hence $x = \psi(a) \in \psi(F)$. Therefore $\psi(F)$ is a closed filter of Y . \square

In the following theorem, a set of equivalent conditions is derived for every filter of a BE -algebra to become a closed filter, which leads to a characterization of $\mathcal{C}(X)$.

THEOREM 3.15. *Let X be a self-distributive BE -algebra. Then the following conditions are equivalent.*

- (1) $\mathcal{C}(X) = X$;
- (2) every element is closed;
- (3) every filter is closed;
- (4) every principal filter is closed.

Proof. (1) \Leftrightarrow (2), (2) \Rightarrow (3) and (3) \Rightarrow (4) are obvious.

(4) \Rightarrow (2): Assume that every principal filter is closed. Let $x \in X$. Clearly $xNN \in \langle xNN \rangle$. Since $\langle xNN \rangle$ is a closed filter, it yields that $x \in \langle xNN \rangle$. Hence $xNN \leq x$. Since $x \leq xNN$, we get $x = xNN$. Therefore every element of X is closed. \square

4. Dense elements of BE -algebras

In this section, the notion of dense elements is introduced in BE -algebras. Some properties of the class of dense elements are studied. The concept of dense BE -algebras is introduced and characterized. The notion of D -filters is introduced and characterized with the help of closed elements.

DEFINITION 4.1. An element x of a BE -algebra X is called *dense* if $xN = 0$.

It is obvious that 1 is a dense element of X . Let us denote the class of all dense elements of a BE -algebra X by $\mathcal{D}(X)$. Then clearly $\mathcal{D}(X)$ is a subalgebra of X .

EXAMPLE 4.2. Let $X = \{1, a, b, 0\}$ be a set and $*$ a binary operation defined on X as follows:

$*$	1	a	b	0
1	1	a	b	0
a	1	1	1	a
b	1	a	1	0
0	1	1	1	1

Clearly $(X, *, 0, 1)$ is a bounded BE -algebra with smallest element 0. Observe that $aN = a$; $bN = 0$. Therefore b and 1 are dense but a is not a dense element.

PROPOSITION 4.3. For any transitive BE -algebra X , $\mathcal{D}(X)$ is a closed filter of X .

Proof. Clearly $1 \in \mathcal{D}(X)$. Let $x, x * y \in \mathcal{D}(X)$. Then $xN = 0$ and $(x * y)N = 0$. By Proposition 2.4 (6), we get $1 = 0N = (x * y)NN \leq xNN * yNN = 0N * yNN = yNN$. Thus $yN = 0$, which yields $y \in \mathcal{D}(X)$. Therefore $\mathcal{D}(X)$ is a filter of X . Let $xNN \in \mathcal{D}(X)$. Then $xN = xNNN = 0$, which yields $x \in \mathcal{D}(X)$. Therefore $\mathcal{D}(X)$ is closed. \square

THEOREM 4.4. *The following conditions hold in a BE -algebra X :*

- (1) $a \in \mathcal{D}(X)$ implies $aN \in \mathcal{B}(X)$,
- (2) $a \in \mathcal{D}(X)$ if and only if $aNN \in \mathcal{D}(X)$,
- (3) $\mathcal{D}(X)$ is a subalgebra of X ,
- (4) if a is a dense element in X , then $f(a)$ is a dense element in Y where $f : X \rightarrow Y$ is a bounded BE -morphism.

Proof. (1). Let $a \in \mathcal{D}(X)$. Then $\langle aN \rangle \cap \langle aNN \rangle = \langle 0 \rangle \cap \langle 1 \rangle = \{1\}$. Therefore $aN \in \mathcal{B}(X)$.

(2). Since $aNNN = aN$, it follows immediately.

(3). Let $a, b \in \mathcal{D}(X)$. Then from (2), it immediately infers $a * b \in \mathcal{D}(X)$.

(4). Let $a \in \mathcal{D}(X)$. Then $f(a)N = f(aN) = f(0) = 0$. Therefore $f(a) \in \mathcal{D}(Y)$. \square

In the following, the notion of dense BE -algebras is introduced.

DEFINITION 4.5. A BE -algebra X is called a *dense BE -algebra* if every non-zero element of X is dense (i.e. $xN = 0$ for all $0 \neq x \in X$).

Clearly the two-element bounded BE -algebra $\{0, 1\}$ is a dense BE -algebra. The bounded BE -algebra given in Example 4.2 is not a dense BE -algebra.

EXAMPLE 4.6. Let $X = \{1, a, b, c, 0\}$ be a non-empty set. Define a binary operation $*$ on X as follows:

$*$	1	a	b	c	0
1	1	a	b	c	0
a	1	1	b	b	0
b	1	a	1	a	0
c	1	1	1	1	0
0	1	1	1	1	1

Clearly $(X, *, 0, 1)$ is a bounded BE -algebra with smallest element 0. Observe that $aN = bN = cN = 0$. Hence $\mathcal{D}(X) = X - \{0\}$. Therefore X is a dense BE -algebra.

It is observed that the set $\mathcal{D}(X)$ is a filter in a transitive BE -algebra but it is not sure that $\mathcal{D}(X)$ is a filter in a BE -algebra. However, in the following theorem, a necessary and sufficient condition is derived for $\mathcal{D}(X)$ to become a maximal filter which leads to a characterization of dense BE -algebras.

THEOREM 4.7. *A BE -algebra X is dense if and only if $\mathcal{D}(X)$ is a maximal filter.*

Proof. Assume that X is dense. Clearly $1 \in \mathcal{D}(X)$. Let $a, a * b \in \mathcal{D}(X)$. Suppose $b = 0$. Since $aN = 0$, we get $0 = (a * b)N = (a * 0)N = 0N = 1$. which is a contradiction. Therefore $b \neq 0$. Since X is dense, it yields $bN = 0$. Hence $b \in \mathcal{D}(X)$. Therefore $\mathcal{D}(X)$ is a filter of X . Suppose P is a proper filter of X such that $\mathcal{D}(X) \subset P$. Choose $x \in P - \mathcal{D}(X)$. Clearly $x \neq 0$. Since X is dense, it gives $xN = 0$. Hence $x \in \mathcal{D}(X)$, which is contradiction. Therefore $\mathcal{D}(X)$ is a maximal filter of X .

Conversely, assume that $\mathcal{D}(X)$ is a maximal filter of X . Suppose X is non-dense. Then there exists $0 \neq x \in X$ such that $xN \neq 0$. Hence $x \notin \mathcal{D}(X)$. Then $\mathcal{D}(X) \subset \langle \mathcal{D}(X) \cup \{x\} \rangle$. Since $\mathcal{D}(X)$ is maximal, we get $\langle \mathcal{D}(X) \cup \{x\} \rangle = X$. Hence $x * (d * 0) = d * (x * 0) = 1$ for some $d \in \mathcal{D}(X)$. Hence $xN = 1$. Thus $x = 0$, which is a contradiction. Therefore X is a dense BE -algebra. \square

THEOREM 4.8. *Let X be a transitive BE -algebra. Then $\mathcal{D}(X)$ is contained in the intersection of all maximal filters of X .*

Proof. Let $x \in \mathcal{D}(X)$ and M a maximal filter of X such that $x \notin M$. Then $\langle M \cup \{x\} \rangle = X$. Thus $0 \in \langle M \cup \{x\} \rangle$. Hence $x^n * 0 \in M$ for some $n \in \mathbb{Z}^+$. Since $x * 0 = xN = 0$, we get $x^n * 0 = x^{n-1} * (x * 0) = x^{n-1} * 0 = x^{n-2} * (x * 0) = \dots x * 0 = 0$. Hence $0 = xN = x * 0 = x^n * 0 \in M$, which is a contradiction. Thus $x \in M$ for all maximal filters of X . Therefore $\mathcal{D}(X) \subseteq M$ for all maximal filters of X . \square

The following result is an immediate consequence of the above two results.

THEOREM 4.9. *The following are equivalent in a transitive and dense BE -algebra X .*

- (1) X has a unique maximal filter;
- (2) $\mathcal{D}(X)$ is maximal;
- (3) $\mathcal{D}(X)$ is the intersection of all maximal filters.

In the following theorem, a set of equivalent conditions is derived for every BE -algebra to become a dense BE -algebra.

THEOREM 4.10. *The following are equivalent in a transitive BE -algebra X .*

- (1) X is dense;
- (2) $(a * b)N = (b * a)N$ for all $0 \neq a, b \in X$;
- (3) $a * bNN = b * aNN$ for all $0 \neq a, b \in X$;
- (4) $aNN * bNN = bNN * aNN$ for all $0 \neq a, b \in X$.

Proof. (1) \Rightarrow (2): Assume that X is dense. Let $0 \neq a, b \in X$. Suppose $a * b = 0$. Then we get $b \leq a * b = 0$, which is a contradiction. Hence, it yields $a * b \neq 0$. Similarly $b * a \neq 0$. Then there exist $0 \neq c \in X$ and $0 \neq d \in X$ such that $a * b = c$ and $b * a = d$. Since X is dense and $c \neq 0$, we get $(a * b)N = cN = 0$. Similarly, we can obtain $(b * a)N = dN = 0$. Therefore $(a * b)N = (b * a)N$ for all $0 \neq a, b \in X$.

(2) \Rightarrow (3): Assume that $(a * b)N = (b * a)N$ for all $0 \neq a, b \in X$. Then

$$\begin{aligned}
 1 &= (b * bNN)NN \\
 &= (bNN * b)NN \\
 &\leq ((a * bNN) * (a * b))NN \\
 &\leq ((a * bNN) * (a * b)NN)NN \\
 &= (a * bNN) * (a * b)NN
 \end{aligned}$$

Hence, it yields that $a * bNN \leq (a * b)NN$. Again, since $b \leq bNN$, we get $(a * b)NN \leq (a * bNN)NN = a * bNN$. From the above two observations, it is concluded that $(a * b)NN = a * bNN$. Therefore $a * bNN = (a * b)NN = (b * a)NN = b * aNN$ for all $0 \neq a, b \in X$.

(3) \Rightarrow (4): Assume that the condition (3) holds. Let $a, b \in X - \{0\}$. Then by Proposition 2.4 (4), it follows that $aNN * bNN = a * bNN = b * aNN = bNN * aNN$.

(4) \Rightarrow (1): Assume that the condition (4) holds. Let $0 \neq a \in X$. Then by Proposition 2.4 (4), it implies that $aNN = 1 * aNN = 1NN * aNN = aNN * 1NN = a * 1NN = a * 1 = 1$. Hence, it yields that $aN = 0$. Therefore X is dense. \square

If X is dense, then it can be routinely verified that X/F is also dense. Though the converse of this statement is not true, in the following, a necessary and sufficient condition is derived for the quotient algebra X/F to become dense.

THEOREM 4.11. *If F is a maximal filter of a self-distributive BE -algebra X , then X/F is dense.*

Proof. Let F be a maximal filter of a self-distributive BE -algebra X . Clearly $a \in F$ for all $0 \neq a \in X$. Hence $aNN \in F$ for all $0 \neq a \in X$. Let $F_0 \neq F_x \in X/F$ be a non-zero element. Then clearly $x \neq 0$. Since F is maximal, we get $x \in F$ and hence $xNN \in F$. Since $1 \in F$ and F is a congruence class, it yields $F_{xNN} = F_1$. It implies $F_xN = F_{xN} = F_0$. Thus F_x is a dense element in X/F . Therefore X/F is dense. \square

For any $x \in F$, it can be seen that $F_x = F_{xNN}$. For, consider $x \in X$. Then clearly $x * xNN \in F$. Since $x \in F$, we get $xNN \in F$. Therefore it concludes that $(x, xNN) \in \theta_F$. In the following theorem, a set of equivalent conditions is derived for every closed filter of a self-distributive BE -algebra to become a maximal filter.

THEOREM 4.12. *Let X be a self-distributive BE -algebra. Then the following conditions are equivalent.*

- (1) X is dense;
- (2) for every filter F , X/F is dense;
- (3) every closed filter is maximal;
- (4) $\mathcal{D}(X)$ is maximal.

Proof. (1) \Rightarrow (2): It is obvious.

(2) \Rightarrow (3): Assume that the condition (2) holds. Let F be a closed filter of X . Then by the hypothesis, we get X/F is dense. Let $0 \neq x \in X$. It is enough to show that $x \in F$. Since $x \neq 0$, we get $F_x \neq F_0$. Since X/F is dense, it yields $F_{xN} = F_xN = F_0$. Hence $F_{xNN} = F_{0N} = F_1$ and so $(xNN, 1) \in \theta_F$. Thus $xNN = 1 * xNN \in F$. Since F is a closed filter of X , it yields that $x \in F$. Therefore F is a maximal filter of X .

(3) \Rightarrow (4): Since $\mathcal{D}(X)$ is closed, it follows immediately.

(4) \Rightarrow (1): Assume that $\mathcal{D}(X)$ is maximal. By Theorem 4.10, it is clear. \square

The notion of D -filters of BE -algebras is now introduced in the following:

DEFINITION 4.13. A proper filter F of a BE -algebra is called D -filter if $\mathcal{D}(X) \subseteq F$.

In any BE -algebra X , clearly $\mathcal{D}(X)$ is a D -filter and every closed filter is a D -filter. Since every maximal filter of a transitive BE -algebra is closed it is also a D -filter. In the following theorem, some equivalent conditions are derived for every filter of a transitive BE -algebra to become D -filter.

THEOREM 4.14. *The following conditions are equivalent in a transitive BE -algebra.*

- (1) Every filter is a D -filter;
- (2) every principal filter is a D -filter;
- (3) X has a unique dense element.

Proof. (1) \Rightarrow (2) is clear.

(2) \Rightarrow (3): Assume that every principal filter is a D -filter. Hence, we get that $\langle 1 \rangle$ is a D -filter of X . Thus, it implies that $\mathcal{D}(X) \subseteq \langle 1 \rangle$. Hence, it concludes that $\mathcal{D}(X) = \{1\}$. Therefore X has the unique dense element, precisely 1.

(3) \Rightarrow (1): It is obvious. \square

For any transitive BE -algebra X , define a mapping $\nu : X \rightarrow X/\mathcal{D}(X)$ by $\nu(x) = \mathcal{D}(X)_x$ for all $x \in X$. Then clearly ν is an epimorphism. In the following theorem, we derive a set of equivalent conditions for ν to become an isomorphism.

THEOREM 4.15. *The following are equivalent in a transitive BE -algebra X .*

- (1) Every filter is a D -filter;
- (2) for any $x, y \in X$, $(x * y)N = 0, (y * x)N = 0$ implies $x = y$;
- (3) X is isomorphic to $X/\mathcal{D}(X)$.

Proof. (1) \Rightarrow (2): Assume that every filter of X is a D -filter. Then by above theorem, X has a unique dense element. Let $x, y \in X$. Assume that $(x * y)N = 0$ and $(y * x)N = 0$. Hence $x * y \in \mathcal{D}(X) = \{1\}$ and $y * x \in \mathcal{D}(X) = \{1\}$. Hence $x * y = 1$ and $y * x = 1$. Therefore, it concludes that $x = y$.

(2) \Rightarrow (3): Assume that condition (2) holds. Clearly $\nu : X \rightarrow X/\mathcal{D}(X)$ is an epimorphism. Let $x, y \in X$ be such that $\mathcal{D}(X)_x = \mathcal{D}(X)_y$. Hence $x * y \in \mathcal{D}(X)$ and $y * x \in \mathcal{D}(X)$, which yield $(x * y)N = 0$ and $(y * x)N = 0$. Then by condition (2), we get $x = y$. Therefore ν is one-one and so ν is an isomorphism.

(3) \Rightarrow (1): Assume that X is isomorphic to $X/\mathcal{D}(X)$. Let $x, y \in \mathcal{D}(X)$. Clearly $x * y, y * x \in \mathcal{D}(X)$. Hence $\mathcal{D}(X)_x = \mathcal{D}(X)_y$. Since ν is an isomorphism, it yields $x = y$. Hence X has a unique dense element. Thus every filter of X is a D -filter. \square

It is clear that every closed filter of a BE -algebra is a D -filter but the converse is not true. For consider the filter $F = \{1, b, c\}$ of the BE -algebra X given in the Example 4.6. Clearly F is a D -filter but not closed. However, in the following theorem, a set of equivalent conditions is derived for every D -filter to become closed.

THEOREM 4.16. *The following are equivalent in a self-distributive BE-algebra X .*

- (1) *Every D -filter is closed;*
- (2) *$\mathcal{D}(X)_x = \mathcal{D}(X)_{xNN}$ for all $x \in X$;*
- (3) *for any D -filter F , $xNN * x \in F$ for all $x \in X$.*

Proof. (1) \Rightarrow (2): Assume that every D -filter is a closed. Let $x \in X$. Clearly $\langle \mathcal{D}(X) \cup \{xNN\} \rangle$ is a D -filter of X . By the hypothesis, $\langle \mathcal{D}(X) \cup \{xNN\} \rangle$ is closed. Since $xNN \in \langle \mathcal{D}(X) \cup \{xNN\} \rangle$, we get $x \in \langle \mathcal{D}(X) \cup \{xNN\} \rangle$. Therefore $xNN * x \in \mathcal{D}(X)$. Clearly $x * xNN = 1 \in \mathcal{D}(X)$. Hence $(x, xNN) \in \theta_{\mathcal{D}(X)}$. Therefore $\mathcal{D}(X)_x = \mathcal{D}(X)_{xNN}$.

(2) \Rightarrow (3): Assume that $\mathcal{D}(X)_x = \mathcal{D}(X)_{xNN}$ for all $x \in X$. Let F be a D -filter of X . Let $x \in X$. Then by (2), we get $(x, xNN) \in \theta_{\mathcal{D}(X)}$. Therefore $xNN * x \in \mathcal{D}(X) \subseteq F$.

(3) \Rightarrow (1): Assume that the condition (3) holds. Let F be a D -filter of X . Let $x \in X$ and $xNN \in F$. By condition (3), we get that $xNN * x \in F$. Since F is a filter, it yields that $x \in F$. Therefore F is closed. \square

References

- [1] S. S. Ahn, Y. H. Kim, and J. M. Ko, *Filters in commutative BE-algebras*, Commun. Korean. Math. Soc. **27** (2012), no.2, 233-242.
- [2] Z. Ciloglu and Y. Ceven, *Commutative and bounded BE-algebras*, Algebra, **2013** (2013), Article ID 473714, 5 pages.
- [3] J. Gispert and A. Torrens, *Boolean representation of bounded BCK-algebras*, Soft Comput. **12** (2008), 941-954.
- [4] K. Iseki and S. Tanaka, *An introduction to the theory of BCK-algebras*, Math. Japon. **23** (1979), no.1, 1-26.
- [5] H. S. Kim and Y. H. Kim, *On BE-algebras*, Sci. Math. Jpn. **66** (2007), no.1, 113-116.
- [6] B. L. Meng, *On filters in BE-algebras*, Sci. Math. Japon, Online, e-2010, 105-111.
- [7] C. Muresan, *Dense Elements and Classes of Residuated Lattices*, Bull. Math. Soc. Sci. Math. Roumanie Tome **53** (2010), no. 1, 11-24.
- [8] D. Piciu and D. D. Tascu, *The localization of commutative bounded BCK-algebras*, Adv. Pure Math. **1** (2011), 367-377.
- [9] Y. H. Yon, S. M. Lee, and K. H. Kim, *On congruences and BE-relations in BE-algebras*, Int. Math. Forum, **5** (2010), no.46, 2263-2270.
- [10] A. Walendziak, *On commutative BE-algebras*, Sci. Math. Jpn. **69** (2009), no.2, 281-284.

*

Department of Mathematics, Aditya Engineering College(A),
Surampalem, Kakinada,
Andhra Pradesh, India-533 437.
E-mail: prabhakar_mb@yahoo.co.in

**

Department of Mathematics,
JNTUK University College of Engineering,
Vizianagaram, Andhra Pradesh, India-535003.
E-mail: valijntuv@gmail.com

Department of Mathematics, MVGR College of Engineering(A),
Vizianagaram, Andhra Pradesh, India-535005.
E-mail: mssraomaths35@rediffmail.com