

THE RELIABLE MODIFIED OF ADOMIAN DECOMPOSITION METHOD FOR SOLVING INTEGRO-DIFFERENTIAL EQUATIONS

AHMED A. HAMOUD* AND KIRTIWANT P. GHADLE**

ABSTRACT. In this article, we discussed semi-analytical approximated methods for solving mixed Volterra-Fredholm integro-differential equations, namely: Adomian decomposition method and modified Adomian decomposition method. Moreover, we prove the uniqueness results and convergence of the techniques. Finally, an example is included to demonstrate the validity and applicability of the proposed techniques.

1. Introduction

The integro-differential equations be an important branch of modern mathematics. It arises frequently in many applied areas which include engineering, electrostatics, mechanics, the theory of elasticity, potential, and mathematical physics [3, 4, 6, 10, 29].

In this work, we consider the mixed Volterra-Fredholm integro-differential equation of the second kind as follows:

$$(1.1) \quad \sum_{j=0}^k p_j(x, z) u_x^{(j)}(x, z) = f(x, z) + \int_a^x \int_{\Omega} K(x, z) G(u^{(l)}(x, z)) dx dz,$$

with the initial or boundary conditions:

$$(1.2) \quad \begin{aligned} u^{(r)}(a, z) &= \alpha_r(z), & r = 0, 1, 2, \dots, (n-1), \\ u^{(r)}(b, z) &= \beta_r(z), & r = n, (n+1), \dots, (k-1), \end{aligned}$$

Received May 05, 2019; Accepted August 21, 2019.

2010 Mathematics Subject Classification: Primary 45K05; Secondary 49M27.

Key words and phrases: Volterra-Fredholm integro-differential equation, Adomian decomposition method, modified Adomian decomposition method, approximate solution.

* Corresponding author.

where $u^{(j)}$ is the j^{th} derivative of the unknown function u that will be determined, $K(x, z)$ is the kernel of the equation, $f(x, z)$ and $p_j(x, z)$ are analytic functions, $G(u^{(l)}(x, z))$, $l \geq 0$ is nonlinear analytic function of u and λ , α_r , $0 \leq r \leq (n - 1)$ and β_r , $n \leq r \leq (k - 1)$ are real finite constants, $a \leq z \leq x$, $a \leq x \leq b$, $\Omega = [a, b]$. Recently, Wazwaz (2001) presented an efficient and numerical procedure for solving boundary value problems for higher-order integro-differential equations. A variety of methods, exact, approximate and purely numerical techniques are available to solve nonlinear integro-differential equations. These methods have been of great interest to several authors and used to solve many nonlinear problems. Some of these techniques are Adomian decomposition method [4, 25], modified Adomian decomposition method [30, 32], Variational iteration method [7, 11] and many techniques for solving integro-differential equations [2, 3, 6, 16, 17, 18, 19, 20, 21, 28].

More details about the sources where these equations arise can be found in physics, biology, and engineering applications as well as in advanced integral equations. Some works based on an iterative scheme have been focusing on the development of more advanced and efficient methods for integro-differential equations such as the variational iteration method which is a simple and Adomian decomposition method [8, 9, 25, 31], and the modified decomposition method for solving Volterra-Fredholm integral and integro-differential equations which is a simple and powerful method for solving a wide class of nonlinear problems [25]. The Taylor polynomial solution of integro-differential equations has been studied in [29]. The use of Lagrange interpolation in solving integro-differential equations was investigated by Marzban [27].

A variety of powerful methods has been presented, such as the homotopy analysis method [30], homotopy perturbation method [6, 30], operational matrix with Block-Pulse functions method [3], variational iteration method [5, 30] and the Adomian decomposition method [4, 25, 30].

Some fundamental works on various aspects of modifications of the Adomian's decomposition method are given by Araghi and Behzadi [1]. The modified form of Laplace decomposition method has been introduced by Manafianheris [26]. Babolian et. al, [3], applied the new direct method to solve nonlinear Volterra-Fredholm integral and integro-differential equation using operational matrix with block-pulse functions. The Laplace transform method with the Adomian decomposition method to establish exact solutions or approximations of the nonlinear Volterra integro-differential equations, Wazwaz [31].

Recently, the authors have used several methods for the numerical or the analytical solutions of linear and nonlinear Volterra and Fredholm integro-differential equations [11, 12, 13, 14, 15, 22, 24, 25, 30].

In this work, our aim is to solve a general form of nonlinear Volterra-Fredholm integro-differential equations using approximate methods, namely, Adomian decomposition method and modified Adomian decomposition method. Also, we prove the uniqueness results and convergence of the techniques.

2. Nonlinear Mixed Volterra-Fredholm Integro-Differential Equation of Second Kind

We consider the mixed Volterra-Fredholm integro-differential equation of the second kind as follows:

$$(2.1) \quad \sum_{j=0}^k p_j(x, z) u_x^{(j)}(x, z) = f(x, z) + \int_a^x \int_{\Omega} K(x, z) G(u^{(l)}(x, z)) dx dz,$$

We can write Eq.(2.1) as follows:

$$(2.2) \quad \begin{aligned} p_k(x, z) u_x^{(k)}(x, z) + \sum_{j=0}^{k-1} p_j(x, z) u_x^{(j)}(x, z) &= f(x, z) + \\ &+ \int_a^x \int_{\Omega} K(x, z) G(u^{(l)}(x, z)) dx dz, \\ u_x^{(k)}(x, z) &= \frac{f(x, z)}{p_k(x, z)} + \int_a^x \int_{\Omega} \frac{K(x, z) G(u^{(l)}(x, z))}{p_k(x, z)} dx dz \\ &\quad - \sum_{j=0}^{k-1} \frac{p_j(x, z)}{p_k(x, z)} u_x^{(j)}(x, z). \end{aligned}$$

Let us set L^{-1} is the multiple integration operator as follows:

$$(2.3) \quad L^{-1}(\cdot) = \int_a^x \int_a^x \cdots \int_a^x (\cdot) dx dx \dots dx, \quad k - \text{times}.$$

$$(2.4) \quad \begin{aligned} u(x, z) &= L^{-1} \left\{ \frac{f(x, z)}{p_k(x, z)} \right\} + g_0(z) + \sum_{r=0}^{k-2} \int_a^x \frac{(x-z)^r}{r!} g_{r+1}(z) dz \\ &+ L^{-1} \left\{ \int_a^x \int_{\Omega} \frac{K(x, z) G(u^{(l)}(x, z))}{p_k(x, z)} dx dz \right\} \\ &- L^{-1} \left\{ \sum_{j=0}^{k-1} \frac{p_j(x, z)}{p_k(x, z)} u_x^{(j)}(x, z) \right\}. \end{aligned}$$

We can obtain the term $g_0(z) + \sum_{r=0}^{k-2} \int_a^x \frac{(x-z)^r}{r!} g_{r+1}(z) dz$ from the initial conditions. From [23], we have

$$(2.5) \quad \begin{aligned} & L^{-1} \left\{ \int_a^x \int_{\Omega} \frac{K(x, z) G(u^{(l)}(x, z))}{p_k(x, z)} dx dz \right\} \\ &= \int_a^x \int_{\Omega} \frac{(x-z)^k K(x, z) G(u^{(l)}(x, z))}{(k!) p_k(x, z)} dx dz \end{aligned}$$

also

$$(2.6) \quad L^{-1} \left\{ \sum_{j=0}^{k-1} \frac{p_j(x, z)}{p_k(x, z)} u_x^{(j)}(x, z) \right\} = \sum_{j=0}^{k-1} \int_a^x \frac{(x-z)^{k-1} p_j(x, z)}{(k-1)! p_k(x, z)} u_x^{(j)}(x, z) dz.$$

By substituting Eq.(2.5) and Eq.(2.6) in Eq.(2.4) we obtain

$$(2.7) \quad \begin{aligned} u(x, z) &= L^{-1} \left\{ \frac{f(x, z)}{p_k(x, z)} \right\} + g_0(z) + \sum_{r=0}^{k-2} \int_a^x \frac{(x-z)^r}{r!} g_{r+1}(z) dz \\ &+ \int_a^x \int_{\Omega} \frac{(x-z)^k K(x, z) G(u^{(l)}(x, z))}{(k!) p_k(x, z)} dx dz \\ &- \sum_{j=0}^{k-1} \int_a^x \frac{(x-z)^{k-1} p_j(x, z)}{(k-1)! p_k(x, z)} u_x^{(j)}(x, z) dz. \end{aligned}$$

We set,

$$\begin{aligned} L^{-1} \left\{ \frac{f(x, z)}{p_k(x, z)} \right\} + g_0(z) + \sum_{r=0}^{k-2} \int_a^x \frac{(x-z)^r}{r!} g_{r+1}(z) dz &= G_1(x, z), \\ \int_{\Omega} \frac{(x-z)^k K(x, z)}{(k!) p_k(x, z)} dx &= K_1(x, z), \end{aligned}$$

and

$$\frac{(x-z)^{k-1} p_j(x, z)}{(k-1)! p_k(x, z)} = K_2(x, z).$$

So, we have nonlinear integro-differential equation as follows:

$$(2.8) \quad \begin{aligned} u(x, z) &= G_1(x, z) + \int_a^x K_1(x, z) G(u^{(l)}(x, z)) dz \\ &- \sum_{j=0}^{k-1} \int_a^x K_2(x, z) u_x^{(j)}(x, z) dz. \end{aligned}$$

3. Description of the Methods

In this section, we present the semi-analytical techniques based on ADM and MADM to solve nonlinear integro-differential equation Eq.(2.8):

3.1. Adomian Decomposition Method (ADM)

In this part, the ADM is used to find approximate of nonlinear integro-differential equation Eq.(2.8). The nonlinear terms $G(u^{(l)}(x, z))$ and $D^j(u(x, z))$, ($D^j = \frac{\partial^j u(x, z)}{\partial x^j}$ is derivative operator), are usually represented by an infinite series of the so called Adomian polynomials as follows:

$$G(u^{(l)}(x, z)) = \sum_{i=0}^{\infty} A_i, \quad D^j(u(x, z)) = \sum_{i=0}^{\infty} B_{ij},$$

where A_i and B_{ij} ($i \geq 0, j = 0, 1, \dots, k-1$) are the Adomian polynomials were introduced in [30]. According to the ADM, we can write the iterative formula as follows:

$$(3.1) \quad u_0(x, z) = G_1(x, z),$$

$$u_{n+1}(x, z) = \int_a^x K_1(x, z) A_n dz - \sum_{j=0}^{k-1} \int_a^x K_2(x, z) B_{nj} dz, \quad n \geq 0.$$

Then, $u(x, z) = \sum_{i=0}^n u_i(x, z)$ as the approximate solution.

3.2. Modified Adomian Decomposition Method (MADM)

In this part, the extended MADM [30] is used to find approximate of nonlinear integro-differential equation Eq.(2.8).

This method is based on the assumption that the function $G_1(x, z)$ can be divided into two parts, namely $G_{11}(x, z)$ and $G_{12}(x, z)$. Under this assumption we set

$$G_1(x, z) = G_{11}(x, z) + G_{12}(x, z).$$

According to the MADM, we can write the iterative formula as follows:

$$u_0(x, z) = G_{11}(x, z),$$

$$u_1(x, z) = G_{12}(x, z) + \int_a^x K_1(x, z) A_0 dz - \sum_{j=0}^{k-1} \int_a^x K_2(x, z) B_{0j} dz,$$

$$u_{n+1}(x, z) = \int_a^x K_1(x, z) A_n dz - \sum_{j=0}^{k-1} \int_a^x K_2(x, z) B_{nj} dz, \quad n \geq 1.$$

Then, $u(x, z) = \sum_{i=0}^n u_i(x, z)$ as the approximate solution.

4. Uniqueness and Convergence Results

In this section the uniqueness of the obtained solution and convergence of the methods are proved. Consider the Eq.(2.8), we assume $G_1(x, z)$ is bounded for all x, z in Ω and

$$|K_1(z, t)| \leq M_1, \quad |K_2(z, t)| \leq M_{1j}, \quad j = 0, 1, \dots, k-1.$$

Also, we suppose the nonlinear terms $G(u(x, z))$ and $D^j(u(x, z))$ are Lipschitz continuous with

$$\begin{aligned} |G(u(x, z)) - G(u^*(x, z))| &\leq d |u(x, z) - u^*(x, z)| \\ |D^j(u(x, z)) - D^j(u^*(x, z))| &\leq C_j |u(x, z) - u^*(x, z)|. \end{aligned}$$

If we set, $\gamma = (b-a)(dM_1 + kCM)$, $C = \max |C_j|$, $M = \max |M_{1j}|$. Then the following theorems can be proved by using the above assumptions.

THEOREM 4.1. *Two-dimensional nonlinear Volterra-Fredholm integro-differential equation, has a unique solution whenever $0 < \gamma < 1$.*

Proof. Let u and u^* be two different solutions of Eq.(2.8) then

$$\begin{aligned} |u(x, z) - u^*(x, z)| &= \left| \int_a^x K_1(x, z)G(u^{(l)}(x, z))dz - \sum_{j=0}^{k-1} \int_a^x K_2(x, z)D^j(u(x, z))dz \right. \\ &\quad \left. - \int_a^x K_1(x, z)G(u^{(l)*}(x, z))dz + \sum_{j=0}^{k-1} \int_a^x K_2(x, z)D^j(u^*(x, z))dz \right| \\ &\leq \left| \int_a^x K_1(x, z)[G(u^{(l)}(x, z)) - G(u^{(l)*}(x, z))]dz \right. \\ &\quad \left. - \sum_{j=0}^{k-1} \int_a^x K_2(x, z)[D^j(u(x, z)) - D^j(u^*(x, z))]dz \right| \\ &\leq \int_a^x |K_1(x, z)| |G(u^{(l)}(x, z)) - G(u^{(l)*}(x, z))| dz \\ &\quad + \sum_{j=0}^{k-1} \int_a^x |K_2(x, z)| |D^j(u(x, z)) - D^j(u^*(x, z))| dz \\ &\leq M_1 d |u(x, z) - u^*(x, z)|(b-a) + kMC |u(x, z) - u^*(x, z)|(b-a) \\ &\leq (b-a)(M_1 d + kMC) |u(x, z) - u^*(x, z)| \\ &= \gamma |u(x, z) - u^*(x, z)|. \end{aligned}$$

So,

$$\left| u(x, z) - u^*(x, z) \right| \leq \gamma \left| u(x, z) - u^*(x, z) \right|,$$

we get $(1 - \gamma) \left| u - u^* \right| \leq 0$. Since $0 < \gamma < 1$, so $\left| u - u^* \right| = 0$, therefore, $u = u^*$ and this completes the proof. \square

THEOREM 4.2. *The series solution $u(x, z) = \sum_{i=0}^{\infty} u_i(x, z)$ of Eq.(1.1) using MADM convergence when $0 < \gamma < 1$ and $\|u_1\| < \infty$.*

Proof. Denote as $(C[J], \|\cdot\|)$ the Banach space of all continuous functions on J with the norm $\|f(x, z)\| = \max |f(x, z)|$ for all x, z in J . Define the sequence of partial sums s_n , let s_n and s_m be arbitrary partial sums with $n \geq m$. We are going to prove that $s_n = \sum_{i=0}^n u_i$ is a Cauchy sequence in this Banach space:

$$\begin{aligned} \|s_n - s_m\| &= \max_{\forall x, z \in J} \left| s_n - s_m \right| \\ &= \max_{\forall x, z \in J} \left| \sum_{i=0}^n u_i(x, z) - \sum_{i=0}^m u_i(x, z) \right| \\ &= \max_{\forall x, z \in J} \left| \sum_{i=m+1}^n u_i(x, z) \right| \\ &= \max_{\forall x, z \in J} \left| \sum_{i=m+1}^n \left[\int_a^x K_1(x, z) A_i dz - \sum_{j=0}^{k-1} \int_a^x K_2(x, z) L_{i_j} dz \right] \right| \\ &= \max_{\forall x, z \in J} \left| \int_a^x K_1(x, z) \left(\sum_{i=m}^{n-1} A_i \right) dz - \sum_{j=0}^{k-1} \int_a^x K_2(x, z) \left(\sum_{i=m}^{n-1} L_{i_j} \right) dz \right|. \end{aligned}$$

From [23], we have

$$\sum_{i=m}^{n-1} A_i = G(s_{n-1}) - G(s_{m-1}),$$

$$\sum_{i=m}^{n-1} L_{i_j} = D^j(s_{n-1}) - D^j(s_{m-1}).$$

So,

$$\begin{aligned}
\|s_n - s_m\| &= \max_{\forall x, z \in J} \left| \int_a^x K_1(x, z)(G(s_{n-1}) - G(s_{m-1}))dz \right. \\
&\quad \left. - \sum_{j=0}^{k-1} \int_a^x K_2(x, z)(D^j(s_{n-1}) - D^j(s_{m-1}))dz \right| \\
&\leq \max_{\forall x, z \in J} \left(\int_a^x |K_1(x, z)||G(s_{n-1}) - G(s_{m-1})|dz \right. \\
&\quad \left. + \sum_{j=0}^{k-1} \int_a^x |K_2(x, z)||D^j(s_{n-1}) - D^j(s_{m-1})|dz \right) \\
&\leq \max_{\forall x, z \in J} \left(M_1 d |s_{n-1} - s_{m-1}|(b-a) \right. \\
&\quad \left. + \sum_{j=0}^{k-1} (M_{1_j} C_j) |s_{n-1} - s_{m-1}|(b-a) \right) \\
&\leq M_1 d |s_{n-1} - s_{m-1}|(b-a) + kMC |s_{n-1} - s_{m-1}|(b-a) \\
&= (M_1 d + kMC)(b-a) |s_{n-1} - s_{m-1}| \\
&= \gamma \|s_{n-1} - s_{m-1}\|.
\end{aligned}$$

Let $n = m + 1$, then

$$\|s_n - s_m\| \leq \gamma \|s_m - s_{m-1}\| \leq \gamma^2 \|s_{m-1} - s_{m-2}\| \leq \dots \leq \gamma^m \|s_1 - s_0\|.$$

So,

$$\begin{aligned}
\|s_n - s_m\| &\leq \|s_{m+1} - s_m\| + \|s_{m+2} - s_{m+1}\| + \dots + \|s_n - s_{n-1}\| \\
&\leq [\gamma^m + \gamma^{m+1} + \dots + \gamma^{n-m-1}] \|s_1 - s_0\| \\
&\leq \gamma^m [1 + \gamma + \gamma^2 + \dots + \gamma^{n-2m-1}] \|s_1 - s_0\| \\
&\leq \frac{1 - \gamma n - m}{1 - \gamma} \|u_1\|.
\end{aligned}$$

Since $0 < \gamma < 1$, we have $(1 - \gamma^{nm}) < 1$, then

$$\|s_n - s_m\| \leq \frac{\gamma m}{1 - \gamma} \|u_1\|.$$

But $|u_1| < \infty$ (since $G_1(x, z)$ is bounded), so, as $m \rightarrow \infty$, then $\|s_n - s_m\| \rightarrow 0$. We conclude that s_n is a Cauchy sequence, therefore the series is convergence and the proof is complete. \square

5. Numerical Example

In this section, we present the semi-analytical techniques based on ADM and MADM, to solve Volterra-Fredholm integro-differential equations:

Example 1.

Consider the Volterra-Fredholm integro-differential equation as follow (5.1)

$$u_x''(x, z) + u(x, z) \sin(xz) = xz \sin(xz) - \frac{1}{3}z^3 + \int_0^x \int_0^1 xz u_x'(x, z) dx dz,$$

with the initial conditions

$$u_x'(0, z) = u(0, z) = 0.$$

The exact solution is $u(x, z) = xz$, $\epsilon = 10^{-2}$.

TABLE 1. Numerical Results of the Example 1.

(x,z)	Er(MADM _{n=10})	Er(ADM _{n=10})
(0.1,0.05)	0.074634	0.084475
(0.2,0.14)	0.074727	0.084638
(0.4,0.23)	0.075597	0.085795
(0.6,0.27)	0.077224	0.085867
(0.85,0.35)	0.078538	0.085968

6. Conclusion

In this work, the ADM and MADM have been successfully employed to obtain the approximate solutions of a mixed Volterra-Fredholm integro-differential equation. Moreover, we proved the uniqueness results and convergence of the techniques. The results show that these methods are very efficient, convenient and can be adapted to fit a larger class of problems. The comparison reveals that although the numerical results of these methods are similar approximately, MADM is the easiest, the most efficient and convenient.

References

- [1] M.A. Araghi and S.S. Behzadi, *Solving nonlinear Volterra-Fredholm integro-differential equations using the modified Adomian decomposition method*, Comput. Methods in Appl. Math., **9** (2009), 321–331.
- [2] S.H. Behiry and S.I. Mohamed, *Solving high-order nonlinear Volterra-Fredholm integro-differential equations by differential transform method*, Natural Science, **4** (2012), no. 8, 581–587.
- [3] E. Babolian, Z. Masouri, and S. Hatamzadeh, *New direct method to solve nonlinear Volterra-Fredholm integral and integro differential equation using operational matrix with Block-Pulse functions*, Progress in Electromagnetic Research, **B 8** (2008), 59–76.
- [4] S.M. El-Sayed, D. Kaya, and S. Zarea, *The decomposition method applied to solve high-order linear Volterra-Fredholm integro-differential equations*, International Journal of Nonlinear Sciences and Numerical Simulation, **5** (2004), no. 2, 105–112.
- [5] F.S. Fadhel, A.O. Mezaal, and S.H. Salih, *Approximate solution of the linear mixed Volterra-Fredholm integro-differential equations of second kind by using variational iteration method*, Al- Mustansiriyah, J. Sci., **24** (2013), no. 5, 137–146.
- [6] M. Ghasemi, M. kajani, and E. Babolian, *Application of He's homotopy perturbation method to nonlinear integro differential equations*, Appl. Math. Comput., **188** (2007), 538–548.
- [7] J.H. He and S.Q. Wang, *Variational iteration method for solving integro-differential equations*, Phys. Lett., **A 367** (2007), 188–191.
- [8] A.A. Hamoud and K.P. Ghadle, *Existence and uniqueness of the solution for Volterra-Fredholm integro-differential equations*, Journal of Siberian Federal University. Mathematics & Physics, **11** (2018), no. 6, 692–701.
- [9] A.A. Hamoud and K.P. Ghadle, *Modified Laplace decomposition method for fractional Volterra-Fredholm integro-differential equations*, Journal of Mathematical Modeling, **6** (2018), no. 1, 91–104.
- [10] A.A. Hamoud and K.P. Ghadle, *Homotopy analysis method for the first order fuzzy Volterra-Fredholm integro-differential equations*, Indonesian J. Elec. Eng. & Comp. Sci., **11** (2018), no. 3, 857–867.
- [11] A.A. Hamoud and K.P. Ghadle, *Some new existence, uniqueness and convergence results for fractional Volterra-Fredholm integro-differential equations*, J. Appl. Comp. Mech., **5** (2019), no. 1, 58–69.
- [12] A.A. Hamoud and K.P. Ghadle, *Existence and uniqueness of solutions for fractional mixed Volterra-Fredholm integro-differential equations*, Indian J. Math., **60** (2018), no. 3, 375–395.
- [13] A.A. Hamoud and K.P. Ghadle, *The approximate solutions of fractional Volterra-Fredholm integro-differential equations by using analytical techniques*, Probl. Anal. Issues Anal., **7(25)** (2018), no. 1, 41–58.
- [14] A.A. Hamoud, K.P. Ghadle, and S.M. Atshan, *The approximate solutions of fractional integro-differential equations by using modified Adomian decomposition method*, Khayyam Journal of Mathematics, **5** (2019), no. 1, 21–39.

- [15] A.A. Hamoud and K.P. Ghadle, *The reliable modified of Laplace Adomian decomposition method to solve nonlinear interval Volterra-Fredholm integral equations*, Korean J. Math., **25** (2017), no. 3, 323–334.
- [16] A.A. Hamoud, L.A. Dawood, K.P. Ghadle, and S.M. Atshan, *Usage of the modified variational iteration technique for solving Fredholm integro-differential equations*, International Journal of Mechanical and Production Engineering Research and Development , **9** (2019), no. 2, 895–902.
- [17] A.A. Hamoud, K.H. Hussain, and K.P. Ghadle, *The reliable modified Laplace Adomian decomposition method to solve fractional Volterra-Fredholm integro-differential equations*, Dynamics of Continuous, Discrete and Impulsive Systems Series B: Applications & Algorithms, **26** (2019), 171–184.
- [18] A.A. Hamoud and K.P. Ghadle, *Modified Adomian decomposition method for solving fuzzy Volterra-Fredholm integral equations*, J. Indian Math. Soc., **85** (2018), no. 1-2, 52–69.
- [19] A.A. Hamoud and K.P. Ghadle, M. Bani Issa, Giniswamy, *Existence and uniqueness theorems for fractional Volterra-Fredholm integro-differential equations*, Int. J. Appl. Math., **31** (2018), no. 3, 333–348.
- [20] A.A. Hamoud, A.D. Azeez, and K.P. Ghadle, *A study of some iterative methods for solving fuzzy Volterra-Fredholm integral equations*, Indonesian J. Elec. Eng. & Comp. Sci., **11** (2018), no. 3, 1228–1235.
- [21] A.A. Hamoud and K.P. Ghadle, *Usage of the homotopy analysis method for solving fractional Volterra-Fredholm integro-differential equation of the second kind*, Tamkang Journal of Mathematics, **49** (2018), no. 4, 301–315.
- [22] A.A. Hamoud, M. Bani Issa, and K.P. Ghadle, M. Abdulghani, *Existence and convergence results for caputo fractional Volterra integro-differential equations*, Journal of Mathematics and Applications, **41** (2018), 109–122.
- [23] A.A. Hamoud, N.M. Mohammed, K.P. Ghadle, and S.L. Dhondge, *Solving integro-differential equations by using numerical techniques*, International Journal of Applied Engineering Research, **14** (2019), no. 14, 3219–3225.
- [24] A.A. Hamoud, M. Bani Issa, and K.P. Ghadle, *Existence and uniqueness results for nonlinear Volterra-Fredholm integro-differential equations*, Nonlinear Functional Analysis and Applications, **23** (2018), no. 4, 797–805.
- [25] A.M. Jerri, *Introduction to Integral Equations with Applications*, New York, Wiley, 1999.
- [26] J. Manafianheris, *Solving the integro-differential equations using the modified Laplace Adomian decomposition method*, Journal of Mathematical Extension, **6** (2012), no. 1, 41–55.
- [27] H.R. Marzban and S.M. Hoseini, *Solution of nonlinear Volterra-Fredholm integro-differential equations via hybrid of Block-Pulse functions and Lagrange interpolating polynomials*, Hindawi Publishing Corporation, Advances in Numerical Analysis, **868** (2012), no. 279, 1–14.
- [28] S.B. Shadan, *The use of iterative method to solve two-dimensional nonlinear Volterra-Fredholm integro-differential equations*, J. of Communication in Numerical Analysis, (2012), 1–20.
- [29] Y. Salih and S. Mehmet, *The approximate solution of higher order linear Volterra-Fredholm integro-differential equations in term of Taylor polynomials*, Appl. Math. Comput., **112** (2000), 291–308.

- [30] A.M. Wazwaz, *Linear and Nonlinear Integral Equations Methods and Applications*, Springer Heidelberg Dordrecht London New York, 2011.
- [31] A.M. Wazwaz, *The combined Laplace transform-Adomian decomposition method for handling nonlinear Volterra integro-differential equations*, Appl. Math. Comput., **216** (2010), 1304–1309.
- [32] A.M. Wazwaz, *A reliable modification of Adomian decomposition method*, Appl. Math. Comput., **102** (1999), 77–86.

*

Department of Mathematics
Dr. Babasaheb Ambedkar Marathwada University
Aurangabad-431 004, India
Department of Mathematics
Taiz University, Taiz, Yemen
E-mail: drahmedselwi985@gmail.com

**

Department of Mathematics
Dr. Babasaheb Ambedkar Marathwada University
Aurangabad-431 004, India
E-mail: drkp.ghadle@gmail.com