

ON THE STABILITY OF THE GENERAL SEXTIC FUNCTIONAL EQUATION

ICK-SOON CHANG*, YANG-HI LEE**, AND JAIOK ROH***

ABSTRACT. The general sextic functional equation is a generalization of many functional equations such as the additive functional equation, the quadratic functional equation, the cubic functional equation, the quartic functional equation and the quintic functional equation. In this paper, motivating the method of Găvruta [J. Math. Anal. Appl., **184** (1994), 431–436], we will investigate the stability of the general sextic functional equation.

1. Introduction

In this paper, let V , X , and Y be a real vector space, a real normed space, and a real Banach space, respectively. Ulam [25] raised the question about the stability of group homomorphisms in 1940 and Hyers [8] gave a partial answer to this question by solving the stability of the Cauchy functional equation in the following year. Since then, many mathematicians have generalized the Hyers's result [6, 7, 9, 11, 15, 18, 20, 24]. In particular, Găvruta [7] generalized the result of Hyers as follow :

PROPOSITION 1.1. *Let $(G, +)$ be an abelian group and $\varphi : G^2 \rightarrow [0, \infty)$ be a function such that*

$$\tilde{\varphi}(x, y) := \sum_{k=0}^{\infty} 2^{-k} \varphi(x, y) < \infty$$

for all $x, y \in G$. If $f : G \rightarrow Y$ is a mapping such that

$$\|f(x + y) - f(x) - f(y)\| \leq \varphi(x, y)$$

Received July 08, 2021; Accepted August 07, 2021.

2010 Mathematics Subject Classification: Primary 39B82, Secondary 39B52.

Key words and phrases: sextic functional equations, stability, Hyers-Ulam stability.

This work was partially supported by Data Science Convergence Research Center of Hallym University.

for all $x, y \in G$, then there exists a unique additive mapping $T : G \rightarrow Y$ such that

$$\|f(x) - T(x)\| \leq \frac{1}{2} \tilde{\varphi}(x, x)$$

for all $x \in G$.

In this paper, motivating Proposition 1.1, we will study the stability of the general sextic functional equation :

$$(1.1) \quad \sum_{i=0}^7 {}_7C_i (-1)^{7-i} f(x + iy) = 0.$$

More detailed term for the concept of a general sextic mapping can be found in Baker's paper [3] by the term, generalized polynomial mapping of degree at most 6.

Jun-Kim[10] have previously studied the stability of a general cubic functional equation and others studied the stability of a general quadratic functional equation, a general cubic functional equation, and a general quartic functional equation(refer to [13, 14, 16, 19]). Also, many mathematicians have investigated the stability of the sextic functional equation (refer to [1, 2, 4, 5, 23, 21, 22]). But they worked on a special sextic functional equation. The stability of the general sextic functional equation have been investigated in Lee [17] and Roh-Lee-Jung [12]. Roh-Lee-Jung [12] have proved the stability of the general sextic functional equation by applying the fixed point theorem in the sense of Cădariu and Radu. And Lee [17] proved the Hyers-Ulam-Rassias stability of the general sextic functional equation as follows :

PROPOSITION 1.2. (Theorem 3 in [17]) *Let $p \neq 1, 2, 3, 4, 5, 6$ be a fixed nonnegative real number. Suppose that $f : X \rightarrow Y$ is a mapping such that*

$$\left\| \sum_{i=0}^7 {}_7C_i (-1)^{7-i} f(x + iy) \right\| \leq \theta (\|x\|^p + \|y\|^p)$$

for all $x, y \in X$. Then there exists a general sextic mapping F with $F(0) = 0$ and a constant $K(p)$ such that

$$\|f(x) - f(0) - F(x)\| \leq K(p) \theta \|x\|^p, \quad \text{for all } x \in X.$$

In this paper, we will investigate the stability of the general sextic functional equation in sense of Găvruta[7].

In fact, Lee [17] investigated the stability of the general sextic functional equation for the mapping f such that

$$Df(x, y) = \sum_{i=0}^7 {}_7C_i(-1)^{7-i} f(x + iy) \leq \theta(\|x\|^p + \|y\|^p)$$

and Roh-Lee-Jung [12] did for the mapping f such that

$$Df(x, y) = \sum_{i=0}^7 {}_7C_i(-1)^{7-i} f(x + iy) \leq \varphi(x, y),$$

where $\varphi : V^2 \rightarrow [0, \infty)$ is a function for which there exists a constant $0 < L < 1$ such that

$$\varphi(2x, 2y) \leq (4\sqrt{21} \cos \theta - 14)L\varphi(x, y)$$

for all $x, y \in V$ and θ is a real constant such that $0 < \theta < \frac{\pi}{4}$ and $\cos(3\theta) = \frac{-17}{21\sqrt{21}}$.

In this paper, we will investigate the stability of the general sextic functional equation for the mapping f such that

$$Df(x, y) = \sum_{i=0}^7 {}_7C_i(-1)^{7-i} f(x + iy) \leq \varphi(x, y),$$

where $\varphi : V^2 \rightarrow [0, \infty)$ is a function such that

$$\sum_{n=0}^{\infty} 64^n \varphi\left(\frac{x}{2^n}, \frac{y}{2^n}\right) < \infty \quad \text{or} \quad \sum_{n=0}^{\infty} 2^{-n} \varphi(2^n x, 2^n y) < \infty$$

for all $x, y \in V$.

2. Stability of a general sextic functional equation

Throughout this section, for a given mapping $f : V \rightarrow Y$, we use the following abbreviations :

$$f_o(x) := \frac{f(x) - f(-x)}{2}, \quad f_e(x) := \frac{f(x) + f(-x)}{2},$$

$$Df(x, y) := \sum_{i=0}^7 {}_7C_i(-1)^{7-i} f(x + iy),$$

$$\Gamma(x) := Df_o(-6x, 2x) + 6Df_o(-x, x) + 42Df_o(-2x, x) + 112Df_o(-3x, x),$$

$$\Delta f(x) := Df_e(-6x, 2x) + 8Df_e(-x, x) + 56Df_e(-2x, x) + 112Df_e(-3x, x)$$

for all $x, y \in V$. If \tilde{f} is a mapping defined by $\tilde{f}(x) = f(x) - f(0)$, then the mapping \tilde{f} satisfies the properties $D\tilde{f}(x, y) = Df(x, y)$ and $\tilde{f}(0) = 0$. By tedious computation we can get the equalities

$$(2.1) \quad \Gamma f(x) = f_o(8x) - 42f_o(4x) + 336f_o(2x) - 512f_o(x),$$

$$(2.2) \quad \Delta f(x) = f_e(8x) - 84f_e(4x) + 1344f_e(2x) - 4096f_e(x)$$

for all $x \in V$.

LEMMA 2.1. For a given mapping $f : V \rightarrow Y$ with $f(0) = 0$, let $J_n f, J'_n f : V \rightarrow Y$ be the mappings defined by

$$J_n f(x)$$

$$\begin{aligned} &:= \frac{4^n - 20 \cdot 16^n + 64 \cdot 64^n}{45} f_e\left(\frac{x}{2^n}\right) - \frac{80(4^n - 17 \cdot 16^n + 16 \cdot 64^n)}{45} f_e\left(\frac{x}{2^{n+1}}\right) \\ &+ \frac{1024(4^n - 5 \cdot 16^n + 4 \cdot 64^n)}{45} f_e\left(\frac{x}{2^{n+2}}\right) + \frac{2^n - 20 \cdot 8^n + 64 \cdot 32^n}{45} f_o\left(\frac{x}{2^n}\right) \\ &- \frac{40(2^n - 17 \cdot 8^n + 16 \cdot 32^n)}{45} f_o\left(\frac{x}{2^{n+1}}\right) + \frac{256(2^n - 5 \cdot 8^n + 4 \cdot 32^n)}{45} f_o\left(\frac{x}{2^{n+2}}\right) \end{aligned}$$

and

$$J'_n f(x)$$

$$\begin{aligned} &:= \left(\frac{1}{64^n} - \frac{5}{16^n} + \frac{4}{4^n}\right) \frac{f_e(2^{n+2}x)}{2880} - \left(\frac{20}{64^n} - \frac{340}{16^n} + \frac{320}{4^n}\right) \frac{f_e(2^{n+1}x)}{2880} \\ &+ \left(\frac{64}{64^n} - \frac{1280}{16^n} + \frac{4096}{4^n}\right) \frac{f_e(2^n x)}{2880} + \left(\frac{1}{32^n} - \frac{5}{8^n} + \frac{4}{2^n}\right) \frac{f_o(2^{n+2}x)}{720} \\ &- \left(\frac{10}{32^n} - \frac{170}{8^n} + \frac{160}{2^n}\right) \frac{f_o(2^{n+1}x)}{720} + \left(\frac{16}{32^n} - \frac{320}{8^n} + \frac{1024}{2^n}\right) \frac{f_o(2^n x)}{720} \end{aligned}$$

for all $x \in V$ and all nonnegative integers n . Then

$$(2.3) \quad \begin{aligned} J_n f(x) - J_{n+1} f(x) &= \left(\frac{4^n}{45} - \frac{20 \cdot 16^n}{45} + \frac{64 \cdot 64^n}{45}\right) \Delta f\left(\frac{x}{2^{n+3}}\right) \\ &+ \left(\frac{2^n}{45} - \frac{20 \cdot 8^n}{45} + \frac{64 \cdot 32^n}{45}\right) \Gamma f\left(\frac{x}{2^{n+3}}\right) \end{aligned}$$

$$(2.4) \quad \begin{aligned} J'_n f(x) - J'_{n+1} f(x) &= - \left(\frac{4}{4^{n+1}} - \frac{5}{16^{n+1}} + \frac{1}{64^{n+1}}\right) \frac{\Delta f(2^n x)}{2880} \\ &- \left(\frac{4}{2^{n+1}} - \frac{5}{8^{n+1}} + \frac{1}{32^{n+1}}\right) \frac{\Gamma f(2^n x)}{720} \end{aligned}$$

for all $x \in V$ and all nonnegative integers n .

Proof. From the equalities (2.1) and the definitions of $J_n f$ and $J'_n f$, we obtain the equalities

$$\begin{aligned} & J_n f(x) - J_{n+1} f(x) \\ &= \left(\frac{4^n}{45} - \frac{20 \cdot 16^n}{45} + \frac{64 \cdot 64^n}{45} \right) \Delta f \left(\frac{x}{2^{n+3}} \right) \\ &+ \left(\frac{2^n}{45} - \frac{20 \cdot 8^n}{45} + \frac{64 \cdot 32^n}{45} \right) \Gamma f \left(\frac{x}{2^{n+3}} \right) \end{aligned}$$

and similarly we have

$$\begin{aligned} & J'_n f(x) - J'_{n+1} f(x) \\ &= - \left(\frac{4}{4^{n+1}} - \frac{5}{16^{n+1}} + \frac{1}{64^{n+1}} \right) \frac{\Delta f(2^n x)}{2880} \\ &- \left(\frac{4}{2^{n+1}} - \frac{5}{8^{n+1}} + \frac{1}{32^{n+1}} \right) \frac{\Gamma f(2^n x)}{720} \end{aligned}$$

for all $x \in V$ and all nonnegative integers n . □

LEMMA 2.2. *If $f : V \rightarrow Y$ is a mapping such that*

$$Df(x, y) = 0$$

for all $x, y \in V$ with $f(0) = 0$, then

$$J_n f(x) = f(x) \quad \text{and} \quad J'_n f(x) = f(x)$$

for all $x \in V$ and all positive integers n .

Proof. If $f : V \rightarrow Y$ is a mapping such that

$$Df(x, y) = 0$$

for all $x, y \in V$ with $f(0) = 0$, then it follows from the definitions of $\Delta f(x)$ and $\Gamma f(x)$ that $\Delta f(x) = 0$ and $\Gamma f(x) = 0$ for all $x \in V$. Therefore, together with the equality $f(x) - J_n f(x) = \sum_{i=0}^{n-1} (J_i f(x) - J_{i+1} f(x))$ and the equality (2.3), we conclude that

$$J_n f(x) = f(x)$$

for all $x \in V$ and all positive integers n . In the same way we can easily show the equality $J'_n f(x) = f(x)$ for all $x \in V$. □

From Lemma 2.1 and Lemma 2.2, we can prove the following stability theorem.

THEOREM 2.3. Let $\varphi : V^2 \rightarrow [0, \infty)$ be a function such that

$$(2.5) \quad \sum_{n=0}^{\infty} 64^n \varphi\left(\frac{x}{2^n}, \frac{y}{2^n}\right) < \infty$$

for all $x, y \in V$. Suppose that $f : V \rightarrow Y$ is a mapping such that

$$(2.6) \quad \|Df(x, y)\| \leq \varphi(x, y)$$

for all $x, y \in V$. Then there exists a general sextic mapping F such that

$$(2.7) \quad \|f(x) - f(0) - F(x)\| \leq \sum_{n=0}^{\infty} \left[\left(\frac{4^n}{45} - \frac{20 \cdot 16^n}{45} + \frac{64 \cdot 64^n}{45} \right) \Phi\left(\frac{x}{2^{n+3}}\right) + \left(\frac{2^n}{45} - \frac{20 \cdot 8^n}{45} + \frac{64 \cdot 32^n}{45} \right) \Phi'\left(\frac{x}{2^{n+3}}\right) \right]$$

for all $x \in V$ and $F(0) = 0$, where $\varphi_e : V^2 \rightarrow [0, \infty)$ and $\Phi, \Phi' : V \rightarrow [0, \infty)$ are functions defined by

$$\begin{aligned} \varphi_e(x, y) &:= \frac{\varphi(x, y) + \varphi(-x, -y)}{2} \\ \Phi(x) &:= \varphi_e(-6x, 2x) + 8\varphi_e(-x, x) + 56\varphi_e(-2x, x) + 112\varphi_e(-3x, x) \\ \Phi'(x) &:= \varphi_e(-6x, 2x) + 6\varphi_e(-x, x) + 42\varphi_e(-2x, x) + 112\varphi_e(-3x, x). \end{aligned}$$

Proof. If \tilde{f} is a mapping defined by $\tilde{f}(x) = f(x) - f(0)$, then the mapping \tilde{f} satisfies the properties $D\tilde{f}(x, y) = Df(x, y)$ and $\tilde{f}(0) = 0$. By (2.1) and the definitions of Γf and Δf , we have

$$\|\Gamma\tilde{f}(x)\| \leq \Phi'(x) \quad \text{and} \quad \|\Delta\tilde{f}(x)\| \leq \Phi(x),$$

for all $x \in V$. Hence, from (2.3) and (2.5), we have

$$\begin{aligned} &\|J_n\tilde{f}(x) - J_{n+1}\tilde{f}(x)\| \\ &\leq \left(\frac{4^n}{45} - \frac{20 \cdot 16^n}{45} + \frac{64 \cdot 64^n}{45} \right) \Phi\left(\frac{x}{2^{n+3}}\right) \\ &\quad + \left(\frac{2^n}{45} - \frac{20 \cdot 8^n}{45} + \frac{64 \cdot 32^n}{45} \right) \Phi'\left(\frac{x}{2^{n+3}}\right) \end{aligned}$$

for all $x \in V$. So we obtain

$$(2.8) \quad \begin{aligned} \|J_n \tilde{f}(x) - J_{n+m} \tilde{f}(x)\| \leq & \sum_{i=n}^{n+m-1} \left[\left(\frac{4^i}{45} - \frac{20 \cdot 16^i}{45} + \frac{64 \cdot 64^i}{45} \right) \Phi \left(\frac{x}{2^{i+3}} \right) \right. \\ & \left. + \left(\frac{2^i}{45} - \frac{20 \cdot 8^i}{45} + \frac{64 \cdot 32^i}{45} \right) \Phi' \left(\frac{x}{2^{i+3}} \right) \right] \end{aligned}$$

for all $x \in V$ and $n, m \in \mathbb{N} \cup \{0\}$. It follows from (2.5) and (2.8) that the sequence $\{J_n \tilde{f}(x)\}$ is a Cauchy sequence for all $x \in V$. Since Y is complete, the sequence $\{J_n \tilde{f}(x)\}$ converges for all $x \in V$. Hence we can define a mapping $F : V \rightarrow Y$ by

$$F(x) := \lim_{n \rightarrow \infty} J_n \tilde{f}(x)$$

for all $x \in V$. Notice that $J_0 \tilde{f}(x) = f(x) - f(0)$ for all $x \in V$ and $F(0) = 0$ follows from $\tilde{f}(0) = 0$. Moreover, letting $n = 0$ and passing the limit $m \rightarrow \infty$ in (2.8) we get the inequality (2.7). From the definition of F , we easily get

$$\begin{aligned} \|DF(x, y)\| &= \lim_{n \rightarrow \infty} \|DJ_n \tilde{f}(x, y)\| \\ &\leq \lim_{n \rightarrow \infty} \left(\frac{64^{n+1}}{45} \varphi_e \left(\frac{x}{2^n}, \frac{y}{2^n} \right) + \frac{20 \cdot 64^{n+1}}{45} \varphi_e \left(\frac{x}{2^{n+1}}, \frac{y}{2^{n+1}} \right) \right. \\ &\quad + \frac{64^{n+1}}{45} \varphi_e \left(\frac{x}{2^{n+2}}, \frac{y}{2^{n+2}} \right) + \frac{64 \cdot 32^n}{45} \varphi_e \left(\frac{x}{2^n}, \frac{y}{2^n} \right) \\ &\quad \left. + \frac{640 \cdot 32^n}{45} \varphi_e \left(\frac{x}{2^{n+1}}, \frac{y}{2^{n+1}} \right) + \frac{1024 \cdot 32^n}{45} \varphi_e \left(\frac{x}{2^{n+2}}, \frac{y}{2^{n+2}} \right) \right) = 0, \end{aligned}$$

for all $x, y \in V$. To prove the uniqueness of F , let $F' : V \rightarrow Y$ be another general sextic mapping satisfying (2.7) and $F'(0) = 0$. Instead of the condition (2.7), it is sufficient to show that there is a unique mapping satisfying the simpler condition

$$\|\tilde{f}(x) - F(x)\| \leq \sum_{i=0}^{\infty} \left(64^{i+1} \Phi \left(\frac{x}{2^{i+3}} \right) + 64^{i+1} \Phi' \left(\frac{x}{2^{i+3}} \right) \right),$$

for all $x \in V$. By Lemma 2.2, the equality $F'(x) = J_n F'(x)$ holds for all $n \in \mathbb{N}$. So we have

$$\begin{aligned} & \|J_n \tilde{f}(x) - F'(x)\| = \|J_n \tilde{f}(x) - J_n F'(x)\| \\ & \leq 64^{n+1} \left\| \tilde{f}_e\left(\frac{x}{2^n}\right) - F'_e\left(\frac{x}{2^n}\right) \right\| + 64^{n+1} \left\| \tilde{f}_e\left(\frac{x}{2^{n+1}}\right) - F'_e\left(\frac{x}{2^{n+1}}\right) \right\| \\ & + 64^{n+2} \left\| \tilde{f}_e\left(\frac{x}{2^{n+2}}\right) - F'_e\left(\frac{x}{2^{n+2}}\right) \right\| + 64^{n+1} \left\| \tilde{f}_o\left(\frac{x}{2^n}\right) - F'_o\left(\frac{x}{2^n}\right) \right\| \\ & + 64^{n+1} \left\| \tilde{f}_o\left(\frac{x}{2^{n+1}}\right) - F'_o\left(\frac{x}{2^{n+1}}\right) \right\| + 64^{n+1} \left\| \tilde{f}_o\left(\frac{x}{2^{n+2}}\right) - F'_o\left(\frac{x}{2^{n+2}}\right) \right\| \\ & \leq \sum_{i=n+3}^{\infty} 64^i \Phi\left(\frac{x}{2^i}\right) + \sum_{i=n+3}^{\infty} 64^i \Phi'\left(\frac{x}{2^i}\right), \end{aligned}$$

for all $x \in V$ and all positive integer n . Taking the limit in the above inequality as $n \rightarrow \infty$, we obtain the equality $F'(x) = \lim_{n \rightarrow \infty} J_n \tilde{f}(x)$ for all $x \in V$, which means that $F(x) = F'(x)$ for all $x \in V$. \square

THEOREM 2.4. *Let $\varphi : V^2 \rightarrow [0, \infty)$ be a function such that*

$$(2.9) \quad \sum_{n=0}^{\infty} 2^{-n} \varphi(2^n x, 2^n y) < \infty$$

for all $x, y \in V$. Suppose that $f : V \rightarrow Y$ is a mapping such that

$$(2.10) \quad \|Df(x, y)\| \leq \varphi(x, y)$$

for all $x, y \in V$. Then there exists a general sextic mapping F such that

$$(2.11) \quad \begin{aligned} \|f(x) - f(0) - F(x)\| & \leq \sum_{n=0}^{\infty} \left[\left(\frac{4}{4^{n+1}} - \frac{5}{16^{n+1}} + \frac{1}{64^{n+1}} \right) \frac{\Phi(2^n x)}{2880} \right. \\ & \quad \left. + \left(\frac{4}{2^{n+1}} - \frac{5}{8^{n+1}} + \frac{1}{32^{n+1}} \right) \frac{\Phi'(2^n x)}{720} \right], \end{aligned}$$

for all $x \in V$ and $F(0) = 0$, where φ_e, Φ, Φ' are functions defined as Theorem 2.3.

Proof. If \tilde{f} is a mapping defined by $\tilde{f}(x) = f(x) - f(0)$, then the mapping \tilde{f} satisfies the properties $D\tilde{f}(x, y) = Df(x, y)$ and $\tilde{f}(0) = 0$. By the definitions of Γf and Δf , we have

$$\|\Gamma \tilde{f}(x)\| \leq \Phi'(x) \quad \text{and} \quad \|\Delta \tilde{f}(x)\| \leq \Phi(x),$$

for all $x \in V$. So it follows from (2.4) and (2.10) that

$$\begin{aligned} \|J'_n \tilde{f}(x) - J'_{n+1} \tilde{f}(x)\| &\leq \left(\frac{4}{4^{n+1}} - \frac{5}{16^{n+1}} + \frac{1}{64^{n+1}} \right) \left\| \frac{\Delta f(2^n x)}{2880} \right\| \\ &\quad + \left(\frac{4}{2^{n+1}} - \frac{5}{8^{n+1}} + \frac{1}{32^{n+1}} \right) \left\| \frac{\Gamma f(2^n x)}{720} \right\| \\ &\leq \left(\frac{4}{4^{n+1}} - \frac{5}{16^{n+1}} + \frac{1}{64^{n+1}} \right) \frac{\Phi(2^n x)}{2880} \\ &\quad + \left(\frac{4}{2^{n+1}} - \frac{5}{8^{n+1}} + \frac{1}{32^{n+1}} \right) \frac{\Phi'(2^n x)}{720} \end{aligned}$$

for all $x \in V$. Together with the equality

$$J'_n \tilde{f}(x) - J'_{n+m} \tilde{f}(x) = \sum_{i=n}^{n+m-1} (J'_i \tilde{f}(x) - J'_{i+1} \tilde{f}(x))$$

for all $x \in V$, we obtain that

(2.12)

$$\begin{aligned} \|J'_n \tilde{f}(x) - J'_{n+m} \tilde{f}(x)\| &\leq \sum_{i=n}^{n+m-1} \left[\left(\frac{4}{4^{i+1}} - \frac{5}{16^{i+1}} + \frac{1}{64^{i+1}} \right) \frac{\Phi(2^i x)}{2880} \right. \\ &\quad \left. + \left(\frac{4}{2^{i+1}} - \frac{5}{8^{i+1}} + \frac{1}{32^{i+1}} \right) \frac{\Phi'(2^i x)}{720} \right] \end{aligned}$$

for all $x \in V$ and $n, m \in \mathbb{N} \cup \{0\}$. It follows from (2.9) and (2.12) that the sequence $\{J'_n \tilde{f}(x)\}$ is a Cauchy sequence for all $x \in V$. Since Y is complete, the sequence $\{J'_n \tilde{f}(x)\}$ converges for all $x \in V$. Hence we can define a mapping $F : V \rightarrow Y$ by

$$F(x) := \lim_{n \rightarrow \infty} J'_n \tilde{f}(x)$$

for all $x \in V$. Note that $F(0) = 0$ follows from $\tilde{f}(0) = 0$. Notice that $J'_0 \tilde{f}(x) = f(x) - f(0)$ for all $x \in V$. Moreover, letting $n = 0$ and passing the limit $n \rightarrow \infty$ in (2.12) we get the inequality (2.11). From

the definition of F , we easily get

$$\begin{aligned} \|DF(x, y)\| &= \lim_{n \rightarrow \infty} \|DJ'_n f(x, y)\| \\ &\leq \lim_{n \rightarrow \infty} \left(\frac{2\varphi_e(2^n x, 2^n y) + \varphi_e(2^{n+1} x, 2^{n+1} y) + \varphi_e(2^{n+2} x, 2^{n+2} y)}{4^n} \right. \\ &\quad \left. + \frac{2\varphi_e(2^n x, 2^n y) + \varphi_e(2^{n+1} x, 2^{n+1} y) + \varphi_e(2^{n+2} x, 2^{n+2} y)}{2^n} \right) \\ &= 0 \end{aligned}$$

for all $x, y \in V$. To prove the uniqueness of F , let $F' : V \rightarrow Y$ be another general sextic mapping satisfying (2.11) and $F'(0) = 0$. Instead of the condition (2.11), it is sufficient to show that there is a unique mapping satisfying the simpler condition

$$\|\tilde{f}(x) - F(x)\| \leq \sum_{i=0}^{\infty} \frac{\Phi(2^i x) + \Phi'(2^i x)}{2^i}$$

for all $x \in V$. By Lemma 2.2, the equality $F'(x) = J'_n F'(x)$ holds for all $n \in \mathbb{N}$. So we have

$$\begin{aligned} \|J'_n \tilde{f}(x) - F'(x)\| &= \|J'_n \tilde{f}(x) - J'_n F'(x)\| \\ &\leq \frac{2}{4^n} \|(\tilde{f}_e - F'_e)(2^n x)\| + \frac{1}{4^{n+1}} \|(\tilde{f}_e - F'_e)(2^{n+1} x)\| + \frac{1}{4^{n+2}} \|(\tilde{f}_e - F'_e)(2^{n+2} x)\| \\ &\quad + \frac{2}{2^n} \|(\tilde{f}_o - F'_o)(2^n x)\| + \frac{1}{2^{n+2}} \|(\tilde{f}_o - F'_o)(2^{n+1} x)\| + \frac{1}{2^{n+2}} \|(\tilde{f}_o - F'_o)(2^{n+2} x)\| \\ &\leq 4 \sum_{i=0}^{\infty} \frac{\Phi(2^{n+i} x) + \Phi'(2^{n+i} x)}{2^{n+i}} + 2 \sum_{i=0}^{\infty} \frac{\Phi(2^{n+i+1} x) + \Phi'(2^{n+i+1} x)}{2^{n+i+1}} \\ &\quad + 2 \sum_{i=0}^{\infty} \frac{\Phi(2^{n+i+2} x) + \Phi'(2^{n+i+2} x)}{2^{n+i+2}} \\ &\leq 8 \sum_{i=n}^{\infty} \frac{\Phi(2^i x) + \Phi'(2^i x)}{2^i} \end{aligned}$$

for all $x \in V$ and all positive integer n . Taking the limit in the above inequality as $n \rightarrow \infty$, we obtain the equality $F'(x) = \lim_{n \rightarrow \infty} J'_n \tilde{f}(x)$ for all $x \in V$, which means that $F(x) = F'(x)$ for all $x \in V$. \square

References

- [1] S. Alshybani, S. M. Vaezpour and R. Saadati, *Generalized Hyers-Ulam stability of sextic functional equation in random normed spaces*, J. Comput. Anal. Appl., **24** (2018), no. 2, 370-381.
- [2] S. Alshybani, S. M. Vaezpour and R. Saadati, *Stability of the sextic functional equation in various spaces*, J. Inequal. Spec. Funct., **9** (2018), no. 4, 8-27.
- [3] J. Baker, *A general functional equation and its stability*, Proc. Natl. Acad. Sci., **133** (2005), no. 6, 1657-1664.
- [4] Y. J. Cho, M. B. Ghaemi, M. Choubin and M. E. Gordji, *On the Hyers-Ulam stability of sextic functional equations in β -homogeneous probabilistic modular spaces*, Math. Inequal. Appl., **16** (2013), no. 4, 1097-1114.
- [5] I. I. EL-Fassi, *New stability results for the radical sextic functional equation related to quadratic mappings in $(2, \beta)$ -Banach spaces*, J. Fixed Point Theory Appl., **20** (2018), 138, 1-17.
- [6] Z. Gajda, *On stability of additive mappings*, Int. J. Math. Math. Sci., **14** (1991), no. 3, 431-434.
- [7] P. Gävruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl., **184** (1994), 431-436.
- [8] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Natl. Acad. Sci. USA, **27** (1941), 222-224.
- [9] G. Isac and T. M. Rassias, *On the Hyers-Ulam stability of ψ -additive mappings*, J. Approx. Theory, **72** (1993), 131-137.
- [10] K.-W. Jun and H.-M. Kim, *On the Hyers-Ulam-Rassias stability of a general cubic functional equation*, Math. Inequal. Appl., **6** (2003), 289-302.
- [11] S.-M. Jung, *On the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl., **204** (1996), 221-226.
- [12] J. Roh, Y. H. Lee and S. -M. Jung *The Stability of a General Sextic Functional Equation by Fixed Point Theory*, J. Funct. Spaces, **2020** (2020), 1-8.
- [13] Y.-H. Lee, *On the generalized Hyers-Ulam stability of the generalized polynomial function of degree 3*, Tamsui Oxf. J. Math. Sci., **24** (2008), no. 4, 429-444.
- [14] Y.-H. Lee, *On the Hyers-Ulam-Rassias stability of the generalized polynomial function of degree 2*, J. Chungcheong Math. Soc., **22** (2009), no.2, 201-209.
- [15] Y.-H. Lee, *Stability of a monomial functional equation on a restricted domain*, Mathematics, **5** (2017), 53.
- [16] Y.-H. Lee, *On the Hyers-Ulam-Rassias stability of a general quartic functional equation*, East Asian Math. J., **35** (2019), no. 3, 351-356.
- [17] Y.-H. Lee, *On the Hyers-Ulam-Rassias stability of a general quintic functional equation and a general sextic functional equation*, Mathematics, **7** (2019), 510.
- [18] Y.-H. Lee and K. W. Jun, *On the stability of approximately additive mappings*, Proc. Amer. Math. Soc., (2000), 1361-1369.
- [19] A. Najati and C. Park, *Fixed Points and Stability of a Generalized Quadratic Functional Equation* J. Inequal. Appl., **2009**(2009), Article ID 193035, 19 pages.
- [20] Th. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc., **72** (1978), 297-300.
- [21] T. Xu, J. Rassias, M. Rassias and W. Xu, *A fixed point approach to the stability of quintic and sextic functional equations in quasi- β -normed spaces*, J. Inequal. Appl., **2010** (2010), Article ID 423231, 23 pages.

- [22] T. Xu, J. Rassias, M. Rassias and W. Xu, *Stability of quintic and sextic functional equations in non-archimedean fuzzy normed spaces*, Eighth International Conference on Fuzzy Systems and Knowledge Discovery, **2011**.
- [23] K. Ravi and S. Sabarinathan, *Generalized Hyers-Ulam stability of a sextic functional equation in paranormed spaces*, International Journal of Mathematics Trends and Technology, **9** (2014), no. 1, 61-69.
- [24] S. Ostadbashi and M. Soleimaninia, *On the stability of the orthogonal pexiderized quartic functional equations*, Nonlinear Funct. Anal. Appl., **20** (2015), no. 4, 539-549.
- [25] S.M. Ulam, *A Collection of Mathematical Problems*, Interscience, New York, 1960.

*

Department of Mathematics
Chungnam National University
Daejeon 34134, Republic of Korea
E-mail: ischang@math.chungnam.ac.kr

**

Department of Mathematics Education
Gongju National University of Education
Gongju 32553, Republic of Korea
E-mail: yanghi2@hanmail.net

Ilsong College of Liberal Arts
Hallym University
Chuncheon 24252, Republic of Korea
E-mail: joroh@hallym.ac.kr